ON THE PERIODIC POINTS OF A TYPICAL CONTINUOUS FUNCTION

K. SIMON

(Communicated by R. Daniel Mauldin)

ABSTRACT. Let n and k be arbitrary natural numbers. We prove that for a typical continuous function f , every neighborhood of any periodic point of f with period n contains periodic points of f with period $n \cdot k$.

The purpose of this paper is to examine the structure of the set of periodic points of typical continuous functions. We shall denote by C the set of continuous functions mapping the interval $[0, 1]$ into itself. This set becomes a metric space with the supremum metric. By the term "typical continuous function" we mean that the set of all functions having the property under consideration is a residual subset of the metric space C .

In [1] S. J. Agronsky, A. M. Bruckner, and M. Laczkovich proved that for a typical continuous function f any neighborhood of a periodic point contains periodic points of arbitrarily large periods. In this paper we prove that this result is true if "arbitrarily large periods" is replaced by "period $k \cdot n$ ($k = 1, 2, \dots$)" (Theorem 1). As a consequence, for a typical continuous f we have that P_f^n , the set of periodic points with period n , is uncountable, dense in itself, but not closed. Also, P_f^n is a residual subset of $\text{Fix}(f^n)$.

In [1] it is proved that, for a typical continuous f , $\text{Fix}(f^n)$ is nowhere dense and perfect. In this paper we prove (Theorem 2) that $\text{Fix}(f^n)$ is bilaterally strongly Φ -porous. For $f \in C$ and $n \in \mathbb{N}$ we define $f^n(x)$ by $f^1(x) = f(x)$ and $f^n(x) = f(f^{n-1}(x))$. P_f^n denotes the set of periodic points of f with period n . We denote $\text{Fix}(f^n) = \{x : f^n(x) = x\}$; then $P_f^n = \text{Fix}(f^n) \setminus \bigcup_{k < n} \text{Fix}(f^k)$. In particular P_f^n is a relatively open subset of $\text{Fix}(f^n)$. For $f \in C$ and $\varepsilon > 0$ denote

$$B(f, \varepsilon) = \{g \in C : \|f - g\| < \varepsilon\}.$$

Theorem 1. *Let $n, k \in \mathbb{N}$ be arbitrary. Then for a typical continuous function f , every neighborhood of any periodic point of f with period n contains periodic points of f with period $n \cdot k$.*

Received by the editors August 3, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 54H20; Secondary 26A18.

©1989 American Mathematical Society
0002-9939/89 \$1.00 + \$.25 per page

Proof. Fix $n \in \mathbb{N}$. Let $A_0 = \{f \in C: \text{Fix}(f^n) \text{ nowhere dense}\}$ and $A_i = \{f \in C: \text{every neighborhood of any periodic point of } f \text{ with period } i \text{ contains, for every } m \in \mathbb{N}, \text{ periodic points of } f \text{ with period } m \cdot i\} \ (i = 1, 2, \dots)$.

We prove by induction that each A_i is residual. In [1] it is proved that the set A_0 is a residual subset of C . Assume that $\bigcap_{0 \leq i < n} A_i$ is a residual subset of C . We shall prove that A_n is also residual. If $f \in \bigcap_{0 \leq i < n} A_i$ and I is an arbitrary open interval then $P_f^n \cap I = \emptyset$ implies $\text{Fix}(f^n) \cap I = \emptyset$. For $k \in \mathbb{N}$ and $\varepsilon > 0$ we define the set of Q_k^ε by

$$Q_k^\varepsilon = \{f \in C: \text{for any } x_0 \in P_f^n \text{ there is } x \in P_f^{k \cdot n} \text{ such that } |x - x_0| < \varepsilon\}.$$

It will be sufficient to show that each Q_k^ε contains a set which is open and dense. Let $f \in \bigcap_{0 \leq i < n} A_i$ and let $\varepsilon' > 0$ be arbitrary. We prove that there exist $g \in B(f, \varepsilon')$ and $\eta > 0$ such that $B(g, \eta) \subset Q_k^\varepsilon$. Let $\{a_i\}_{i=0}^l \subset [0, 1] \setminus \text{Fix}(f^n)$ be a finite sequence such that $a_0 = 0$, $a_l = 1$ and $0 < a_{i+1} - a_i < \varepsilon$ for every $0 \leq i < l$. Put $J_i = [a_{i-1}, a_i]$ ($i = 1, \dots, l$). Choose a finite sequence $\{b_j\}_{j=1}^z \subset P_f^n$ as follows: if $J_i \cap P_f^n \neq \emptyset$ ($i = 1, \dots, l$) then there should exist an $x_0 \in J_i \cap P_f^n$ such that the sequence $\{b_j\}_{j=1}^z$ contains the orbit of x_0 . Let $z' = z/n$. We can choose $b_1, b_2, \dots, b_z$ so that they are contained in different orbits. Choose $\eta_1 \in \mathbb{R}$ so that

- (i) $\eta_1 > 0$,
- (ii) $\eta_1 < \frac{1}{3} \cdot \min_{\substack{0 \leq i_1 < i_2 \leq l \\ 0 \leq j_1 < j_2 \leq z \\ 0 \leq i \leq l, 0 \leq j \leq z}} \{|a_{i_1} - a_{i_2}|, |b_{j_1} - b_{j_2}|, |a_i - b_j|\}$,
- (iii) $\eta_1 < \varepsilon'/3$, and
- (iv) if for some i $J_i \cap P_f^n = \emptyset$ and $\|g - f\| < 3\eta_1$ then $J_i \cap P_g^n = \emptyset$.

This is possible since $f \in \bigcap_{i < n} A_i$. Let $0 < u < \eta_1$ be such that if $|x - y| < u$ then $|f(x) - f(y)| < \eta_1$. Then $[b_i - u/2, b_i + u/2] \cap [b_j - u/2, b_j + u/2] = \emptyset$ for every $1 \leq i < j \leq z$.

We can now define the function $g \in C$. First we define g on

$$\bigcup_{i=1}^n [f^i(b_m) - u/2, f^i(b_m) + u/2] \quad (m = 1, \dots, z')$$

using the following procedure. Let $1 \leq m \leq z'$ be fixed but arbitrary and take $x_0 = b_m$. Put $I_j = [f^{j-1}(x_0) - u/2, f^{j-1}(x_0) + u/2]$. Let us define g on the set $\bigcup_{j=1}^n I_j$ so that the graph of g on this set will be included in the squares $E_j = I_j \times I_{j+1}$ (see Figure 1). If $j > 1$ then let the graph of g in E_j be the increasing diagonal of E_j . (See Figure 1.) Let $x_1, \dots, x_k, x_{k+1} = x_1$ be distinct points in $(0, u)$ and let the point W_i be defined by $W_i = (x_i, x_{i+1})$ ($i = 1, \dots, k$); then $W_i \in L = [0, u] \times [0, u]$. Let $v = \min_{i \neq j} \frac{1}{3} |x_i - x_j|$ and define

$$B_i = [x_i - \frac{v}{2}, x_i + \frac{v}{2}] \times [x_{i+1} - \frac{v}{2}, x_{i+1} + \frac{v}{2}].$$

$$n = 4, k = 3$$

FIGURE 1

Let $h: [0, u] \rightarrow [0, u]$ be piecewise linear and continuous such that the graph of h in each B_i is the diagonal of B_i , if $1 \leq i < k$ then h is increasing in B_i but h is decreasing in B_k . (See Figure 2.)

It is easy to see that if $\eta = v/10$ and $q \in B(h, \eta)$ then q has periodic points of period k in the interval $(x_i - v/2, x_i + v/2)$. Now we define the graph of g on I_1 by translating the graph of h into E_1 (using the translation that maps L onto E_1). Thus we have defined g on $\bigcup_{i=1}^2 [b_i - u/2, b_i + u/2]$. Put $g(a_i) = f(a_i)$ ($i = 0, 1, \dots, l$). If $J_i \cap P_f^n = \emptyset$ then let $g|_{J_i} = f|_{J_i}$. For those x 's for which g has not yet been defined, let us define g so that $\|f - g\| < \eta_1$; by the definition of u this is possible. Now if $P_f^n \cap J_i = \emptyset$ then $P_g^n \cap J_i = \emptyset$ or else $P_g^{n \cdot k} \cap J_i \neq \emptyset$. Thus $g \in B(f, \varepsilon') \cap Q_k^\varepsilon$ and it is easy to see that if $\eta = v/10$ then $B(g, \eta) \subset B(f, \varepsilon') \cap Q_k^\varepsilon$. It follows that Q_k^ε contains

FIGURE 2

an open dense set, and since $\bigcap_{\epsilon, k} Q_k^\epsilon \subset A$ we have that A is a residual subset of C . This completes the proof of the theorem.

We show that, for a typical continuous f , P_f^k is a dense relatively open subset of $\text{Fix}(f^k)$ from which it will follow that, P_f^k is a residual subset of $\text{Fix}(f^k)$ and P_f^k is uncountable.

Corollary 1. *For a typical continuous f , P_f^k is a dense relatively open subset of $\text{Fix}(f^k)$.*

Proof. If $x \in \text{Fix}(f^k)$ then there is $q|_k$ such that $x \in P_f^q$. It follows from Theorem 1 that any neighborhood of x contains periodic points with period $q \cdot k/q$. As we have seen in the introduction, P_f^k is relatively open in $\text{Fix}(f^k)$.

Corollary 2. *For a typical continuous f , P_f^k is not closed for every $k > 1$.*

Proof. It follows from Theorem 1 that for every $x \in \text{Fix}(f)$, $x \in \text{cl}(P_f^k)$ and $x \notin P_f^k$.

Corollary 3. *For a typical continuous f , P_f^k is dense in itself.*

Proof. Since, for a typical f , $\text{Fix}(f^k)$ is dense in itself [1], the statement follows from Corollary 1.

Remark. For a typical f there is $x \in \text{Fix}(f^n)$ such that in a suitable neighborhood of x , the periods of the periodic points are all multiples of n . Moreover, this is true for the points of a residual subset of $\text{Fix}(f^n)$. Indeed, let $f \in C$ and $I_0 \subset [0, 1]$ be an open interval and let $I_k = f^k(I_0)$. Define $P_k^n(f) = \{I_0: \text{ for every } 0 \leq 1 \neq j \leq k-1, I_i \cap I_j = \emptyset \text{ and } I_n \subset I_0 \text{ and } \min\{|I_0|, \dots, |I_n|\} < 1/k\}$. (See [1, Proposition 1].)

In [1] the authors proved that for a typical continuous function f , $\text{Fix}(f^n) \cap \bigcap_{k=1}^{\infty} G_k^n$ is a residual subset of $\text{Fix}(f^n)$. Put $B_f^n = \{x \in P_f^n: \text{ there is } V \text{ neighborhood of } x \text{ such that for any periodic point } y \in V \text{ there is } k \text{ such that period of } y \text{ equal to } n \cdot k\}$. It is easy to see $P_f^n \cap \bigcap_{k=1}^{\infty} G_k^n \subset B_f^n$ and therefore B_f^n is a residual subset of $\text{Fix}(f^n)$.

In [1] it is proved that for a typical continuous f the set $\text{Fix}(f^k)$ is nowhere dense and perfect. In Theorem 2 we shall prove that for every porosity pre-measure (i.e., a continuous map $\Phi: (0, 1] \rightarrow (0, 1]$), $\text{Fix}(f^k)$ is bilaterally strongly Φ -porous for a typical continuous f . This means that for every $x \in \text{Fix}(f^k)$ there are sequences of intervals $\{I_n\}_{n=1}^{\infty}$ and $\{J_n\}_{n=1}^{\infty}$ such that $I_n \subset (x - 1/n, x) \setminus \text{Fix}(f^k)$, $J_n \subset (x, x + 1/n) \setminus \text{Fix}(f^k)$ and

$$\lim_{n \rightarrow \infty} \frac{\text{dist}(x, I_n)}{\Phi(|I_n|)} = \lim_{n \rightarrow \infty} \frac{\text{dist}(x, J_n)}{\Phi(|J_n|)} = 0.$$

We shall apply the method used by P. Humke and M. Laczkovich in [2] to this problem.

The pair of sequences (α, β) is said to be proper if $\alpha = \{\alpha_n\}_{n=1}^{\infty}$, $\beta = \{\beta_n\}_{n=1}^{\infty}$ and $\beta_n \rightarrow 0$ and $0 < \alpha_n < \beta_n$. If (α, β) is a proper pair of sequences then the sequence $x = \{x_n\}_{n=1}^{\infty}$ is called an (α, β) -sequence if $x_n \rightarrow x_0$ and $\alpha_n \leq x_n - x_0 \leq \beta_n$ for each $n \in \mathbb{N}$. Let the natural number $l > 1$ and the proper pair (α, β) be fixed. (See [2, p. 245].) Put

$B_N = \{f \in C: \text{ there is } \{x_i\}_{i=1}^{\infty}(\alpha, \beta)\text{-sequence such that } f^l(x_i) = x_i \text{ if } i \geq N\}$.

Lemma. B_N is nowhere dense.

Proof. First we shall prove that B_N is closed. Let $\{f_k\}_{k=1}^{\infty} \subset B_N$ and $f_k \rightarrow f$ in C . Thus, for every fixed k there is a sequence $x^k = \{x_n^k\}_{n=1}^{\infty}$ converging to x_0^k such that x^k is an (α, β) -sequence and $f_k^l(x_i^k) = x_i^k$ if $i \geq N$. Using a diagonal procedure we obtain a subsequence $\{k_i\}_{i=1}^{\infty}$ such that for every n there is x_n^* such that $\{x_n^{k_i}\}_{i=1}^{\infty} \rightarrow x_n^*$ and $x^* = \{x_n^*\}_{n=1}^{\infty}$ is a (α, β) -sequence. As $\{f_k\}_{k=1}^{\infty} \rightarrow f$ in C and since for $n > N$ $f_{k_i}^l(x_n^{k_i}) = x_n^{k_i}$ it follows that $f^l(x_n^*) = x_n^*$ if $n > N$.

If p is a polynomial then $\{x: p^l(x) = x\}$ is finite and hence $p \notin B_N$. Since the set of polynomials is dense in C , the complement of B_N is dense and, as B_N is closed, B_N is nowhere dense.

Let $\Phi: (0, 1] \rightarrow (0, 1]$ be continuous arbitrary fixed.

Theorem 2. *For a typical continuous f , $\text{Fix}(f^l)$ is bilaterally strongly Φ -porous set.*

Proof. Let (α, β) be such that $\alpha_n = \beta_{n+1}$, $\beta_n < 1/n$ and $\Phi(\beta_n - \beta_{n+1}) \geq n\beta_{n+1}$. Let $M = \{x: f^l(x) = x\}$. If $f \in \bigcup_n B_n$ then for each $x \in M$ there is a sequence $\{n_i\} \rightarrow \infty$ such that $M \cap [x + \alpha_{n_i}, x + \beta_{n_i}] = \emptyset$. Let $J_i = [x + \alpha_{n_i}, x + \beta_{n_i}]$. Thus $J_i \subset (x, x + 1/i) \setminus M$ and

$$\text{dist}(x, J_i) = \alpha_{n_i} = \beta_{n_i+1} < \frac{\Phi(\beta_{n_i} - \beta_{n_i+1})}{n_i} = \frac{\Phi(|J_i|)}{n_i}$$

and hence

$$\lim_{n \rightarrow \infty} \frac{\text{dist}(x, J_i)}{\Phi(|J_i|)} = 0.$$

Using a similar procedure we can find a sequence $\{I_i\}_{i=1}^\infty$ such that

$$I_i \subset (x - 1/i, x) \setminus M \quad \text{and} \quad \lim_{i \rightarrow \infty} \frac{\text{dist}(x, I_i)}{\Phi(|I_i|)} = 0.$$

This completes the proof of the theorem.

ACKNOWLEDGMENT

The author would like to thank Miklós Laczkovich for his useful advice.

REFERENCES

1. S. J. Agronsky, A. M. Bruckner and M. Laczkovich, *Dynamics of typical continuous functions*, J. London Math. Soc. (2) (to appear).
2. P. Humke and M. Laczkovich, *Typical continuous functions are virtually nonmonotone*, Proc. Amer. Math. Soc. **92** (1985), 244–248.

DEPARTMENT OF MATHEMATICS, TECHNICAL UNIVERSITY, MISKOLC, HUNGARY