GABRIEL AND KRULL DIMENSIONS OF MODULES OVER RINGS GRADED BY FINITE GROUPS

PIOTR GRZESZCZUK AND EDMUND R. PUCZYŁOWSKI

(Communicated by Donald S. Passman)

ABSTRACT. Let R be a ring graded by a finite group G with the identity component R_e and let M be a left R -module. It is proved that $\text{Gdim}_R M = \text{Gdim}_{R_e} M$, $\text{Kdim}_R M = \text{Kdim}_{R_e} M$ and $\text{Ndim}_R M = \text{Ndim}_{R_e} M$, where Gdim , Kdim and Ndim denote, respectively, Gabriel, Krull and dual Krull dimensions. The proofs are based on the use of lattice theory, a method which also gives alternative proofs of known results about normalizing extensions.

INTRODUCTION

In [10] it was proved that a module over a ring R graded by a finite group G satisfies ACC on R -submodules if and only if it satisfies the condition on R_e -submodules, where R_e is the identity component of R . The corresponding question for DCC is, like the analogous one concerning normalizing extensions (cf. [2, 5, 12]), more difficult and seems that rather no direct extension of arguments of [10] or those used for normalizing extensions could provide such a result. In this paper, following the experience of [5, 9], we approach the problem via lattice theory. It appears that on a level of lattices differences between ACC and DCC disappear and the idea of [10] can be employed. It allows us not only to handle the DCC case but also prove that none of Gabriel, Krull and dual Krull dimensions of an R -module changes when it is considered over R or R_e .

1. PRELIMINARIES

Throughout this paper R is a ring with unity graded by a finite group G , i.e., $R = \bigoplus_{g \in G} R_g$ for additive subgroups R_g , satisfying $R_g R_h \subseteq R_{gh}$ for all $g, h \in G$. Obviously R_e is a subring of R and $1 \in R_e$. All considered modules will be left unital modules and $N \leq M$ (or $N \leq_s M$ if we need to specify the ring of scalars) means that N is a submodule of M . For proper submodules we write $N < M$.

Received by the editors March 25, 1987 and, in revised form, December 14, 1987.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 16A03, 16A55; Secondary 06C05.

©1989 American Mathematical Society
 0002-9939/89 \$1.00 + \$.25 per page

In this paper we use some lattices and their sublattices. All those are modular with 0 and 1 (we consider only sublattices containing 0 and 1 of the original lattice). Given a module M we write $L(M)$ for the lattice of submodules of M . We denote by L^0 the dual lattice to the lattice L , i.e., the lattice having the same elements as L but carrying the reverse ordering. For all elements $a, b \in L$ with $a \leq b$, $[a, b]$ denotes the interval $\{x \in L \mid a \leq x \leq b\}$. If θ is a congruence on L then $\theta(a)$ denotes the congruence class determined by a . A lattice isomorphism will be denoted by $\approx$. All undefined terms and used results on lattices can be found in [1].

An ideal I of a lattice L is called essential if for every nonzero ideal J of L we have $I \cap J \neq 0$.

Lemma 1. *Let $\{s_g \mid g \in G\}$, where G is a finite group, be a set of order preserving maps of a lattice L into itself such that $s_e = \text{id}$ and for every $g, h \in G$ and $x \in L$, $s_g s_h(x) \leq s_{gh}(x)$. If $f: L \rightarrow L'$ is a homomorphism of L onto a nonzero lattice L' and I is an essential ideal of L' , then $f(\bigcap_{g \in G} s_g^{-1} f^{-1}(I)) \neq 0$.*

Proof. For every subset X of G containing e , let $I_X = \bigcap_{g \in X} s_g^{-1} f^{-1}(I)$. Obviously $f(I_{\{e\}}) = I$, so $f(I_{\{e\}}) \neq 0$. Let X be a subset of maximal cardinality such that $e \in X$ and $f(I_X) \neq 0$. If there is $h \in G \setminus X$ then by maximality of $|X|$, $f(s_h(a)) \neq 0$ for some $a \in I_X$. Thus, since the ideal I is essential, $I \cap [0, f(s_h(a))] \neq 0$. Hence there is $b \in L$ with $b \leq s_h(a)$ and $0 \neq f(b) \in I$. Since $s_{gh^{-1}}$ preserves the order of L , for every $g \in X$, $s_{gh^{-1}}(b) \leq s_{gh^{-1}}s_h(a)$. Now by the assumption, $s_{gh^{-1}}(b) \leq s_{gh^{-1}}s_h(a) \leq s_g(a)$. Since $a \in I_X$ and $g \in X$, $f(s_g(a)) \in I$. Consequently, $f(s_{gh^{-1}}(b)) \in I$. Thus $b \in I_{Xh^{-1} \cup \{e\}}$ and since $f(b) \neq 0$, we have $f(I_{Xh^{-1} \cup \{e\}}) \neq 0$. This contradicts maximality of $|X|$, so $X = G$ and the lemma follows.

A sublattice K of a lattice L is said to be dense in L if for each element k in $K \setminus \{1\}$, every essential ideal I of the lattice $[k, 1]$ contains an element $k_1 \in K$, $k_1 \neq k$.

The following is the key proposition of this paper. A similar proposition is true for normalizing extensions, as proved in §4.

Proposition 1. *If M is an R -module and f is a homomorphism of $L_{(R_c)}M$ onto a nonzero lattice L , then $f(L_{(R_c)}M) (f(L^0_{(R_c)}M))$ is a dense sublattice of $L (L^0)$. In particular, $L_{(R_c)}M (L^0_{(R_c)}M)$ is a dense sublattice of $L_{(R_c)}M (L^0_{(R_c)}M)$.*

Proof. For every $N \leq_R M$ there is a natural isomorphism $i: L_{(R_c)}(M/N) \rightarrow [N, M] \subseteq L_{(R_c)}M$ such that $i(L_{(R_c)}(M/N)) = [N, M] \cap L_{(R_c)}M$. Now, f maps $[N, M]$ onto $[f(N), 1]$. Thus for every $k \in f(L_{(R_c)}M)$ there is an R -module K and a homomorphism $\bar{f}$ of $L_{(R_c)}K$ onto $[k, 1] \subseteq L$ such that $\bar{f}(L_{(R_c)}K) = f(L_{(R_c)}M) \cap [k, 1]$. Hence to prove that $f(L_{(R_c)}M)$ is a dense sublattice of L it suffices to show that every essential ideal I of L contains a nonzero element of

$f(L({}_R M))$. Define for each $g \in G$, $s_g: L({}_{R_e} M) \rightarrow L({}_{R_e} M)$ by $s_g(N) = R_g N$. Obviously for every $g, h \in G$ and $T \in L({}_{R_e} M)$ we have $s_g s_h(T) \leq s_{gh}(T)$ and $s_e = \text{id}$. Hence by Lemma 1 there is $N \leq_{R_e} M$ such that $f(N) \neq 0$ and $f(R_g N) \in I$ for all $g \in G$. Obviously $f(\sum_{g \in G} R_g N)$ is a nonzero element of $I \cap f(L({}_R M))$.

Going the same line as above in the dual situation but now applying the maps $s_g^0: L^0({}_{R_e} M) \rightarrow L^0({}_{R_e} M)$ given by $s_g^0(N) = R_{g^{-1}}^{-1}(N) = \{m \in M \mid R_{g^{-1}} m \leq N\}$ one obtains that $f(L^0({}_R M))$ is a dense sublattice of L^0 .

2. KRULL DIMENSION

In this section we study the Krull dimension (Kdim) and the dual Krull dimension (Ndim) of modules over G -graded rings. Recall that for a module M , $\text{Kdim } M$ ($\text{Ndim } M$) is defined (cf. [3, 7]) by induction as follows:

$\text{Kdim } M = -1$ ($\text{Ndim } M = -1$) if and only if $M = 0$;

if α is an ordinal number and $\text{Kdim } M \not\leq \alpha$ ($\text{Ndim } M \not\leq \alpha$) then $\text{Kdim } M = \alpha$ ($\text{Ndim } M = \alpha$) provided for every descending (ascending) chain $M_1 \geq M_2 \geq \dots$ ($M_1 \leq M_2 \leq \dots$) of submodules of M there is an n such that $\text{Kdim } M_i/M_{i+1} < \alpha$ ($\text{Ndim } M_{i+1}/M_i < \alpha$) for $i \geq n$.

One of basic properties of these dimensions says that if $N \leq M$ then $\text{Kdim } M = \sup\{\text{Kdim } N, \text{Kdim } M/N\}$ if either side exists. The same is true for Ndim .

Proposition 2. *Let M be a module, $\alpha \geq 0$ an ordinal number and let $\kappa_L(\eta_L)$ be the relation on $L = L(M)$ defined by $M_1 \kappa_L M_2$ ($M_1 \eta_L M_2$) if and only if $\text{Kdim}(M_1 + M_2/M_1 \cap M_2) < \alpha$ ($\text{Ndim}(M_1 + M_2/M_1 \cap M_2) < \alpha$). Then*

- (a) κ_L and η_L are congruences on L ;
- (b) $\text{Kdim } M = \alpha$ ($\text{Ndim } M = \alpha$) if and only if the lattice L/κ_L (L/η_L) is nonzero and satisfies DCC (ACC).

Proof. We prove the proposition for Kdim ; the proof for Ndim is similar. In view of Lemma 1.3.8 of [8], to prove that κ_L is a congruence it suffices to check that if $M_1 \leq M_2$ are submodules of M such that $\text{Kdim } M_2/M_1 < \alpha$ then for every submodule T of M , $\text{Kdim } M_2 + T/M_1 + T < \alpha$ and $\text{Kdim } M_2 \cap T/M_1 \cap T < \alpha$. However these are immediate consequences of isomorphisms $M_2 + T/M_1 + T \simeq M_2/M_1 + T \cap M_2$, $M_2 \cap T/M_1 \cap T \simeq M_1 + T \cap M_2/M_1$ and the property of Kdim formulated before the proposition.

It is clear that if the lattice L/κ_L is nonzero and satisfies DCC then $\text{Kdim } L = \alpha$. Conversely, let $\kappa_L(M_1) \geq \kappa_L(M_2) \geq \dots$ be a descending chain of L/κ_L . Obviously if $N_i = M_1 \cap \dots \cap M_i$ for $i = 1, 2, \dots$, then $N_1 \geq N_2 \geq \dots$ and $\kappa_L(N_i) = \kappa_L(M_i)$. Hence assuming $\text{Kdim } M = \alpha$ we get that for some n , $\kappa_L(M_n) = \kappa_L(N_n) = \kappa_L(N_{n+1}) = \kappa_L(M_{n+i})$, $i = 1, 2, \dots$, so the lattice L/κ_L satisfies DCC.

To obtain the main result of this section we also need the following

Lemma 2. *A dense sublattice K of a lattice L satisfies ACC if and only if the lattice L satisfies ACC.*

Proof. Obviously if L satisfies ACC then the same is true for K . Suppose that K satisfies ACC and L does not. Let M be an element of K maximal with respect to the property “ $[m, 1]$ does not satisfy ACC”. Let $a_1 < a_2 < \dots$ be a strictly ascending chain of elements of $[m, 1]$ and let $I = \bigcup_{k \geq 1} [m, a_k]$. Of course I is an ideal of $[m, 1]$. By Zorn’s Lemma there is an ideal J of $[m, 1]$ maximal with respect to the property $I \cap J = \{m\}$. Using modularity one can easily check that the ideal $I \vee J$ of $[m, 1]$ generated by $I \cup J$ is essential in $[m, 1]$. Hence the density property implies that there is $m_1 \in K \cap (I \vee J)$, $m_1 > m$. Hence for some $a \in I$, $b \in J$ we have $m < m_1 \leq a \vee b$. By the definition of I , $a \leq a_i$ for some i . Thus $[m_1, 1]$ contains the chain

$$a_i \vee b \leq a_{i+1} \vee b \leq \dots$$

The choice of m implies that it stabilizes. Hence for some $k \geq 0$, $a_{i+k+1} \vee b = a_{i+k} \vee b$. Now applying the modular law one obtains $a_{i+k+1} = a_{i+k+1} \wedge (a_{i+k+1} \vee b) = a_{i+k+1} \wedge (a_{i+k} \vee b) = a_{i+k} \vee (a_{i+k+1} \wedge b)$. But $I \cap J = \{m\}$, so we have $a_{i+k+1} \wedge b = m$. Hence $a_{i+k+1} = a_{i+k}$, contradiction.

Proposition 1 and Lemma 2 give immediately

Corollary 1. *If M is an R -module then*

- (i) $[10] \text{ } {}_R M \text{ is Noetherian if and only if } {}_{R_e} M \text{ is Noetherian;}$
- (ii) ${}_R M \text{ is Artinian if and only if } {}_{R_e} M \text{ is Artinian.}$

Now we prove the main result of this section.

Theorem 1. *For every R -module M , $\text{Kdim}_R M = \text{Kdim}_{R_e} M$ ($\text{Ndim}_R M = \text{Ndim}_{R_e} M$) if either side exists.*

Proof. We will concentrate here on Kdim ; the proof for Ndim is parallel.

The inequality $\text{Kdim}_R M \leq \text{Kdim}_{R_e} M$ is obvious. The proof of the other inequality is by induction on $\alpha = \text{Kdim}_R M$. The case $\alpha = 0$ is done in Corollary 1(ii). Now suppose that $\alpha > 0$ and the result holds for $\beta < \alpha$. The induction assumption gives $\kappa_K = \kappa_L \cap (K \times K)$, where $L = L({}_{R_e} M)$, $K = L({}_R M)$ and κ_L, κ_K are defined as in Proposition 2. Hence $K/\kappa_K \approx \{\kappa_L(N) \mid N \in K\} \subseteq L/\kappa_L$. Applying Proposition 1 and Lemma 2 we obtain that the lattice K/κ_K satisfies DCC if and only if the lattice L/κ_L satisfies DCC. This and Proposition 2 end the proof.

Theorem 1, and hence also Corollary 1, was proved in [4] when the grading is assumed inner.

3. GABRIEL DIMENSION

We will use the following Lanski’s [11] characterization of the Gabriel dimension, $\text{Gdim } M$, of a module M . It is given inductively starting with

$\text{Gdim } M = 0$ if and only if $M = 0$. Let α be a nonlimit ordinal and suppose that $\text{Gdim } M = \beta$ has been defined for $\beta < \alpha$. Call A an α -simple module if for every $0 \neq N \leq A$, both $\text{Gdim } N \not\leq \alpha$ and $\text{Gdim } A/N < \alpha$. Then $\text{Gdim } M = \alpha$ if $\text{Gdim } M \not\leq \alpha$ and if for each $N < M$, M/N contains a β -simple module for $\beta \leq \alpha$. When α is a limit ordinal, $\text{Gdim } M = \alpha$ if $\text{Gdim } M \not\leq \alpha$ and if for each $N < M$, M/N contains a β -simple module for $\beta < \alpha$.

Now we state some auxiliary facts which will be used in the proof of the main result of this section.

Proposition 3 [11]. (a) For any $N \leq M$,

$$\text{Gdim } M = \sup\{\text{Gdim } N, \text{Gdim } M/N\}$$

if either side exists.

(b) If $M_i \leq M$, $i \in I$ then $\text{Gdim } \sum_{i \in I} M_i = \sup\{\text{Gdim } M_i \mid i \in I\}$ if either side exists.

Proposition 4. Let α be an ordinal number and let γ_L be the relation on $L = L(M)$ given by $M_1 \gamma_L M_2$ if and only if $\text{Gdim}(M_1 + M_2/M_1 \cap M_2) < \alpha$. Then

- (i) γ_L is a congruence relation;
- (ii) if α is nonlimit and if N is an α -simple submodule of M then $\gamma_L(N)$ is an atom of L/γ_L ;
- (iii) if α is nonlimit and if for some $N \leq M$, $\gamma_L(N)$ is an atom of L/γ_L then there is $T < N$ such that $\text{Gdim } T < \alpha$ and N/T is α -simple.

Proof. The proof of (i) follows easily from Proposition 3(a). The condition (ii) is a direct consequence of the definition of γ_L . Now we prove (iii). Let $T = \sum\{U \leq N \mid \text{Gdim } U < \alpha\}$. By Proposition 3(b) $\text{Gdim } T < \alpha$. Now if $T < T' \leq N$ then Proposition 3(a) and the choice of T' guarantee that $\text{Gdim } T'/T \not\leq \alpha$. On the other hand, since $\gamma_L(N)$ is an atom of L/γ_L and $\gamma_L(T') \neq 0$, we have $\gamma_L(N) = \gamma_L(T')$. Hence, by the definition of γ_L , $\text{Gdim } N/T' < \alpha$.

Lemma 3. If $K = \{0, 1\}$ is a dense sublattice of a lattice L then $1 = a_1 \vee \cdots \vee a_n$ for some atoms $a_1, \dots, a_n \in L$.

Proof. By Lemma 2 the lattice L satisfies ACC, so it suffices to prove that for every $x \in L$ there is $y \in L$ such that $x \wedge y = 0$ and $x \vee y = 1$. Let J be an ideal of L maximal with respect to the property $[0, x] \cap J = \{0\}$. Then the ideal $[0, x] \vee J$ generated by $[0, x] \cup J$ is essential in L . Hence, since $\{0, 1\}$ is a dense sublattice of L , $1 \in [0, x] \vee J$. It means that $1 = x \vee y$ for some $y \in J$. Obviously $x \wedge y \in [0, x] \cap J = \{0\}$.

Lemma 4. Suppose that $\delta > 0$ is an ordinal number and M is an R -module such that for every $0 \neq N \leq_{R_e} M$ there exists $0 \neq N' \leq_{R_e} N$, $_{R_e} N' - \beta$ -simple for some $\beta < \delta$. Then there exists $0 \neq T \leq_{R_e} M$ such that for every $g \in G$, $R_g T = 0$ or $R_g T$ is a β -simple R_e -module for some $\beta < \delta$.

Proof. Let H be a subset of G containing e , of maximal cardinality such that for some $0 \neq X \leq_{R_e} M$ and all $h \in H$, $R_h X = 0$ or $R_h X$ is β -simple

R_e -module for some $\beta < \delta$. If there is $g \in G \setminus H$ then $R_g X \neq 0$. Hence there exists $0 \neq Y \leq_{R_e} R_g X$, $R_e Y$ β -simple for some $\beta < \delta$. Now for every $h \in H$, $R_{hg^{-1}} Y \leq R_{hg^{-1}} R_g X \leq R_h X$. Thus for every $\bar{h} \in Hg^{-1} \cup \{e\}$, $R_{\bar{h}} Y = 0$ or $R_{\bar{h}} Y$ is a β -simple R_e -module for some $\beta < \delta$. This contradiction proves that $G = H$, which gives the result.

Now we are ready to prove the main result of this section.

Theorem 2. *For every R -module M , $\text{Gdim}_R M = \text{Gdim}_{R_e} M$ if either side exists.*

Proof. We will prove by induction on α that if one side is equal to α then so is the other. It is clear for $\alpha = 0$. Suppose now that $\alpha > 0$ and the result holds for $\beta < \alpha$. Observe that this assumption says in particular that if γ_L and γ_K are the congruences defined in Proposition 4 on $L = L(R_e M)$ and $K = L(R M)$ respectively then $\gamma_K = \gamma_L \cap (K \times K)$. Thus $K/\gamma_K \approx \{\gamma_L(N) \mid N \in K\} \subseteq L/\gamma_L$ and by Proposition 1 K/γ_K can be treated as a dense sublattice of L/γ_L .

Suppose that $\text{Gdim}_R M = \alpha$. Since $\gamma_K = \gamma_L \cap (K \times K)$, $\text{Gdim}_{R_e} M \neq \alpha$. Thus we have to show that for every $N <_{R_e} M$, M/N contains a nonzero β -simple R_e -module, where $\beta \leq \alpha$ if α is nonlimit and $\beta < \alpha$ otherwise. Let T be the largest R -submodule of M which is contained in N . Passing to the module M/T we can assume that $T = 0$. Since $\text{Gdim}_R M = \alpha$ there is $0 \neq T' \leq_R M$, $R T'$ β -simple for some $\beta \leq \alpha$ if α is nonlimit and $\beta < \alpha$ otherwise. If $\beta < \alpha$ then by the induction assumption $\text{Gdim}_{R_e}(T'/T' \cap N) < \alpha$, so $0 < \text{Gdim}_{R_e}(T' + N/N) < \alpha$ and we are done. If $\beta = \alpha$ then by Proposition 4(ii) $\gamma_L(T')$ is an atom of the sublattice $\{\gamma_L(N) \mid N \in K\}$ of L/γ_L . Thus by Proposition 1 and Lemma 3 there are atoms $\gamma_L(N_1), \dots, \gamma_L(N_k)$ of L/γ_L such that $\gamma_L(T') = \gamma_L(N_1) \vee \dots \vee \gamma_L(N_k)$. It follows that the length of $[\gamma_L(N), \gamma_L(T' + N)]$ is finite. Hence by Proposition 4(iii) there is $_{R_e} \bar{N}$ such that $N < \bar{N} < T' + N$ and $_{R_e}(\bar{N}/N)$ is β -simple for some $\beta \leq \alpha$, as required.

Suppose that $\text{Gdim}_{R_e} M = \alpha$. By the induction assumption $\text{Gdim}_R M \neq \alpha$. Thus we have to show that for every $N <_R M$, M/N contains a β -simple R -submodule with $\beta \leq \alpha$ if α is nonlimit and $\beta < \alpha$ otherwise. We can obviously assume that $N = 0$. Observe that M satisfies the assumption of Lemma 4 with $\delta = \alpha$ in the limit case and $\delta = \alpha + 1$ otherwise. Hence there is $0 \neq X \leq_{R_e} M$ such that for every $g \in G$, $R_g X = 0$ or $R_g X$ is β -simple R_e -module for some $\beta \leq \alpha$ if α is nonlimit and $\beta < \alpha$ otherwise. Now $T = \sum_{g \in G} R_g X$ is a nonzero R -submodule of M . If for all $g \in G$, $\text{Gdim}_{R_e}(R_g X) < \alpha$ then the induction assumption gives $\text{Gdim}_R T < \alpha$ and we are done. If for some $g \in G$, $\text{Gdim}_{R_e}(R_g X) = \alpha$ then by Proposition 4(ii) $\gamma_L(T)$ is a join of atoms of L/γ_L . Hence also the lattice K/γ_K contains an atom $\gamma_K(T') \leq \gamma_K(T)$. Thus by Proposition 4(iii) there exists $T'' \leq_R T$ such that $\text{Gdim}_R T'' < \alpha$ and $_{R_e}(T'/T'')$ is α -simple. If $T'' \neq 0$ then it contains a

nonzero β -simple R -module for some $\beta < \alpha$. If $T'' = 0$ then T' is α -simple and we are done. This ends the proof.

4. REMARKS

1. The methods of this paper can be almost directly applied to obtain analogous results for some other than $R_e \subseteq R$ ring extensions of "finite type". Here we comment how they could be used to obtain some known results on normalizing extensions.

Recall that an extension $R \subseteq S$ of rings with the same 1 is finite normalizing if for some $s_1 = 1, s_2, \dots, s_n \in S$, $S = Rs_1 + \dots + Rs_n$ and $Rs_j = s_j R$ for $1 \leq j \leq n$.

It is known [2, 6, 12, 13], that if $R \subseteq S$ is a finite normalizing extension, M is an S -module then $\text{Kdim}_S M = \text{Kdim}_R M$ and $\text{Gdim}_S M = \text{Gdim}_R M$. It is clear that proving the above result by the method of this paper one will have some problems with Proposition 1 and Lemma 4 only. In the present situation instead of Proposition 1 one can apply the following

Proposition 1'. *Let M be an S -module and $\varphi_i, \varphi_i^0: L(RM) \rightarrow L(RM)$ be given by $\varphi_i(N) = s_i N$ and $\varphi_i^0(N) = \{m \in M \mid s_i m \in N\}$. If f is a homomorphism of $L(RM)$ onto a nonzero lattice L such that $f(X) = f(Y)$ implies $f(\varphi_i(X)) = f(\varphi_i(Y))$ ($f(\varphi_i^0(X)) = f(\varphi_i^0(Y))$) for $1 \leq i \leq n$, then $f(L(SM))$ ($f(L^0(SM))$) is a dense sublattice of L (L^0).*

Proof (Sketch). Similarly as in the proof of Proposition 1 to get that $f(L(RM))$ is a dense sublattice of L it suffices to show that every essential ideal I of L contains a nonzero element of $f(L(SM))$. The assumption on f and φ_i implies that for every $1 \leq i \leq n$, $g_i: L \rightarrow L$ given by $g_i(x) = f(\varphi_i(X))$ where $x = f(X)$, is a well-defined map. One easily checks that $g_i(0) = 0$, for every $x, y \in L$, $g_i(x \vee y) = g_i(x) \vee g_i(y)$ and if $x \leq y$ then $g_i([x, y]) = [g_i(x), g_i(y)]$. These imply that $g_i^{-1}(I)$, $1 \leq i \leq n$, are essential ideals of L . Hence $\bigcap_{i=1}^n g_i^{-1}(I) \neq 0$. Now if $0 \neq x \in \bigcap_{i=1}^n g_i^{-1}(I)$ then, since $g_1 = \text{id}$, $y = \bigvee_{i=1}^n g_i(x) \neq 0$. Clearly $y \in I \cap f(L(SM))$.

Applying φ_i^0 instead of φ_i one gets the dual result.

Instead of Lemma 4 one can apply the following.

Lemma 4'. *Suppose that $\delta > 0$ is an ordinal number, M is an S -module and N is a δ -simple R -submodule of M . Then for every $1 \leq i \leq n$, $s_i N$ is δ -simple R -module or $\text{Gdim}_R s_i N < \delta$.*

The proof follows from the fact that $L(Rs_i N) \approx L(R(N/K))$, where $K = \{x \in N \mid s_i x = 0\}$.

2. In [13] there was introduced the notion of the dual Gabriel dimension, $\text{Gcodim } M$, of a module M . The same proof as for Gdim gives the inequality $\text{Gcodim}_{R_e} M \geq \text{Gcodim}_R M$ (if $\text{Gcodim}_R M$ exists !) but we have not been able to obtain the other inequality.

REFERENCES

1. G. Birkhoff, *Lattice theory*, 3rd ed., Amer. Math. Soc., Providence, R.I., 1967.
2. J. Bit-David, *Normalizing extensions*. II, Proc. Ring Theory Conf., Lecture Notes in Math., vol. 825, Springer-Verlag, Berlin, Heidelberg and New York, pp. 6–9.
3. L. Chambless, *N-dimension and N-critical modules, application to Artinian modules*, Comm. Algebra **8** (1980), 1561–1592.
4. M. Cohen, *Smash product, inner actions and quotient rings*, Pacific J. Math. **125** (1986), 45–66.
5. J. W. Fisher, *Chain conditions for modular lattices with finite group actions*, Canad. J. Math. **31** (1978), 558–564.
6. J. W. Fisher, C. Lanski and J. K. Park, *Gabriel dimension of finite normalizing extensions*, Comm. Algebra **8** (1980), 1493–1503.
7. R. Gordon and J. C. Robson, *Krull dimension*, Mem. Amer. Math. Soc. No. 133, 1973.
8. G. Grätzer, *General lattice theory*, Birkhäuser-Verlag, Basel, 1978.
9. P. Grzeszczuk and E. R. Puczyłowski, *Goldie dimension and chain conditions for modular lattices with finite group actions*, Canad. Math. Bull. **29** (1986), 274–280.
10. P. Grzeszczuk, *On G-systems and G-graded rings*, Proc. Amer. Math. Soc. **95** (1985), 348–352.
11. C. Lanski, *Gabriel dimension and rings with involutions*, Houston J. Math. **4** (1978), 397–415.
12. B. Lemonnier, *Dimension de Krull et codeviation; application au théorème d'Eakin*, Comm. Algebra **6** (1978), 1647–1665.
13. B. Lemonnier, *Dimension et codimension de Gabriel dans les extensions triangulaires*, Comm. Algebra **14** (1986), 941–950.

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WARSAW, BIAŁYSTOK DIVISION,
AKADEMICKA 2, 15–267 BIAŁYSTOK, POLAND

INSTITUTE OF MATHEMATICS, UNIVERSITY OF WARSAW, PKIN, 00-901 WARSAW, POLAND