A NOTE ON ROE'S CHARACTERIZATION OF THE SINE FUNCTION

RALPH HOWARD

(Communicated by George R. Sell)

ABSTRACT. Let $f^{(n)}$, $n = 0, \pm 1, \pm 2, \dots$ be a sequence of complex valued functions on the real line with $(d/dx)f^{(n)} = f^{(n+1)}$ and satisfying inequalities $|f^{(n)}(x)| \leq M_n(1 + |x|)^k$ where as $n \rightarrow \infty$ the growth conditions $\lim_{n \rightarrow \infty} M_n(1 + \varepsilon)^{-n} = 0$ and $\lim_{n \rightarrow -\infty} M_{-n}(1 + \varepsilon)^{-n} = 0$ hold for all $\varepsilon > 0$. Then $f^{(0)}(x) = p(x)e^{ix} + q(x)e^{-ix}$ where p and q are polynomials of degree at most k .

In his paper [1] J. Roe proves

Theorem (Roe [1]). Let $\{f^{(n)}\}_{n=-\infty}^{\infty}$ be a two way infinite sequence of real valued functions defined on the real line R . Assume $f^{(n+1)}(x) = (d/dx)f^{(n)}(x)$ and that there is a constant M so that $|f^{(n)}(x)| \leq M$ for all n and x . Then $f^{(0)}(x) = a \sin(x + \phi)$ for some real constants a and ϕ .

This gives a rather striking characterization of the sine functions $a \sin(x + \phi)$ in terms of the size of their derivatives and antiderivatives. In this note we show that the bounds $|f^{(n)}(x)| \leq M$ can be relaxed to $|f^{(n)}(x)| \leq M_n(1 + |x|)^\alpha$ with $0 \leq \alpha < 1$ and where the constants only need to have supexponential growth. More generally:

Theorem. Let $\{f^{(n)}\}_{n=-\infty}^{\infty}$ be a sequence of complex valued functions defined on the real numbers with

$$(1) \quad f^{(n+1)}(x) = \frac{d}{dx} f^{(n)}(x)$$

and so that there are constants $M_n \geq 0$, $\alpha \in [0, 1)$, and a nonnegative integer k satisfying

$$(2) \quad |f^{(n)}(x)| \leq M_n(1 + |x|)^{k+\alpha}.$$

If

$$(3) \quad \lim_{n \rightarrow \infty} \frac{M_n}{(1 + \varepsilon)^n} = 0 \quad \text{all } \varepsilon > 0$$

Received by the editors July 28, 1986 and, in revised form, June 1, 1988.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 33A10; Secondary 42A38.

© 1989 American Mathematical Society
0002-9939/89 \$1.00 + \$.25 per page

and

$$(4) \quad \lim_{n \rightarrow \infty} \frac{M_{-n}}{(1 + \varepsilon)^n} = 0 \quad \text{all } \varepsilon > 0,$$

then

$$(5) \quad f^{(0)}(x) = p(x)e^{ix} + q(x)e^{-ix}$$

where $p(x)$ and $q(x)$ are polynomials of degree at most k .

Remark 1. The conclusion of the theorem can be sharpened by giving a more precise description of the functions $f^{(n)}$. If $k = 0, 1, 2, \dots$ and $n = 0, \pm 1, \pm 2, \dots$ then define

$$\begin{aligned} p_{n,k}(x) &= \left(1 + \frac{d}{dx}\right)^n x^k \\ &= \sum_{m=0}^k \frac{n(n-1)\cdots(n+m-k+1)}{(k-m)!} k(k-1)\cdots(m+1)x^m \end{aligned}$$

where for negative n we expand $(1 + d/dx)^n$ formally by use of Taylor's theorem. Then $p_{n,k}(x)$ is a polynomial of degree k and

$$p_{n,k}(x) + p'_{n,k}(x) = \left(1 + \frac{d}{dx}\right) p_{n,k}(x) = p_{n+1,k}(x).$$

When $n \geq 0$ these are Laguerre polynomials. This is because

$$(-1)^k p_{n,k}(-x) = \left(1 - \frac{d}{dx}\right)^n x^k = e^x \frac{d^n}{dx^n} (x^k e^{-x}).$$

See for example [0, p. 204]. This last equation implies that if λ is a complex number and

$$f_{k,\lambda}^{(n)}(x) = \lambda^n p_{n,k}(\lambda x) e^{\lambda x}$$

then $\{f_{k,\lambda}^{(n)}\}_{n=-\infty}^{\infty}$ satisfies equation (1). Then if $\{f^{(n)}\}_{n=-\infty}^{\infty}$ is a sequence of functions satisfying the hypothesis of the theorem then there are complex numbers $a_0, \dots, a_k, b_0, \dots, b_k$ so that

$$f^{(n)} = \sum_{m=0}^k (a_m f_{m,i}^{(n)} + b_m f_{m,-i}^{(n)})$$

where $(i)^2 = -1$. The proof of this from the theorem is done by induction on k . The details are left to the reader.

Remark 2. The functions $f_{k,\lambda}^{(n)}(x)$ just defined satisfy

$$|f_{k,\lambda}^{(n)}(x)| \leq (k+1)! n^k \max(|\lambda|^n, |\lambda|^{n+k}) (1 + |x|)^k e^{\operatorname{Re}(\lambda)x}.$$

By giving λ pure imaginary values close to i or $-i$ we see that there is no obvious weakening of the growth conditions (2), (3), or (4).

Remark 3. It is impossible to replace the interval $(-\infty, \infty)$ by a half infinite interval. The functions $f^{(n)}(x) = (-1)^n e^{-x}$ on $(0, \infty)$ yield a counterexample. (This observation is due to David Richman.)

Proof of the theorem. Let $f(x) = f^{(0)}(x)$. The, following [1], we will show the support of the Fourier transform $\hat{f}$ of f contained in the set $\{1, -1\}$. As the integral defining the Fourier transform may diverge, we define it as a distribution, that is as a linear functional on the vector spaces $\mathcal{S}$ of rapidly decreasing functions on R . Explicitly the value of $\hat{f}$ on $\phi \in \mathcal{S}$ is

$$\langle \hat{f}, \phi \rangle = \langle B, \hat{\phi} \rangle = \int_{-\infty}^{\infty} f(x) \hat{\phi}(x) dx.$$

Here we follow the notation of [2, Chapter 7].

Suppose it has been shown that the support of $\hat{f}$ is contained in $\{1, -1\}$. Then a standard result [2, Theorem 6.25, p. 150] implies there is an $m \geq 0$ and complex numbers a_j, b_j $0 \leq j \leq m$ so that

$$\hat{f} = \sum_{j=0}^m a_j \delta_1^{(j)} + \sum_{j=0}^m b_j \delta_{-1}^{(j)}$$

where δ_1 (resp. δ_{-1}) is the delta function at 1 (resp. at -1) and $\delta_1^{(j)}$ is the j th distributional derivative of δ_1 . This Fourier transform can be inverted to give that $f(x) = f^{(0)}(x)$ has the form given by (5) with $p(x)$ and $g(x)$ polynomials (of degree at most m). But $|f(x)| \leq |f^{(0)}(x)| \leq M_0(1 + |x|)^{k+\alpha}$. This implies the polynomials have degree at most k .

This reduces the proof to showing

Lemma 1. *The conditions (2) and (3) imply the support of $\hat{f}$ is disjoint from $(1, \infty)$ and $(-\infty, -1)$.*

Lemma 2. *The conditions (2) and (4) imply the support of $\hat{f}$ is disjoint from $(-1, 1)$.*

Proof of Lemma 1. We will only show the support of $\hat{f}$ is disjoint from $(1, \infty)$, the proof for $(-\infty, -1)$ being identical. Let ϕ be a smooth function with its support, $\text{spt}(\phi)$, in $(1, \infty)$. Then $\text{spt}(\phi) \subseteq [r, \infty)$ for some $r > 1$. We now need to show $\langle \hat{f}, \phi \rangle = 0$. Let $n \geq 0$ and

$$(6) \quad \psi_n(t) = \frac{\phi(t)}{(-it)^n}.$$

This is smooth as $\phi = 0$ near $t = 0$. Thus differentiating under the integral gives

$$\hat{\psi}_n^{(n)}(x) = \frac{d^n}{dx^n} \int_{-\infty}^{\infty} \frac{\phi(t)}{(-it)^n} e^{-itx} dx = \hat{\phi}(x).$$

So

$$\begin{aligned}
 \langle \hat{f}, \phi \rangle &= \langle f, \hat{\phi} \rangle \\
 &= \langle f, \hat{\psi}_n^{(n)} \rangle \\
 &= (-1)^n \langle f^{(n)}, \hat{\psi}_n \rangle \\
 &= (-1)^n \int_{-\infty}^{\infty} f^{(n)}(x) \hat{\psi}_n(x) dx.
 \end{aligned}$$

By (2) this implies

$$(7) \quad |\langle \hat{f}, \phi \rangle| \leq M_n \int_{-\infty}^{\infty} (1 + |x|^{k+\alpha}) |\hat{\psi}_n(x)| dx.$$

We now estimate $|\hat{\psi}_n(x)|$. First using that $\text{spt}(\phi) \subseteq [r, \infty)$,

$$(8) \quad |\hat{\psi}_n(x)| \leq \int_r^{\infty} \frac{|\phi(t)|}{t^n} dt \leq \frac{1}{r^n} \|\phi\|_{L^1}.$$

Also if $x \neq 0$ then integration by parts $(k+2)$ times yields

$$\begin{aligned}
 |\hat{\psi}_n(x)| &= \left| \int_r^{\infty} \left(\frac{d^{k+2}}{dt^{k+2}} \frac{\phi(t)}{(-it)^n} \right) \frac{e^{-ixt}}{(-ix)^{k+2}} dt \right| \\
 &\leq \frac{1}{|x|^{k+2}} \int_r^{\infty} \left| \frac{d^{k+2}}{dt^{k+2}} \frac{\phi(t)}{t^n} \right| dt \\
 &\leq \frac{1}{|x|^{k+2}} \sum_{j=0}^{k+2} \binom{k+2}{j} n(n+1) \\
 &\quad \cdots (n+j-1) \|\phi^{(k+2-j)}\|_{\infty} \int_r^{\infty} \frac{dt}{t^{n+j}} \\
 &= \frac{1}{|x|^{k+2}} \sum_{j=0}^{k+2} \binom{k+2}{j} \frac{n(n+1) \cdots (n+j-1) \|\phi^{(k+2-j)}\|}{(n+j-1)r^{n+j-1}} \\
 &\leq \frac{c_1(k, \phi) n^{k+1}}{|x|^{k+2} r^{n-1}}
 \end{aligned}$$

where $c_1(k, \phi)$ is a constant depending only on k and ϕ . This can be combined with (8) to give

$$(10) \quad |\psi_n(x)| \leq \begin{cases} 1/r^n \|\phi\|_{L^1} & |x| \leq 1, \\ \frac{c_1(k, \phi) n^{k+1}}{|k|^{k+2} r^{n-1}} & |x| > 1. \end{cases}$$

Using this in (7) gives an estimate of the form

$$(11) \quad |\langle \hat{f}, \phi \rangle| \leq c_2(k, \alpha, \phi) \frac{n^{k+1}}{r^{n-1}} M_n.$$

Let ε be so that $1 < 1 + \varepsilon < r$. Then for large n

$$\frac{n^{k+1}}{r^{n-1}} < \frac{1}{(1 + \varepsilon)^n}.$$

Using this and the condition (3) and (11)

$$|\langle \hat{f}, \phi \rangle| \leq c_2(k, \alpha, \phi) \lim_{n \rightarrow \infty} \frac{M_n}{(1 + \varepsilon)^n} = 0.$$

Therefore $\langle \hat{f}, \phi \rangle = 0$ for all ϕ with support in $(1, \infty)$, i.e., the support of $\hat{f}$ is disjoint from $(1, \infty)$ as claimed.

Proof of Lemma 2. The proof is very similar to the proof of Lemma 1. Let ϕ be a smooth function with support in $(-1, 1)$. Then for some $r < 1$ the inclusion $\text{spt}(\phi) \subseteq [-r, r]$ holds.

$$\begin{aligned} \langle \hat{f}, \phi \rangle &= \langle f^{(0)}, \hat{\phi} \rangle = \left\langle \frac{d^n}{dx^n} f^{(-n)}, \hat{\phi} \right\rangle \\ &= (-1)^n \langle f^{(-n)}, \hat{\phi}^{(n)} \rangle \\ &= (-1)^n \int_{-\infty}^{\infty} f^{(-n)}(x) \hat{\phi}^{(n)}(x) dx. \end{aligned}$$

By (2) this implies

$$(12) \quad |\langle \hat{f}, \phi \rangle| \leq M_{-n} \int_{-\infty}^{\infty} (1 + |x|)^{k+\alpha} |\hat{\phi}^{(n)}(x)| dx.$$

Differentiating under the integral and using $\text{spt}(\phi) \subseteq [-r, r]$

$$(13) \quad \hat{\phi}^{(n)}(x) = \int_{-r}^r (-it)^n \phi(t) e^{-itx} dt.$$

Thus

$$(14) \quad |\hat{\phi}^{(n)}(x)| \leq \int_{-r}^r |t|^n |\phi(t)| dt \leq 2r^n \|\phi\|_{L^1}.$$

Also for $x \neq 0$, integration by parts $(k+2)$ times and calculations similar to those of inequality (9) yield

$$\begin{aligned} |\hat{\phi}^{(n)}(x)| &= \left| \int_{-r}^r \left(\frac{d^{k+2}}{dt^{k+2}} (-it)^n \phi(t) \right) \frac{e^{-itx}}{(-ix)^{k+2}} dt \right| \\ (15) \quad &\leq \frac{1}{|x|^{k+2}} \int_{-r}^r \left| \frac{d^{k+2}}{dt^{k+2}} (t^n \phi(t)) \right| dt \\ &\leq \frac{c_3(k, \phi) h^{k+2}}{|x|^{k+2}} r^{n-k-2}. \end{aligned}$$

Putting the last two estimates together

$$(16) \quad |\hat{\phi}^{(n)}(x)| \leq \begin{cases} 2r^n \|\phi\|_{L^1} & |x| \leq 1, \\ \frac{c_3(k, \phi) h^{k+2}}{|x|^{k+2}} r^{n-k-2} & |x| > 1. \end{cases}$$

Putting this in (12) gives an estimate

$$|\langle \hat{f}, \phi \rangle| \leq c_4(k, \alpha, \phi) M_{-n} n^{k+2} r^{n-k-2}.$$

The proof is now completed in the same manner as the proof of Lemma 1.

ACKNOWLEDGMENT

I would like to thank Charles Nicol, David Richman, and Anton Schep for some interesting discussions related to this paper. I would also like to thank the referee for the observation that the polynomials $p_{n,k}$ in Remark 1 are Laguerre polynomials.

RÉFÉRENCES

0. E. D. Rainville, *Special functions*, Chelsea, New York, 1971.
1. J. Roe, *A characterization of the sine function*, Math. Proc. Cambridge Philos. Soc. **87** (1980), 69–73.
2. W. Rudin, *Functional analysis*, McGraw-Hill, New York, 1973.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTH CAROLINA, COLUMBIA, SOUTH CAROLINA 29208