SURJECTIVE ISOMETRIES OF WEIGHTED BERGMAN SPACES

CLINTON J. KOLASKI

(Communicated by Paul S. Muhly)

ABSTRACT. Let Ω be a bounded, simply connected domain in $\mathbb{C}^n = \mathbb{R}^{2n}$, let $F \in L^1(m_\Omega)$ be positive and continuous on Ω , and let $B_F^p(\Omega) = L^p(Fdm) \cap H(\Omega)$ ($0 < p < \infty$) denote the weighted Bergman space over Ω . We characterize those automorphisms Φ of Ω such that the map $f \rightarrow g \cdot (f \circ \Phi)$ is a surjective isometry of $B_F^p(\Omega)$, including an explicit description of $|g|$.

1. INTRODUCTION

In [2] it was shown that a surjective (linear) isometry of the Bergman space $B^p(\Omega) = L^p(m) \cap H(\Omega)$ ($0 < p < \infty$, $p \neq 2$) must have the form

$$(1.1) \quad f \rightarrow g \cdot (f \circ \Phi)$$

where Φ is an automorphism of the bounded Runge domain Ω , and g is related to Φ by (1.2).

Conversely, given an automorphism Φ of Ω , if there is a function g in $B^p(\Omega)$ related to Φ by

$$(1.2) \quad \int_{\Omega} (h \circ \Phi) |g|^p dm = \int_{\Omega} h dm$$

for every function h continuous on $\bar{\Omega} = \Omega \cup \partial\Omega$, then (1.1) defines a surjective isometry of $B^p(\Omega)$. In (1.2) dm denotes Lebesgue measure.

If the domain Ω is sufficiently nice, e.g., a ball or polydisc, then for every automorphism Φ there is a corresponding function $g = g_\Phi$ in $B^p(\Omega)$ which satisfies (1.2). It was conjectured in [2] that this would always be the case if the group of automorphisms acted transitively on Ω .

In this paper we shall characterize those automorphism Φ of a bounded, simply connected domain Ω which generate surjective isometries of the weighted Bergman space $B_F^p(\Omega) \equiv L^p(Fdm) \cap H(\Omega)$, where the weight function $F \in L^1(m_\Omega)$ is assumed only to be positive and continuous on Ω . Moreover, when

Received by the editors May 31, 1988.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 46E30; Secondary 30H05, 46J15.

Key words and phrases. Bergman space, isometry, automorphism.

the corresponding function g_Φ exists, an explicit description of $|g_\Phi|$ is given. One of the consequences is to answer the above conjecture affirmatively (cf. Remark (ii) and Theorem (2)).

2. OUR SETTING

Let n be a fixed positive integer, let Ω be a bounded domain in $\mathbb{C}^n = \mathbb{R}^{2n}$, Let $H(\Omega)$ denote the class of holomorphic functions on Ω , let $\text{Aut}(\Omega)$ denote the group of automorphisms (i.e., biholomorphic maps) of Ω , and let m_Ω denote Lebesgue measure on Ω normalized so that $m_\Omega(\Omega) = 1$. The closure of Ω in $\mathbb{C}^n$ is denoted by $\overline{\Omega}$.

When referring to the space

$$(2.1) \quad B_F^p(\Omega) \equiv L^p(Fdm_\Omega) \cap H(\Omega) \quad (0 < p < \infty)$$

we shall always assume that the weight function F satisfies

$$(2.2) \quad F \in L^1(m_\Omega) \text{ is POSITIVE and CONTINUOUS on } \Omega,$$

and that this space is given the usual metric of $L^p(Fdm_\Omega)$.

The following short argument shows $B_F^p(\Omega)$ is a closed subspace of $L^p(Fdm_\Omega)$ ($0 < p < \infty$).

Let $a \in \Omega$, choose $r > 0$ so that the closed ball $a + 2r\overline{B}$ lies in Ω , and let $\delta > 0$ denote the minimum of F on this closed ball.

For $f \in B_F^p(\Omega)$, and for z in the closed ball $a + r\overline{B}$, the subharmonicity of $|f|^p$ gives

$$(2.3) \quad |f(z)|^p \leq c \int_{z+r\overline{B}} |f|^p dm \leq \frac{c}{\delta} \int_{z+r\overline{B}} |f|^p F dm \leq \frac{c}{\delta} \int_\Omega |f|^p F dm$$

where c is a constant independent of f .

If $\{f_j\}$ is a Cauchy sequence in $B_F^p(\Omega)$, then (2.3) implies that $\{f_j\}$ converges uniformly on compact subsets of Ω . Hence, $B_F^p(\Omega)$ is closed in $L^p(Fdm_\Omega)$ ($0 < p < \infty$).

3. SURJECTIVE ISOMETRIES OF $B_F^p(\Omega)$

We now turn our attention to determining when an automorphism generates a surjective isometry.

It is easily shown that, given $\Phi \in \text{Aut}(\Omega)$, if there is a function g in $B_F^p(\Omega)$ ($0 < p < \infty$) related to Φ by

$$(3.1) \quad \int_\Omega (h \circ \Phi) |g|^p F dm = \int_\Omega h F dm$$

for every function h continuous on $\overline{\Omega}$, then (1.1) defines a surjective isometry of $B_F^p(\Omega)$. In the cases considered previously [2, 3] there was always such a function g , but in the present setting this need not be true. Before proving this we shall require a lemma which gives a converse to (3.1).

Lemma 1. Let $p > 0$, and let $\Phi \in \text{Aut}(\Omega)$. If there is a function g in $B_F^p(\Omega)$ such that the map $f \rightarrow g \cdot (f \circ \Phi)$ is a surjective isometry of $B_F^p(\Omega)$, then (3.1) holds.

Proof. Let X be the set of all functions continuous on $\overline{\Omega}$ which satisfy (3.1). By our hypothesis X contains all functions for the form $|f|^p$, where f is any analytic polynomial. It is clear from (3.1) that X is a linear space which is closed under complex conjugation. We shall show X is an algebra.

If $p = 2$, then $(f + h)(\bar{f} + \bar{h}) = |f + h|^2 \in X$, for every pair of analytic polynomials f and h . It now follows easily that $f\bar{h} \in X$. Since X is clearly closed under the supremum norm, an application of the Stone-Weierstrass theorem completes the proof of the case $p = 2$.

In a similar manner the case $p \neq 2$ follows from [5, Theorem 7.5.3] with $M =$ algebra of analytic polynomials, $Af = f \circ \Phi$, $d\mu_1 = Fdm_\Omega$ and $d\mu_2 = |g|^p Fdm_\Omega$.

Before we characterize the automorphisms which generate isometries, let us first observe that this class forms a subgroup of $\text{Aut}(\Omega)$.

Proposition 1. The automorphisms of Ω which generate isometries of $B_F^p(\Omega)$ ($0 < p < \infty$) form a subgroup of $\text{Aut}(\Omega)$. Moreover, if $\Phi, \Psi \in \text{Aut}(\Omega)$ have (respectively) corresponding functions $g, h \in B_F^p(\Omega)$, then

(a) $h \cdot (g \circ \Psi)$ corresponds to $\Phi \circ \Psi$ and

(b) $1/(g \circ \Phi^{-1})$ corresponds to Φ^{-1} .

Proof. (a) is obvious. To see (b) define U and T on $B_F^p(\Omega)$ by

$$Tf = g \cdot (f \circ \Phi) \text{ and } Uf = [1/(g \circ \Phi^{-1})](f \circ \Phi^{-1}).$$

By our hypothesis, T is a surjective isometry of $B_F^p(\Omega)$, so the range of T is equal to $B_F^p(\Omega)$. It is easy to show that $UT(f) = f$, for all $f \in B_F^p(\Omega)$, and it follows that $U = T^{-1}$; i.e., U is a surjective isometry of $B_F^p(\Omega)$. This proves (b), and the proposition follows.

We are now able to characterize those automorphisms which generate isometries.

Theorem 1. Let Ω be a bounded, simply connected domain in $\mathbb{C}^n$, and let $\Phi \in \text{Aut}(\Omega)$. Let $p > 0$. Then there is a function g in $B_F^p(\Omega)$ such that the map $f \rightarrow g \cdot (f \circ \Phi)$ is a surjective isometry of $B_F^p(\Omega)$ if and only if

$$(3.2) \quad \log(F \circ \Phi/F)$$

is pluriharmonic in Ω . Moreover, if (3.2) holds, then g is unique (modulo a unimodular constant), is never zero in Ω , and g is related to Φ by

$$(3.3) \quad |g|^p = (F \circ \Phi/F) \cdot |J\Phi|^2$$

where $J\Phi$ denotes the complex Jacobian of Φ .

Proof. Let $p > 0$, and let $\Phi \in \text{Aut}(\Omega)$.

Let us first suppose there is a function g in B_F^p such that the map $f \rightarrow g \cdot (f \circ \Phi)$ defines a surjective isometry of $B_F^p(\Omega)$. Then, by Lemma 1, Φ and g are related by (3.1). Thus, letting

$$(3.4) \quad G = |g|^p F / (J_R \Phi)$$

where $J_R \Phi$ denotes the (real) Jacobian of Φ , we find for every continuous function h on $\overline{\Omega}$,

$$(3.5) \quad \begin{aligned} \int_{\Omega} h F dm &= \int_{\Omega} (h \circ \Phi) |g|^p F dm \\ &= \int_{\Omega} \{[h \cdot (G \circ \Phi^{-1})] \circ \Phi\} (J_R \Phi) dm \\ &= \int_{\Omega} h \cdot (G \circ \Phi^{-1}) dm. \end{aligned}$$

Consequently, $F = G \circ \Phi^{-1}$; which with (3.4) shows

$$(3.6) \quad (F \circ \Phi) / F = |g|^p / (J_R \Phi).$$

Because our isometry is surjective, there is a function $f \in B_F^p(\Omega)$ such that $1 = g \cdot (f \circ \Phi)$. It follows that g is never zero in Ω ; thus $|g|^p = |g^p|$, where we use the principal branch of the logarithm in defining roots. We recall [5]; there is a holomorphic function $J\Phi$, the complex Jacobian of Φ , which is related to $J_R \Phi$ by $J_R \Phi = |J\Phi|^2$.

We may now write (3.6) in the form

$$(3.7) \quad (F \circ \Phi) / F = |g^p / (J\Phi)^2|,$$

which shows the function $(F \circ \Phi / F)$ is the modulus of a holomorphic function which is never zero in Ω . Thus (3.2) holds.

Conversely, if we assume (3.2) holds, then [5] our assumption that Ω be simply connected implies there is a function f in $H(\Omega)$ such that $\log(F \circ \Phi / F) = \operatorname{Re}(f)$; thus

$$(3.8) \quad F \circ \Phi / F = \exp[\operatorname{Re}(f)].$$

If we define

$$(3.9) \quad g = (J\Phi)^{2/p} \exp(f/p)$$

then g is a holomorphic function on Ω which (by (3.8)) satisfies (3.3). Moreover,

$$(3.10) \quad \int_{\Omega} |g|^p F dm = \int_{\Omega} (F \circ \Phi / F) |J\Phi|^2 F dm = \int_{\Omega} F dm < \infty.$$

Hence, $g \in B_F^p(\Omega)$.

Finally an argument similar to (3.10) shows g is related to Φ by (3.1); therefore, the map $f \rightarrow g \cdot (f \circ \Phi)$ is a surjective isometry of $B_F^p(\Omega)$. As noted above g is never zero in Ω , and (3.7) must hold. Clearly (3.7) gives (3.3), which then shows g is unique, modulo a unimodular constant.

Remarks. (i) The “only if” part is true for any bounded domain.

(ii) If $F \equiv 1$, then (3.2) is satisfied with $f \equiv 0$. Thus every $\Phi \in \text{Aut}(\Omega)$ generates a surjective isometry of $B_F^p(\Omega)$ ($0 < p < \infty$). Moreover, (3.9) shows the corresponding function g is given by $g = (J\Phi)^{2/p}$. This result is generalized in Theorem (2).

(iii) Note that (3.2) is independent of p . Thus, if an automorphism Φ generates an isometry for one $p > 0$, then it generates an isometry for every $p > 0$.

(iv) It is clear from (3.2) that it is the algebraic structure of the weight function F , not its behavior near the boundary of Ω , that determines which automorphisms generate isometries. As a simple example let Ω be the unit disc in $\mathbb{C}$. If $F(z) = (1 - |z|^4)^{-x}$ ($0 < x < 1$), or if $F(z) = \exp[-1/(1 - |z|^2)]$, then direct computation shows that (3.2) holds if and only if the automorphism is a rotation.

(v) Again let Ω be the unit disc in $\mathbb{C}$. If $F(z) = \exp(z^2 \bar{z} + z \bar{z}^2)$, then (3.2) is satisfied only by the identity automorphism.

(vi) If the analytic polynomials are dense in $B_F^p(\Omega)$, then the proof of Theorem (2.1) in [2] can be modified to show that every surjective isometry of $B_F^p(\Omega)$ ($p \neq 2$) is given by (1.1). Question: Is this true when the analytic polynomials are not dense in $B_F^p(\Omega)$?

4. SURJECTIVE ISOMETRIES OF $A^{p,r}(\Omega)$

The “classical” weighted Bergman spaces have the weight function F expressed as a power of the Bergman kernel. To be more precise, let

$$(4.1) \quad F_r(z) = [B_\Omega(z, z)]^{-r} \quad (z \in \Omega)$$

where B_Ω denotes the Bergman kernel for the bounded domain Ω , and r is greater than some negative constant $K(\Omega)$. The space $B_{F_r}^p(\Omega)$ is usually denoted as $A^{p,r}(\Omega)$ [1, 3, 4].

If the group $\text{Aut}(\Omega)$ acts transitively on Ω (which is true, for example, when Ω is a bounded symmetric domain), then [6],

$$(4.2) \quad B_\Omega(\Phi(z), \Phi(w)) \cdot (J\Phi)(z) \cdot (\overline{J\Phi})(w) = B_\Omega(z, w)$$

for all $(z, w) \in \Omega \times \Omega$, and for every $\Phi \in \text{Aut}(\Omega)$.

It now follows easily from (4.1) and (4.2) that

$$(4.3) \quad (F_r \circ \Phi)/F_r = |J\Phi|^{2r}$$

for every $\Phi \in \text{Aut}(\Omega)$ which, together with Theorem (1), gives

Theorem 2. *If the group $\text{Aut}(\Omega)$ acts transitively on Ω , then every $\Phi \in \text{Aut}(\Omega)$ generates a surjective isometry of $A^{p,r}(\Omega)$ for all $p > 0$ and all $r > K(\Omega)$. Moreover, the corresponding function g is given by*

$$(4.4) \quad g = (J\Phi)^{(2r+2)/p}$$

Remark. (vii) If Ω is taken to be the unit ball in $\mathbb{C}^n$, and if $\Phi \in \text{Aut}(\Omega)$, then [5],

$$(J\Phi)(z) = [(1 - |a|^2)/(1 - \langle z, a \rangle)^2]^{(n+1)/2}$$

where $\Phi(a) = 0$. Thus, Theorem 2 contains Theorem 4 of [3] as a special case (recall there $x = (n + 1)r$).

REFERENCES

1. C. Horowitz, *Zeros of functions in the Bergman spaces*, Duke Math. J. **41** (1974), 693–710.
2. C. Kolaski, *Isometries of Bergman spaces over bounded Runge domains*, Canad. J. Math. **33** (1981), 1157–1164.
3. —, *Isometries of Weighted Bergman spaces*, Canad. J. Math. **34** No. 4 (1982), 910–915.
4. D. H. Luecking, *Representation and duality in weighted spaces of analytic functions*, Indiana Univ. Math. J. **34**, No. 2 (1985), 319–336.
5. W. Rudin, *Function theory in the unit ball of $\mathbb{C}^n$* , Springer-Verlag, New York, 1980.
6. A. Selberg, *Automorphic functions and integral operators*, Seminars on Analytic Functions, II, Institute for Advanced Study, Princeton, N.J., 1957, pp. 152–161.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MINNESOTA-DULUTH, DULUTH, MINNESOTA 55812