NONCOMPACT COMMUTATORS IN THE COMMUTANT OF A CYCLIC OPERATOR

JAMES D. STAFNEY

(Communicated by John B. Conway)

ABSTRACT. We show that the commutant of the operator $S \oplus (I + S^*)$, where S is the shift operator, contains two operators A and B such that $AB - BA$ is not compact operator.

Let S be the shift operator and let $T = S \oplus (I + S^*)$. Herrero [1] has shown that T and T^* are cyclic and the commutant of T is not commutative. In this paper we show that the commutant of T fails to be commutative in the following stronger sense.

Theorem. *The commutant of the operator $S \oplus (I + S^*)$ contains two operators M , M_1 such that $MM_1 - M_1M$ is not a compact operator.*

Proof. For convenience, let $W = S \oplus (I + S^*)$. So, W is an operator (= bounded linear operator) on the Hilbert space $H = H^2 \oplus H^2$. Each operator A on H can be expressed as a matrix

$$A = \begin{pmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{pmatrix},$$

where each $A_{ij} \in \mathcal{L}(H^2)$, the bounded linear operators on H^2 . In particular,

$$W = \begin{pmatrix} S & 0 \\ 0 & I + S^* \end{pmatrix}.$$

An easy computation shows that the operators

$$E_1 = \begin{pmatrix} I & 0 \\ 0 & 0 \end{pmatrix}, \quad E_2 = \begin{pmatrix} 0 & 0 \\ C & 0 \end{pmatrix}$$

are in the commutant of W provided that $X = C$ satisfies the operator equation

$$(1) \quad (I + S^*)X = XS.$$

Received by the editors July 1, 1988.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 47B47.

Key words and phrases. Commutant, cyclic operator, commutator.

Furthermore, a simple computation shows that

$$E_1 E_2 - E_2 E_1 = \begin{pmatrix} 0 & 0 \\ -C & 0 \end{pmatrix}.$$

These observations show that the theorem follows from Lemma 1.

Lemma 1. *The operator equation (1) has a solution X in $\mathcal{L}(H^2)$ which is not a compact operator.*

To complete the proof of the theorem we now prove Lemma 1 based on the following lemmas and propositions. Let T and φ be as in Lemma 4. Let A be the operator on H^2 with matrix $(a_{k,n})$ corresponding to φ as in Proposition 3. Then Proposition 3 and Lemma 4 show that $X = A$ is a solution (in $\mathcal{L}(H^2)$) of (1) which is not compact.

Proposition 2. *The operator $X = A$ is a solution in $\mathcal{L}(H^2)$ of (1) if and only if the matrix $(a_{k,n})$, $k, n = 0, 1, 2, \dots$, of A relative to the orthonormal basis $u_n = e^{in\theta}$, $n = 0, 1, \dots$, satisfies:*

- (i) $a_{k,n} + a_{k+1,n} = a_{k,n+1}$;
- (ii) *the matrix $(a_{k,n})$ determines a bounded operator on l^2 .*

Proof. Suppose A is a solution of (1) and $(a_{k,n})$ is its matrix. Then (ii) is clearly satisfied. To prove (i) we note

$$\begin{aligned} \sum_k a_{k,n+1} u_k &= A u_{n+1} = A S u_n = (I + S^*) A u_n = (I + S^*) \sum_k a_{k,n} u_k \\ &= \sum_k a_{k,n} u_k + \sum_k a_{k+1,n} u_k. \end{aligned}$$

The converse follows from similar reasoning.

Let L^2 denote $L^2(\mu)$ where μ is normalized Lebesgue measure on the unit circle. We regard H^2 as a subspace of L^2 whenever appropriate. For each $\varphi \in L^2$, we define the transformation T_φ from L^2 into sequences by

$$T_\varphi f(k) = \int (e^{i\theta} - 1)^k \varphi(e^{i\theta}) f(e^{i\theta}) d\theta / 2\pi,$$

where $f \in L^2$, $k = 0, 1, \dots$.

Proposition 3. *For $\varphi \in L^2$, the matrix*

$$a_{k,n} \equiv \int (e^{i\theta} - 1)^k e^{in\theta} \varphi(e^{i\theta}) d\theta / 2\pi$$

is the matrix, relative to the orthonormal basis in Proposition 2, of a solution (compact solution) $X = A$ of (1) if and only if the restriction of T_φ to H^2 determines a bounded (compact) linear transformation of H^2 into l^2 .

Proof. If $(a_{k,n})$ is the matrix of a bounded operator, then T_φ clearly has the stated boundedness property. If T_φ has the stated boundedness property, then

$(a_{k,n})$ is the matrix of an operator $A \in \mathcal{L}(H^2)$. An easy computation shows that $(a_{k,n})$ satisfies (i) of Proposition 2. Thus, $X = A$ is a solution of (1).

Lemma 4. Let $T = T_\varphi$ where φ is the characteristic function of the arc $\{e^{i\theta} : 0 < \theta < \pi/3\}$. Then

- (i) T is a bounded linear transformation of L^2 into l^2 ;
- (ii) the restriction of T to H^2 is not a compact operator.

Proof. For $f \in L^2$,

$$Tf(k) = \int_0^{\pi/3} (e^{i\theta} - 1)^k f(e^{i\theta}) d\theta / 2\pi.$$

If $b_k = \int_0^1 x^k g(x) dx$, then since the Hilbert matrix is bounded,

$$\sum_k |b_k|^2 \leq C \int_0^1 |g(x)|^2 dx$$

for some constant C . From this and a routine change of variable argument with $x = |e^{i\theta} - 1|$, we conclude that T is bounded.

We now prove (ii). For convenience, let $v(\theta) = e^{i\theta} - 1$. A routine computation shows that

$$T^*Tf(e^{i\theta_1}) = \int_0^{\pi/3} [1/(1 - \overline{v(\theta_1)}v(\theta))] f(\theta) d\theta / 2\pi$$

for $f \in L^2$ and $0 \leq \theta_1 \leq \pi/3$; and, $T^*Tf(e^{i\theta_1}) = 0$, otherwise. We will show that the restriction of T^*T to H^2 is not compact. For convenience, let M denote the restriction of T^*T to H^2 . In particular, M is a bounded linear transformation from H^2 into L^2 . We will use Lemma 5 to show that M is not compact.

Lemma 5. Suppose that H, H' are Hilbert spaces, that A is a bounded linear transformation from H into H' and that $x_n \in H$ for $n = 1, 2, \dots$ such that

- (i) $\|x_n\| = 1$;
- (ii) $(x_m, x_n) \rightarrow 0$ as $n \rightarrow \infty$ for each fixed m ;
- (iii) $\|Ax_n\| \geq \delta > 0$ for all n .

Then A is not compact.

This lemma and Lemma 6 below show that M is not compact, which completes the proof of (ii) in Lemma 4.

For $0 < a < 1$, let $z_a = (1+a)e^{i\pi/3}$ and $f_a(e^{i\theta}) = \sqrt{a}/(e^{i\theta} - z_a)$. Clearly, each $f_a \in H^2$.

Lemma 6. There exist positive constants C_0, C_1, C_2 such that for $0 < a, d < 1$,

- (i) $C_0 \leq \|f_a\| \leq C_1$,

- (ii) $(f_a, f_d) \rightarrow 0$ as $d \rightarrow 0$ for each fixed a , and
 (iii) $\|Mf_a\| \geq C_2 \|f_a\|$ where $\|\cdot\|$, $(\cdot, \cdot)$ denote the norm and inner product on L^2 .

Proof. Let $b = 1 + a$ ($0 < a < 1$) and let $I = \int_{-\pi}^{\pi} 1/|e^{i\theta} - b|^2 d\theta$. By noting that

$$|e^{i\theta} - b|^2 = a^2 + b\theta^2 - 2bR(\theta),$$

where $R(\theta) = \cos \theta - (1 - \theta^2/2)$ one can show that

$$\left(\frac{6}{11}\right) \left(\frac{1}{a}\right) \leq I \leq (3\sqrt{2}\pi) \left(\frac{1}{a}\right);$$

and, (i) easily follows.

To prove (ii) we note that (f_a, f_d) is an integral over the interval $-\pi < \theta < \pi$ which is the sum of integrals over the subsets $J_1 = \{\theta: |\theta - \pi/3| < \delta\}$ and its complement J_2 in $[-\pi, \pi]$. By applying the Schwarz inequality to the integrals over each interval J_1 and J_2 , by noting that $\int_{J_1} |f_a|^2 d\theta \rightarrow 0$ as $\delta \rightarrow 0$ for a fixed and by noting that $f_d \rightarrow 0$ as $d \rightarrow 0$ uniformly in J_2 , we obtain (ii).

We now prove (iii). From the definition of M ,

$$Mf_a(\theta_1) = \int_0^{\pi/3} [1/(1 - \overline{v(\theta_1)}v(\theta))] f_a(\theta) d\theta/2\pi$$

for $0 < \theta_1 < \pi/3$. For convenience, let $F = F(a, \theta_1, \theta)$ denote the integrand in the latter integral. It is easily seen that

1. the argument of $f_a(\theta)$ varies over an interval of length $\leq \pi/2$ for $0 < a < 1$ and $0 \leq \theta \leq \pi/3$.

A geometric argument shows that $\overline{v(\theta_1)}v(\theta)$ lies in the intersection of two circles both of radius one and with centers at $e^{i\pi/3}$ and $e^{-i\pi/3}$ for $0 \leq \theta$, $\theta_1 \leq \pi/3$. From this we conclude that

2. the argument of $1 - \overline{v(\theta_1)}v(\theta)$ varies over an interval of length $\leq \pi/3$ for $0 \leq \theta$, $\theta_1 \leq \pi/3$; and, for some $C > 0$,

$$(2) \quad |1 - \overline{v(\theta_1)}v(\theta)| \leq C(1 - |\overline{v(\theta_1)}v(\theta)|), \quad 0 \leq \theta, \theta_1 \leq \pi/3.$$

Thus, in particular, the argument of F varies over an interval of length $\leq 5\pi/6$ for $0 < a < 1$ and $0 \leq \theta$, $\theta_1 \leq \pi/3$. So, for some complex number λ with $|\lambda| = 1$, the argument of λF lies in the interval $[-5\pi/12, 5\pi/12]$. Therefore,

$$|Mf_a(\theta_1)| \geq C \int_0^{\pi/3} |F(a, \theta_1, \theta)| d\theta, \quad 0 < \theta_1 < \pi/3$$

for some $C > 0$.

Let $t = |v(\theta)|$ and $s = |v(\theta_1)|$. From (2) and the definition of f_a it follows that

$$|F| \geq C(1/(1-st))(1/\sqrt{a}), \quad 0 < s < 1, 1-a < t < 1,$$

for some $C > 0$. By a change of variable we conclude that for some $C > 0$,

$$|Mf_a(\theta_1)| \geq (C/\sqrt{a}) \int_{1-a}^1 1/(1-st) dt \geq (C/\sqrt{a}) \log(a/2(1-s)) \geq 0$$

for $s > 1 - a/2$. Thus, for some $C > 0$, C independent of a , we have

$$\int_0^{\pi/3} |Mf_a(\theta_1)|^2 d\theta_1 \geq (C/a) \int_{1-a/2}^1 [\log(a/2(1-s))]^2 ds = (C/2) \int_0^1 (\log t)^2 dt,$$

which, together with (i), proves (ii).

ACKNOWLEDGMENT

The referee has pointed out that the operator $A = S^* \otimes I$ is cyclic and the commutant of A contains operators B and C such that $BC - CB$ is not compact. However, A^* is not cyclic. So, the significance of Herrero's example is that both T and T^* are cyclic. The referee has also noted that if A denotes the infinite direct sum of the operators $(1/n)I + 10^{-n}T$, $n = 1, 2, \dots$, then A and A^* are cyclic and the commutant of A contains a pair of operators with noncompact commutator. The point of our theorem is that it is not so clear that the operator T has this property. Because of the referees comments, we changed our original abstract and first paragraph.

REFERENCES

1. D. A. Herrero, *On multicyclic operators*, Integral Equations Operator Theory **1** (1978), pp. 57, 102.

UNIVERSITY OF CALIFORNIA, DEPARTMENT OF MATHEMATICS & COMPUTER SCIENCE, RIVERSIDE, CALIFORNIA 92521