CONTINUITY PROPERTIES OF THE SPECTRUM OF OPERATORS ON LEBESGUE SPACES

BRUCE A. BARNES

(Communicated by John B. Conway)

ABSTRACT. Fix $1 \leq p < s \leq \infty$. Let $T_x, x \in [p, s]$, be the collection of bounded linear operators on the Lebesgue spaces L^x determined by some fixed operator T . This paper concerns continuity properties of the map $x \rightarrow \sigma(T_x)$.

1. INTRODUCTION

Let Ω be a nonempty set, and let μ be a σ -finite measure on Ω . If $1 \leq p \leq \infty$, then let L^p denote the usual Lebesgue space $L^p(\Omega, \mu)$ with the usual p -norm. Now fix p and s with $1 \leq p < s \leq \infty$. Let $B_{p,s}$ be the algebra of all linear operators $T: L^p \cap L^s \rightarrow L^p \cap L^s$ with the property that T is continuous on $L^p \cap L^s$ with respect to both the p -norm and the s -norm. The algebra $\mathcal{B}_{p,s}$ is a Banach algebra which is studied in [2]. Also, some applications involving $\mathcal{B}_{p,s}$ can be found in [3]. If $T \in \mathcal{B}_{p,s}$, then by the Riesz Convexity Theorem [6, Theorem 11, p. 525] for each x in the interval $[p, s]$ the linear operator T has a unique extension to a bounded linear operator T_x on L^x (in the case where $s = \infty$, T_∞ is a bounded linear operator on the closure of $L^p \cap L^\infty$ in L^∞). Let $\sigma(T_x)$ and $r(T_x)$ denote the spectrum and spectral radius of this extension. It is well known that $\sigma(T_x)$ can be different for different x in $[p, s]$. An example of this phenomenon is given in [4]: Let $\Omega = (0, \infty)$ and μ be the Lebesgue measure. Take T to be the Cesàro operator, so

$$T_y(f)(t) = t^{-1} \int_0^t f(x) dx \quad (f \in L^y),$$

$1 < y \leq \infty$. Then $\sigma(T_y)$ is the circle with center and radius $(2(1 - y^{-1}))^{-1}$. Note that $\sigma(T_y)$ varies continuously in y in this case.

In this paper we study continuity properties of the map $y \rightarrow \sigma(T_y)$. A key theorem here is that for $T \in \mathcal{B}_{p,s}$, the function $y \rightarrow r(T_y)$ is continuous on the open interval (p, s) (it need not be continuous on $[p, s]$). A number of other continuity results are derived using this basic theorem as a tool.

Received by the editors July 26, 1988.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 47A10.

Key words and phrases. Linear operator, spectrum, spectral radius.

©1989 American Mathematical Society
0002-9939/89 \$1.00 + \$.25 per page

There are only a few examples in the literature where $\sigma(T_y)$ is specifically computed and this set is different for different values of y . Specific examples where this occurs can be found in [4], [5], and [7]. Next we give a number of additional examples.

Example 1. Let Ω be the positive integers, and let μ be the measure with

$$\mu(k) = 1/k! \quad (k \geq 1).$$

For $1 \leq p < \infty$, let T_p be the linear operator on $l^p(\mu)$ defined by $T_p(\{a_k\}) = \{b_k\}$ where $b_k = (k+1)^{-1}a_{k+1}$, $k \geq 1$. Also, let T_∞ act on c_0 by the same rule.

Note. For $1 < p \leq \infty$, T_p is compact and $\sigma(T_p) = \{0\}$.

Proof. Fix p , $1 < p < \infty$. Define M_n on $l^p(\mu)$ by $M_n(\{a_k\}) = \{b_k\}$ where $b_k = a_k$ for $1 \leq k \leq n$, and $b_k = 0$ for $k > n$. For $\{a_k\} \in l^p(\mu)$,

$$\begin{aligned} \|(T_p - M_n T_p)\{a_k\}\|_p^p &= \sum_{k=n+1}^{\infty} (1/k!)(k+1)^{-p}|a_{k+1}|^p \\ &= \sum_{k=n+1}^{\infty} (k+1)^{1-p}(1/(k+1)!)|a_{k+1}|^p \leq (n+2)^{1-p}\|\{a_k\}\|_p^p. \end{aligned}$$

Therefore

$$\|T_p - M_n T_p\|_p \leq (n+2)^{p^{-1}-1} \rightarrow 0$$

as $n \rightarrow \infty$. This proves T_p is compact. A similar computation shows that T_∞ is compact.

Now it is easy to see that the operator $M_n T_p$ is nilpotent, so $\sigma(M_n T_p) = \{0\}$ for $n \geq 1$. Since T_p is compact, this implies $\sigma(T_p) = \{0\}$ [8, Theorem 3].

Now we consider the case where $p = 1$. Let l^1 denote the usual sequence space (the L^1 -space of the positive integers with respect to counting measure). Define an isometry $W_1: l^1 \rightarrow l^1(\mu)$ by $W_1(\{a_k\}) = \{c_k\}$ where $c_k = k!a_k$, $k \geq 1$. The isometry W_1 maps l^1 onto $l^1(\mu)$. Let B be the unilateral backward shift on l^1 ; $B(\{a_k\}) = \{b_k\}$ where $b_k = a_{k+1}$, $k \geq 1$. A straightforward computation shows

$$W_1^{-1} T_1 W_1 = B \quad \text{on } l^1.$$

As is well known, $\sigma(B) = D$, the closed unit disk, so $\sigma(T_1) = D$.

To summarize: in this case $T \in \mathcal{B}_{1,\infty}$, T_x is compact with $\sigma(T_x) = \{0\}$ for $1 < x \leq \infty$, while T_1 is not compact with $\sigma(T_1) = D$. The spectral radius function is given by

$$r(T_x) = 0 \quad \text{for } 1 < x \leq \infty, \quad r(T_1) = 1.$$

Example II. Let Ω be the positive integers, and let μ be the measure with

$$\mu(k) = 2^k \quad (k \geq 1).$$

Let S and B be defined on any sequence space by

$$\begin{aligned} S(\{a_k\}) &= \{b_k\} & \text{where } b_1 &= 0, b_k = a_{k-1} \text{ for } k > 1; \\ B(\{a_k\}) &= \{c_k\} & \text{where } c_k &= a_{k+1} \text{ for } k \geq 1. \end{aligned}$$

Let S_p and B_p denote the operators S and B acting on the sequence space $l^p(\mu)$, $1 \leq p < \infty$. Let l^p be the usual L^p -space on the positive integers with counting measure. For $1 \leq p < \infty$ define $W_p: l^p \rightarrow l^p(\mu)$ by $W_p(\{a_k\}) = \{d_k\}$ where $d_k = 2^{-k/p} a_k$, $k \geq 1$. Then W_p is a linear isometry of l^p onto $l^p(\mu)$. Each of the following properties is established by a simple computation.

- (II.1) $S_2^* = 2B_2$, so $S_2 + 2B_2$ is selfadjoint on $l^2(\mu)$;
- (II.2) $W_p^{-1} S_p W_p = 2^{1/p} S$ on l^p ;
- (II.3) $W_p^{-1} B_p W_p = 2^{-1/p} B$ on l^p .

Now the spectrum of an operator of the form $a + bS + cB$, $a, b, c \in \mathbb{C}$, on the space l^p has been specifically computed by I. Gokhberg and M. Zambitsky. This result can be found in [9, Proposition 2]. Combining this information with (II.1)–(II.3), we can compute the spectrum of any operator of the form $a + bS_p + cB_p$, $1 \leq p < \infty$. We state these results for a specific operator:

Set $T_p = S_p + 2B_p \in \mathcal{B}(l^p(\mu))$, $1 \leq p < \infty$. Then

- (II.4) $\sigma(T_2) = \{\lambda \in \mathbb{R} : -2\sqrt{2} \leq \lambda \leq 2\sqrt{2}\}$;
- (II.5) For $p \neq 2$, $p^{-1} + q^{-1} = 1$ ($q^{-1} = 0$ when $p = 1$), $\sigma(T_p)$ is the set of all $\lambda = x + iy$, $x, y \in \mathbb{R}$, which lie inside or on the ellipse $x^2/a^2 + y^2/b^2 = 1$ where $a = (2^{1/p} + 2^{1/q})$ and $b = (2^{1/p} - 2^{1/q})$.
- (II.6) $r(T_p) = 2^{1/p} + 2^{1-1/p}$, $1 \leq p < \infty$.

The operator T_2 in this example is selfadjoint. In this case it is always true that when $2 \leq x \leq y$ or $y \leq x \leq 2$, then $\sigma(T_x) \subseteq \sigma(T_y)$; see [2, Theorem 4.4]. Also, the fact that $\sigma(T_p) \neq \sigma(T_2)$ for $p \neq 2$ shows that the hypothesis (4.1) in §4 of [1] cannot be omitted and still obtain the results in [1, Theorem 4.8].

2. CONTINUITY PROPERTIES

The main result of this paper concerns the continuity of the spectral radius function $x \rightarrow r(T_x)$; this result is stated in Theorem 3. On the way to proving this theorem, we derive some additional interesting continuity properties of the spectrum.

Proposition 1. Assume $T \in \mathcal{B}_{p,s}$ and $r(T_y) = 0$ for some $y \in [p, s]$. Then $r(T_x) = 0$ for all $x \in (p, s)$.

Proof. Fix $x \in (p, s)$. We may assume $x < y$. Choose z such that $z \in (p, s)$ and $z < x$. Then $\exists t \in (0, 1)$ with $x^{-1} = tz^{-1} + (1-t)y^{-1}$. Applying the Riesz Convexity Theorem, we have for all $n \geq 1$

$$\|(T^n)_x\| \leq \|(T^n)_z\|^t \|(T^n)_y\|^{1-t}.$$

Therefore

$$r(T_x) \leq r(T_z)^t r(T_y)^{1-t} = 0.$$

Corollary 2. Assume $T \in \mathcal{B}_{p,s}$ and for some $y \in [p, s]$ $\sigma(T_y)$ is finite. Then $\sigma(T_x) = \sigma(T_y)$ for all $x \in (p, s)$.

Proof. Assume $\sigma(T_y) = \{\lambda_1, \dots, \lambda_n\}$. Let q be the polynomial $q(z) = \prod_{k=1}^n (z - \lambda_k)$. Then $\sigma(q(T_y)) = q(\sigma(T_y)) = \{0\}$. It follows from Proposition 1 that $\sigma(q(T_x)) = \{0\}$ for all $x \in (p, s)$. Thus, $\sigma(T_x) \subseteq \{\lambda_1, \dots, \lambda_n\}$ for all such x . Fix any $x \in (p, s)$, $x \neq y$. We may assume $x < y$. Now $T \in \mathcal{B}_{x,y}$, and both $\sigma(T_x)$ and $\sigma(T_y)$ are finite. Then by [2, Cor. 5.2], $\sigma(T_x) = \sigma(T_y)$.

Now we prove the main result.

Theorem 3. Assume $T \in \mathcal{B}_{p,s}$. Then $r(T_x)$ is a continuous function on (p, s) .

Proof. If $r(T_y) = 0$ for some $y \in (p, s)$, then $r(T_x) = 0$ for all $x \in (p, s)$ by Proposition 1. Thus, we may assume $r(T_x) > 0$ for all $x \in (p, s)$. For each $n \geq 1$, define

$$\varphi_n(w) = n^{-1} \log(\|(T^n)_{w^{-1}}\|).$$

By the Riesz Convexity Theorem [6, Theorem 11, p. 525] φ_n is a convex function on (s^{-1}, p^{-1}) . Now $\varphi_n(w)$ converges pointwise to $\varphi(w) = \log(r(T_{w^{-1}}))$. By hypothesis $\varphi(w)$ has only finite values, and φ must be convex on (s^{-1}, p^{-1}) . It follows from [10, Theorem 3.2] that $\varphi(w)$ is continuous on (s^{-1}, p^{-1}) , and thus, $r(T_x)$ is continuous on (p, s) .

Concerning this result, note that Example I shows that $x \rightarrow r(T_x)$ need not be continuous on the closed interval $[p, s]$.

Next we derive some consequences of the continuity of the spectral radius function. The first is a type of upper semicontinuity property for the function $x \rightarrow \sigma(T_x)$. For K a compact subset of $\mathbb{C}$, let $\hat{K}$ denote the polynomial convex hull of K . E. Stout's book [11] is a good source for information on polynomial convexity.

Theorem 4. Assume $T \in \mathcal{B} = \mathcal{B}_{p,s}$ and fix $x \in (p, s)$. Let U be an open set in $\mathbb{C}$ with $\sigma(T_x)^\wedge \subseteq U$. Then $\exists \delta > 0$ such that $\sigma(T_y) \subseteq U$ whenever $|y - x| < \delta$.

Proof. Suppose no such δ exists. Then we can choose a sequence $\{y_n\} \subseteq (p, s)$ with $y_n \rightarrow x$ such that for each n there exists $\lambda_n \in \sigma(T_{y_n})$ with $\lambda_n \notin U$. The sequence $\{\lambda_n\}$ is contained in the compact set $\sigma_{\mathcal{B}}(T)$, and so some subsequence converges to a number $\lambda \notin U$. We may assume that $\lambda_n \rightarrow \lambda$.

Set $K = \sigma(T_x)^\wedge$, and if q is a polynomial, then let $\|q\|_K = \sup\{|q(\mu)| : \mu \in K\}$. Now $\lambda \notin K$, so by definition there is a polynomial q such that $|q(\lambda)| > \|q\|_K = 1$. Choose a positive integer n sufficiently large that $|q^n(\lambda)| > 2$. Set $p = q^n$. Thus,

$$|p(\lambda)| > 2 \quad \text{and} \quad \|p\|_K = 1.$$

Since $r(p(T_x)) = 1$, by the continuity of the spectral radius (Theorem 3) $r(p(T_{y_k})) \rightarrow 1$. Also, $p(\lambda_k) \in \sigma(p(T_{y_k}))$ and $p(\lambda_k) \rightarrow p(\lambda)$. This provides a contradiction since for k sufficiently large

$$r(p(T_{y_k})) < 2 \quad \text{and} \quad |p(\lambda_k)| > 2.$$

Next we prove a stronger upper semicontinuity property of the map $x \rightarrow \sigma(T_x)$ which holds under the assumption that the spectrum of T in the algebra $\mathcal{B}_{p,s}$ is “thin”. First we need a preliminary result that uses the continuity of the spectral radius function.

Lemma 5. *Assume $T \in \mathcal{B} = \mathcal{B}_{p,s}$ and $\lambda \notin \sigma_{\mathcal{B}}(T)$. Then $d(x) = \text{dist}(\lambda; \sigma(T_x))$ is a continuous function on (p, s) .*

Proof. For all $x \in (p, s)$ $((\lambda - T)^{-1})_x = (\lambda - T_x)^{-1}$. Thus since

$$\sigma((\lambda - T_x)^{-1}) = \{(\lambda - \mu)^{-1} : \mu \in \sigma(T_x)\},$$

it follows that for $x \in (p, s)$

$$r(((\lambda - T)^{-1})_x) = \sup\{|\lambda - \mu|^{-1} : \mu \in \sigma(T_x)\} = (d(x))^{-1}.$$

By Theorem 3 this function is continuous on (p, s) which proves the lemma.

Theorem 6. *Assume $T \in \mathcal{B} = \mathcal{B}_{p,s}$ and that $\sigma_{\mathcal{B}}(T)$ has empty interior. If $x \in (p, s)$ and U is an open set with $\sigma(T_x) \subseteq U$, then $\exists \delta > 0$ such that $\sigma(T_y) \subseteq U$ whenever $|x - y| < \delta$.*

Proof. Suppose no such δ exists. Then there is a sequence $\{y_n\} \subseteq (p, s)$ such that $y_n \rightarrow x$ and for each n there exists $\lambda_n \in \sigma(T_{y_n})$ with $\lambda_n \notin U$. We may assume (by taking a subsequence if necessary) that $\{\lambda_n\}$ converges to a number $\mu \notin U$. Let $\varepsilon = \text{dist}(\mu; \sigma(T_x)) > 0$. Set $D = \{\alpha \in \mathbb{C} : |\alpha - \mu| < \varepsilon/4\}$. Since $\sigma_{\mathcal{B}}(T)$ has no interior, we can choose $\lambda \in D$ with $\lambda \notin \sigma_{\mathcal{B}}(T)$. By Lemma 5 $d(y) = \text{dist}(\lambda; \sigma(T_y))$ is a continuous function on (p, s) . By the choices of ε and λ , $d(x) > \varepsilon/2$. But since $\lambda_n \rightarrow \mu$, we have $d(y_n) < \varepsilon/2$ for all n sufficiently large, a contradiction.

Next we turn to some results concerning the situation where $\sigma(T_x)^{\wedge}$ is disconnected or has an isolated point at some $x \in (p, s)$. What we prove is that this property extends to all y in some neighborhood of x .

We need the following technical lemma.

Lemma 7. *Let K and J be compact subsets of $\mathbb{C}$ with $\hat{K}$ and $\hat{J}$ disjoint. Then there exists U and V , open sets with compact closure, such that*

- (i) $\hat{K} \subseteq U$ and $\hat{J} \subseteq V$; and
- (ii) $\overline{U}^{\wedge}$ and $\overline{V}^{\wedge}$ are disjoint.

Proof. For each $\mu \in \hat{J}$, choose a polynomial q_{μ} such that $\|q_{\mu}\|_K = 1$ and $|q_{\mu}(\mu)| > 3$. Let

$$V_{\mu} = \{\lambda \in \mathbb{C} : |q_{\mu}(\lambda)| > 3\}, \quad \text{and} \\ U_{\mu} = \{\lambda \in \mathbb{C} : |q_{\mu}(\lambda)| < 2\}.$$

Let $\{V_{\mu_1}, \dots, V_{\mu_n}\}$ be a finite cover for $\hat{J}$. For convenience, in the notation that follows the subscript μ_k will be replaced by k . Note that $\hat{J} \subseteq \bigcup_{k=1}^n V_k$ and $\hat{K} \subseteq \bigcap_{k=1}^n U_k$. Choose U with compact closure such that

$$\hat{K} \subseteq U \subseteq \overline{U} \subseteq \bigcap_{k=1}^n U_k.$$

If $\lambda \in \overline{U}^\wedge$, then $\|q_k\|_{\overline{U}} < 2$, so $|q_k(\lambda)| < 2$ for $1 \leq k \leq n$. Thus, $\overline{U}^\wedge \subseteq \bigcap_{k=1}^n U_k$. Since $\bigcap_{k=1}^n U_k$ and $\bigcup_{k=1}^n V_k$ are disjoint, it follows that $\overline{U}^\wedge$ and $\hat{J}$ are disjoint.

Now repeat the argument above with $\overline{U}$ in place of J and J in place of K . The argument proves $\exists V$ an open set with compact closure such that $\hat{J} \subseteq V$ and $\overline{V}^\wedge$ and $\overline{U}^\wedge$ are disjoint.

Theorem 8. Let $T \in \mathcal{B}_{p,s}$. Assume for some $x \in (p, s)$, $\sigma(T_x)^\wedge$ is disconnected. Then $\sigma(T_y)$ is disconnected for all y in some neighborhood of x .

Proof. Assume K and J are disjoint nonempty compact sets with $\sigma(T_x)^\wedge = K \cup J$. Then $\hat{K} = K$ and $\hat{J} = J$, so by Lemma 7, there exist open sets U and V with compact closure such that $K \subseteq U$, $J \subseteq V$, and $\overline{U}^\wedge$ and $\overline{V}^\wedge$ are disjoint. By Theorem 4 $\exists \delta > 0$ such that $\sigma(T_y) \subseteq U \cup V$ whenever $|x - y| < \delta$. Fix y with this property, and assume (without loss of generality) that $x < y$. Then $T \in \mathcal{B} = \mathcal{B}_{x,y}$. By [2, Theorem 5.1(4)]

$$\sigma_{\mathcal{B}}(T) \subseteq \sigma_{\mathcal{B}}(T)^\wedge = (\sigma(T_x) \cup \sigma(T_y))^\wedge \subseteq (\overline{U} \cup \overline{V})^\wedge = \overline{U} \cup \overline{V},$$

where the last equality follows from [11, Lemma 29.21(b)]. Now since $\sigma(T_x)$ has nonempty intersection with both U and V , the same is true for $\sigma_{\mathcal{B}}(T)$. Then using the fact that $S \rightarrow S_y$ is a continuous algebra monomorphism of $\mathcal{B}$ into $\mathcal{B}(L^y)$, it follows from [2, Theorem 4.5] that $\sigma(T_y) \cap U$ and $\sigma(T_y) \cap V$ are nonempty. Thus, $\sigma(T_y)$ is disconnected.

Corollary 9. Let $T \in \mathcal{B}_{p,s}$. Assume for some $x \in (p, s)$ that λ_0 is an isolated point of $\sigma(T_x)^\wedge$. Then λ_0 is an isolated point of $\sigma(T_y)$ for all y in some neighborhood of x .

Proof. Set $K = \{\lambda_0\}$ and $J = \sigma(T_x)^\wedge \setminus \{\lambda_0\}$. Choose $U, V, \delta > 0, y$, and $\mathcal{B}$ is in the proof of Theorem 8. Then $\sigma_{\mathcal{B}}(T) \subseteq \overline{U} \cup \overline{V}$. Let $Q \in \mathcal{B}$ be the spectral projection corresponding to the set $\sigma_{\mathcal{B}}(T) \cap \overline{U}$ which is a nonempty open and closed subset of $\sigma_{\mathcal{B}}(T)$. Then by the Spectral Mapping Theorem

$$\sigma((QT)_x) = \sigma(Q_x T_x) = \{\lambda_0\}.$$

It follows from Corollary 2 that for all $w \in (x, y)$, $\sigma((QT)_w) = \{\lambda_0\}$. Therefore λ_0 is an isolated point of $\sigma(T_w)$ when $x < w < y$. This argument proves that λ_0 is an isolated point of $\sigma(T_w)$ whenever $|x - w| < \delta$.

Added in proof. It has been pointed out to me that a general upper semicontinuity property of the map $y \rightarrow \sigma(T_y)$ has been proved by I. Ya. Sneiberg in

[Mat. Issled., 9 (1974), 214–229 (in Russian)]. Both Theorems 4 and 6 are special cases of this result. On the positive side, this same result implies that Theorem 8 and Corollary 9 have more general forms. Specifically, $\sigma(T_x)^\wedge$ can be replaced by $\sigma(T_x)$ in the hypotheses of these two results without altering the conclusions.

REFERENCES

1. B. Barnes, *The spectrum of integral operators on Lebesgue spaces*, J. of Operator Th. **18** (1987), 115–132.
2. —, *Interpolation of spectrum of bounded operators on Lebesgue spaces*, to appear in Rocky Mt. J. Math.
3. —, *Essential spectrum in a Banach algebra applied to linear operators*, recently submitted.
4. D. Boyd, *The spectrum of the Cesàro operator*, Acta. Sci. Math. **29** (1968), 31–34.
5. K. Dayanithy, *Interpolation of spectral operators*, Math. Z. **159** (1978), 1–2.
6. N. Dunford and J. Schwartz, *Linear operators*, Part I, Interscience, New York-London, 1964.
7. D. Herrero, *Operator theory*, Advances and Applications, Vol. 11, Birkhauser-Verlag, Basel, 1983, 191–232.
8. J. Newburgh, *The variation of spectra*, Duke Math. J. **18** (1951), 165–176.
9. J. Nieto, *On the essential spectrum of symmetrizable operators*, Math. Annalen **178** (1968), 145–153.
10. W. Rudin, *Real and complex analysis*, Third Ed., McGraw-Hill, New York, 1987.
11. E. Stout, *The theory of uniform algebras*, Bodgen & Quigley, Tarrytown-on-Hudson, New York, 1971.

MATHEMATICS DEPARTMENT, UNIVERSITY OF OREGON, EUGENE, OREGON 97403