NON-NEGATIVE SOLUTIONS FOR A CLASS OF RADially SYMMETRIC NON-POSITONE PROBLEMS

ALFONSO CASTRO AND R. SHIVAJI

(Communicated by Walter D. Littman)

ABSTRACT. We consider the existence of radially symmetric non-negative solutions for the boundary value problem

$$\begin{aligned} -\Delta u(x) &= \lambda f(u(x)) & \|x\| \leq 1, \quad x \in \mathbb{R}^N \quad (N \geq 2) \\ u(x) &= 0 & \|x\| = 1 \end{aligned}$$

where $\lambda > 0$, $f(0) < 0$ (non-positone), $f' \geq 0$ and f is superlinear. We establish existence of non-negative solutions for λ small which extends some work of our previous paper on non-positone problems, where we considered the case $N = 1$. Our work also proves a recent conjecture by Joel Smoller and Arthur Wasserman.

1. INTRODUCTION

Here we consider the existence of radially symmetric non-negative solutions for the boundary value problem

$$(1.1) \quad -\Delta u(x) = \lambda f(u(x)) \quad \|x\| < 1, \quad x \in \mathbb{R}^N, \quad N \geq 2$$

$$(1.2) \quad u(x) = 0 \quad \|x\| = 1$$

where $\lambda > 0$ and $f: [0, \infty) \rightarrow \mathbb{R}$ is such that $f' \geq 0$. As is well documented, the study of (1.1)–(1.2) is equivalent to the problem

$$(1.3) \quad -u'' - (n/r)u' = \lambda f(u); \quad r \in (0, 1)$$

$$(1.4) \quad u'(0) = 0$$

$$(1.5) \quad u(1) = 0,$$

where $n = N - 1$. We will assume that

$$(1.6) \quad \lim_{u \rightarrow +\infty} (f(u))/u = +\infty, \text{ i.e., } f \text{ is superlinear,}$$

Received by the editors October 26, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 35J65, 34B15.

Key words and phrases. Non-negative solution, non-positone problem, singular o.d.e., Dirichlet problem.

Part of the first author's research was done while visiting Argonne National Laboratory.

The second author was supported by a Mississippi State University research initiation grant.

$$(1.7) \quad f(0) < 0 \text{ (non-positone),}$$

and for some $k \in (0, 1)$,

$$(1.8) \quad \Lambda = \lim_{d \rightarrow +\infty} (d/f(d))^{N/2} \{F(kd) - [(N-2)/(2N)]df(d)\} = +\infty$$

where $F(x) = \int_0^x f(r) dr$.

If $f(0) > 0$ (positone) and $\lambda > 0$ small, it is known that (1.1)–(1.2) has two solutions: one near zero, the other bifurcating from infinity. However, the popular method of sub-super solutions used in positone problems seems rather difficult to apply when $f(0) < 0$, since $v \equiv 0$ is no longer a sub-solution. In fact, it is a super solution. This is why we have been motivated to undertake this study. Our main result is given in Theorem 1.1.

Theorem 1.1. *Under the above assumptions, there exists $\lambda_0 > 0$ such that if $0 < \lambda < \lambda_0$, then (1.1)–(1.2) has a non-negative solution u_λ such that $u_\lambda > 0$ and decreasing on $[0, 1]$ and $u'_\lambda(1) < 0$.*

Castro and Shivaji [3] have made an extensive study of the one-dimensional problem ($N = 1$). Our proof of Theorem 1.1 is based on the shooting method. That is, to prove that (1.3)–(1.5) has a solution, we consider the problem (1.3)–(1.4) subject to $u(0) = d$. By analyzing this problem depending on the parameter d , we show that for an adequate value of d , u satisfies also (1.5). To prove Lemma 3.2 we use an identity of Pohozaev type (see §2) used by Castro and Kurepa [1, 2] to study oscillatory solutions of other radially symmetric problems. For other applications and extensions of this type of identity, see Ni and Serrin [4].

Our work also proves a recent conjecture by Smoller and Wasserman [6]. In their paper they proved an existence result applicable to functions of the type $f(u) = u^q - \varepsilon$ where $\varepsilon > 0$, $1 < q < N/(N-2)$ and conjectured that an optimal result would be to extend it to $1 < q < (N+2)/(N-2)$. In fact, our work includes this optimal result since if $f(u) = u^q - \varepsilon$ where $\varepsilon > 0$, $1 < q < (N+2)/(N-2)$ then (1.8) is satisfied with k chosen larger than $[(q+1)(N-2)/(2N)]^{(1/(q+1))}$. Note here that if $q < (N+2)/(N-2)$ then $(q+1)(N-2)/(2N) < \{(N+2)/(N-2) + 1\}(N-2)/(2N) = 1$.

We will restrict our proofs in this paper to the case $N > 2$. When $N = 2$ the proof is easier along the same lines as in the case $N > 2$.

2. PRELIMINARIES AND NOTATIONS

First of all we extend f to $(-\infty, \infty)$ by defining $f(x) = f(0)$ for $x < 0$. By (1.6) we see that $\lim_{d \rightarrow \infty} F(d) = \infty$. Hence (see (1.7)) there exist positive real numbers $\beta < \theta$ such that

$$(2.1) \quad 0 = f(\beta) = F(\theta).$$

Since (see (1.8)) $\Lambda = \infty$, we see that there exists $\gamma > (\theta/k)$ such that

$$(2.2) \quad 2NF(kd) - (N-2)df(d) \geq 0 \quad \text{for } d \geq \gamma.$$

Now for each real number d , the initial value (1.3), (1.4), $u(0) = d$ has a unique solution $u(t, d, \lambda)$. This solution depends continuously on (d, λ) in the sense that if $\{(d_n, \lambda_n)\} \rightarrow (d, \lambda)$, then $\{u(\cdot, d_n, \lambda_n)\}$ converges uniformly to $u(\cdot, d, \lambda)$ on $[0, 1]$. To see this, we observe that for each (d, λ) the map

$$(2.3) \quad u(s) \rightarrow d + \lambda \int_0^s t^{-n} \int_0^t r^n (-f(u(r))) dr$$

defines a contraction on $C([0, \varepsilon], R)$ for ε small enough.

Next given $d \in R$, $\lambda \in R$, we define

$$(2.4) \quad E(t, d, \lambda) = \frac{(u'(t, d, \lambda))^2}{2} + \lambda F(u(t, d, \lambda)),$$

$$(2.5) \quad H(t, d, \lambda) = tE(t, d, \lambda) + \frac{N-2}{2}u(t, d, \lambda)u'(t, d, \lambda).$$

Multiplying (1.3) by $r^N u'$ and integrating over $[\hat{t}, t]$, and then multiplying (1.3) by $r^n u$ and integrating over $[\hat{t}, t]$, we obtain

$$(2.6) \quad \begin{aligned} t^{N-1}H(t, d, \lambda) &= \hat{t}^{N-1}H(\hat{t}, d, \lambda) + \int_{\hat{t}}^t r^n \lambda [NF(u(r, d, \lambda)) \\ &\quad - \frac{N-2}{2}f(u(r, d, \lambda))u(r, d, \lambda)] dr. \end{aligned}$$

This identity is a form of "Pohozaev identity." For more details see Castro and Kurepa [1] and Pucci and Serrin [5].

Further, for $d \geq \gamma$ let $t_0 := t_0(d, \lambda)$ be such that $d \geq u(t_0, d, \lambda) \geq kd$ for all $t \in [0, t_0)$ and $u(t_0, d, \lambda) = kd$. Multiplying by r^n (1.3)–(1.4) and $u(0) = d$ gives $u'(t, d, \lambda) = -\lambda t^{-n} \int_0^t r^n f(u(r, d, \lambda)) dr$. Hence $-\lambda t f(kd) \geq Nu'(t, d, \lambda) \geq -\lambda t f(d)$, and integrating on $[0, t_0]$ we have

$$(2.7) \quad C_1 \{d/(\lambda f(kd))\}^{1/2} \geq t_0 \geq C_1 \{d/(\lambda f(d))\}^{1/2}$$

where $C_1 = \{(1-k)2N\}^{1/2} > 0$. Also choosing $\hat{t} = 0$, $t = t_0$, (2.6) gives

$$(2.8) \quad \begin{aligned} t_0^n H(t_0, d, \lambda) &= \lambda \int_0^{t_0} r^n \{NF(u(r, d, \lambda)) \\ &\quad - [(N-2)/2]f(u(r, d, \lambda))u(r, d, \lambda)\} dr \\ &\geq \lambda \int_0^{t_0} r^n \{NF(kd) - [(N-2)/2]f(d)d\} dr \\ &\geq \lambda \{NF(kd) - [(N-2)/2]f(d)d\} t_0^N / N \\ &\geq \lambda \{NF(kd) - [(N-2)/2]f(d)d\} \cdot \{C_1^N / N\} \cdot \{d/(\lambda f(d))\}^{N/2} \\ &= C_2 \lambda^{(1-N/2)} \{F(kd) - [(N-2)/(2N)]df(d)\} \cdot \{d/f(d)\}^{N/2} \end{aligned}$$

where $C_2 = (C_1)^N > 0$.

3. MAIN LEMMAS AND PROOF OF THEOREM 1.1

Lemma 3.1. *If $\lambda \in (0, \lambda_1 := N(\gamma - \beta)/f(\gamma))$, then $u(t, \gamma, \lambda) \geq \beta$ for all $t \in [0, 1]$.*

Proof. Let $t_1 := \sup\{t \leq 1; u(r, \gamma, \lambda) \geq \beta, \text{ for all } r \in (0, t)\}$. Since $u'(t, \gamma, \lambda) = -\lambda t^{-n} \int_0^t s^n f(u(s, \gamma, \lambda)) ds$, u is decreasing on $[0, t_1]$. Also if $\lambda \in (0, \lambda_1)$, $t \in [0, t_1]$, we have

$$|u'(t, \gamma, \lambda)| \leq \lambda t f(\gamma)/N < \gamma - \beta.$$

Hence $u(t_1, \gamma, \lambda) > \gamma - (\gamma - \beta)t_1$. In particular, if $t_1 < 1$ this gives $u(t_1, \gamma, \lambda) > \beta$ contradicting the definition of t_1 . Thus $t_1 = 1$ and the lemma is proven.

Lemma 3.2. *There exists $\lambda_2 > 0$ such that if $\lambda \in (0, \lambda_2)$, then $\{u(t, d, \lambda)\}^2 + \{u'(t, d, \lambda)\}^2 > 0$ for $t \in [0, 1]$, $d \in [\gamma, +\infty)$.*

Proof. Now for $t \geq t_0$, (2.6) and (2.8) gives

$$(3.1) \quad \begin{aligned} t^n H(t) &\geq C_2 \lambda^{(1-N/2)} \{Fkd - [(N-2)/(2N)]df(d)\} \{d/f(d)\}^{N/2} \\ &\quad + \lambda \int_{t_0}^t r^n \{NF(u(r, d, \lambda)) - [(N-2)/2]f(u(r, d, \lambda))u(r, d, \lambda)\} dr. \end{aligned}$$

Now by (1.8), our definition of $f(x)$ for $x < 0$ and the fact that $f(0) < 0$, there exists a constant $B < 0$ such that $G(s) = NF(s) - [(N-2)/2]f(s)s \geq B$ for all s . Further using (1.8), we may assume without loss of generality that γ is large enough so that $\{F(kd) - [(N-2)/(2N)]df(d)\} \{d/f(d)\}^{N/2} \geq 1$ for $d \geq \gamma$. Hence by (3.1) we have, for $t \in [t_0, 1]$,

$$(3.2) \quad \begin{aligned} t^n H(t) &\geq C_2 \lambda^{(1-N/2)} \{F(kd) - [(N-2)/(2N)]df(d)\} \cdot \{d/f(d)\}^{N/2} \\ &\quad + \lambda B \{t^N - t_0^N\}/N \\ &\geq C_2 \lambda^{(1-N/2)} + \lambda B/N \\ (3.3) \quad &= \lambda \{C_2 \lambda^{-N/2} + B/N\}. \end{aligned}$$

That is, there exists λ_2 such that for $\lambda \in (0, \lambda_2)$, $H(t)$ (and hence $[u(t, \lambda, d)]^2 + [u'(t, \lambda, d)]^2$) is positive for every $t \in [0, 1]$ and every $d \in [\gamma, +\infty)$ and the lemma is proven.

Lemma 3.3. *Given any $\lambda > 0$, there exists $d > \gamma$ such that $u(t, d, \lambda) < 0$ for some $t \in [0, 1]$.*

Proof. Let $\rho > 0$ and ω be such that $\omega'' + (n/t)\omega' + \rho\omega = 0$, $\omega(0) = 1$, $\omega'(0) = 0$ and the first zero of ω is $\frac{1}{\lambda}$. By (1.6), there exists $d_0(\lambda) \geq \theta/k$ such that if $x \geq d_0$ then

$$(3.4) \quad (f(x)/x) \geq (\rho/\lambda).$$

Suppose now that for every $d > \gamma$, $u(t, d, \lambda) \geq 0$ for all $t \in [0, 1]$. First we show that there exists $d_1(\lambda) \geq d_0(\lambda)$ such that for $d > d_1(\lambda)$ and $t_1 \in (0, 1]$

$$(3.5) \quad \text{if } u'(t_1, d, \lambda) = 0 \text{ then } u''(t_1, d, \lambda) < 0.$$

In fact, let $d_1(\lambda) > d_0(\lambda)$ be such that if $d > d_1(\lambda)$ then $t^n H(t) > 0$ for all $t \in [t_0, 1]$ (see (3.2)). Suppose there exists $t_1 \in [0, 1]$ with $u'(t_1, d, \lambda) = 0$, $u''(t_1, d, \lambda) \geq 0$. By (1.3) we have $u(t_1, d, \lambda) \leq \beta$. Since $kd > \beta$ we have $t_1 > t_0$. Thus $t_1^n H(t_1) = t_1^N \lambda F(u(t_1, d, \lambda)) < 0$, which contradicts the definition of $d_1(\lambda)$. Thus (3.5) holds. By (3.5) we have that if $d > d_1(\lambda)$ and $t \in [0, 1]$ then $u(t, d, \lambda) \cdot u'(t, d, \lambda) \leq 0$. Hence, by (3.2) and the fact that $E(t, d, \lambda) \geq \lambda F(kd)$ for $t \in [0, t_0]$, there exists $d_2(\lambda) > d_1(\lambda)$ such that for $d > d_2$

$$(3.6) \quad E(t, d, \lambda) \geq \lambda F(d_0) + 2d_0^2$$

for all $t \in [0, 1]$. Now let $d > d_2$. Since $(d\omega)'' + (n/r)(d\omega)' + \rho(d\omega) = 0$ and $u'' + (n/r)u' + \lambda(f(u)/u)u = 0$, we get

$$(3.7) \quad u(t)\{t^n v'(t)\} - v(t)\{t^n u'(t)\} = \int_0^t s^n \left\{ \frac{\lambda f(u/s)}{u(s)} - \rho \right\} ds$$

where $v = d\omega$. Hence if $u(t, d, \lambda) \geq d_0$ for all $t \in [0, \frac{1}{4}]$ then by (3.4) and facts that $v(\frac{1}{4}) = 0$, $v'(\frac{1}{4}) < 0$ we obtain a contradiction to (3.6). Thus there exists $t^* \in (0, \frac{1}{4})$ such that

$$(3.8) \quad u(t^*, d, \lambda) = d_0.$$

Also u is decreasing on $(0, t^*)$ and (3.6) implies $u'(t^*, d, \lambda) \leq -2d_0$. Since we are assuming $u(t, d, \lambda) \geq 0$ for all $t \in (t^*, 1]$, we have $u' \leq 0$ for $t \in [t^*, 1]$. Therefore $0 \leq u(t, d, \lambda) \leq d_0$ for $t \in (t^*, 1]$ and by (3.6) we have $u'(t, d, \lambda) \leq -2d_0$ for all $t \in (t^*, 1]$. Hence integrating we have

$$u(t^* + \frac{1}{2}, d, \lambda) - d_0 \leq -2d_0 \cdot (\frac{1}{2}),$$

that is, $u(t^* + \frac{1}{2}, d, \lambda) \leq 0$ where $t^* + \frac{1}{2} < 1$, with $u'(t^* + \frac{1}{2}, d, \lambda) \leq -2d_0$. Thus there exists $T \in (0, 1)$ such that $u(T, d, \lambda) < 0$ which is a contradiction, hence the lemma is proven.

Proof of Theorem 1.1. Let $\lambda_0 = \min\{\lambda_1, \lambda_2\}$ and $\lambda \in (0, \lambda_0)$. Let $\hat{d}(\lambda) := \hat{d} = \sup\{d \in [\gamma, +\infty); u(t, d, \lambda) \geq 0 \text{ for all } t \in [0, 1]\}$. By Lemma 3.3 we have that $\hat{d} < +\infty$. Now we claim that:

- (A) $u(1, \hat{d}, \lambda) = 0$,
- (B) $u(t, \hat{d}, \lambda) > 0 \forall t \in [0, 1)$,
- (C) $u'(1, \hat{d}, \lambda) < 0$, and
- (D) u is decreasing on $[0, 1]$.

Suppose there exists $T_1 < 1$ such that $u(T_1, \hat{d}, \lambda) = 0$. Then Lemma 3.2 gives $u'(T_1, \hat{d}, \lambda) \neq 0$ and without loss of generality we can assume $u'(T_1, \hat{d}, \lambda) < 0$. Thus there exists $T_2 \in (T_1, 1)$ such that $u(T_2, \hat{d}, \lambda) < 0$, a contradiction to the definition of $\hat{d}$. This proves (B). That is, $u(1, \hat{d}, \lambda) \geq 0$. Suppose $u(1, \hat{d}, \lambda) > 0$. Then there exists $\eta > 0$ such that $u(t, \hat{d}, \lambda) \geq \eta$ for all $t \in [0, 1]$. Thus there exists $\delta > 0$ such that $u(t, \hat{d} + \delta, \lambda) \geq \eta/2$ for all $t \in [0, 1]$, which contradicts the definition of $\hat{d}$. Hence $u(1, \hat{d}, \lambda) = 0$ and

(A) is proven. Finally (C) follows from Lemma 3.2 and (D) follows by Gidas, Ni and Nirenberg [7].

Remark. Unlike the case $N = 1$ (see Castro and Shivaji [1987, Theorem 1.2]), for $N \geq 2$ the problem (1.1)–(1.2) does not have non-negative solutions with interior zeros. This follows because if there exists $t_0 \in (0, 1)$ for which $u'(t_0) = u(t_0) = 0$ then $E(t_0) = 0$. By (1.3) we obtain $dE/dt = -n(u')^2/t \leq 0$. But $E(1) \geq 0$. Thus $E = 0$ for all $t \in [t_0, 1]$ which is possible only if $u' = 0$ and hence $u = 0$ for all $t \in [t_0, 1]$. But from (1.3) we see that this is impossible with $f(0) < 0$.

REFERENCES

1. A. Castro and A. Kurepa, *Energy analysis of a nonlinear singular differential equation and applications*, Rev. Colombiana Mat. (to appear).
2. A. Castro and A. Kurepa, *Infinitely many solutions to a superlinear Dirichlet problem in a ball*, Proc. Amer. Math. Soc. **101** (1) (1987), 57–64.
3. A. Castro and R. Shivaji, *Non-negative solutions for a class of non-positone problems*, Proc. Royal Soc. Edinburgh Sect. A **108**, (1988) 291–302.
4. W. Mo Ni and J. Serrin, *Non-existence theorems for quasi linear partial differential equations*, Rend. Circ. Mat. Palermo (5) Suppl (1986).
5. P. Pucci and J. Serrin, *A general variational identity*, Indiana Univ. Math. J. **35** (3) (1986), 681–703.
6. J. Smoller and A. Wasserman, *Existence of positive solutions for semilinear elliptic equations in general domains*, Arch. Rational Mech. Anal. **98** (3) (1987), 229–249.
7. B. Gidas, W. M. Ni and L. Nirenberg, *Symmetry and related properties via the maximum principle*, Comm. Math. Phys. **68** (1979), 209–243.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF NORTH TEXAS, DENTON, TEXAS 76203-5116

DEPARTMENT OF MATHEMATICS, MISSISSIPPI STATE UNIVERSITY, MISSISSIPPI STATE, MISSISSIPPI 39762