A WEAK-TYPE ESTIMATE FOR FOURIER-BESSEL MULTIPLIERS

JOHN GOSSELIN AND KRZYSZTOF STEMPAK

(Communicated by Richard R. Goldberg)

ABSTRACT. We apply Hörmander's technique to prove a weak-type $(1, 1)$ estimate for multiplier operators with respect to the Fourier-Bessel transform. This improves a result in [4, 5].

1. INTRODUCTION

Consider the measure space $(\mathbf{R}^+, d\mu(x))$ where $d\mu(x) = x^r dx$, r being a fixed positive real number. By $L^p(\mu)$, $1 \leq p \leq \infty$, we denote the corresponding Lebesgue spaces equipped with the norms

$$\|f\|_p = \left(\int_0^\infty |f|^p d\mu \right)^{1/p}.$$

Let $\hat{f}(\lambda)$, $\lambda \geq 0$, denote the Fourier-Bessel transform of the function $f \in L^1(\mu)$. Explicitly this means

$$\hat{f}(\lambda) = \int_0^\infty f(x) \phi_\lambda(x) d\mu(x),$$

where $\phi_\lambda(x) = a(r)(\lambda x)^{-(r-1)/2} J_{(r-1)/2}(\lambda x)$, $x \geq 0$, $a(r) = 2^{(r-1)/2} \Gamma((r+1)/2)$, and J_α denotes the Bessel function of the first kind of order α . The functions ϕ_λ , $\lambda \geq 0$, are eigenfunctions of the second order Bessel differential operator

$$L = - \left(\frac{d}{dx^2} + \frac{r}{x} \frac{d}{dx} \right).$$

More precisely, we have

$$(1.1) \quad L\phi_\lambda = \lambda^2 \phi_\lambda.$$

If $r = n - 1$, $n \geq 2$ being an integer, then

$$\hat{f}(\|\bar{\lambda}\|) = \frac{1}{2} \pi^{-n/2} \Gamma\left(\frac{n}{2}\right) \int_{\mathbf{R}^n} f(\|\bar{x}\|) \exp(-i\langle \bar{\lambda}, \bar{x} \rangle) d\bar{x}$$

Received by the editors May 22, 1987.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 42B15; Secondary 42C15.

Key words and phrases. Fourier-Bessel transform, multipliers, generalized convolution.

© 1989 American Mathematical Society
0002-9939/89 \$1.00 + \$.25 per page

for $\bar{\lambda} \in \mathbf{R}^n$. Thus $\hat{f}$ gives the radial part of the Fourier transform of the radial function $f(\|\bar{x}\|)$ on $\mathbf{R}^n$ (observe that $f \in L^1(\mu)$ if and only if $f(\|\bar{x}\|) \in L^1(\mathbf{R}^n, d\bar{x})$ where $\|\cdot\|$ denotes the Euclidean norm and $d\bar{x}$ is Lebesgue measure on $\mathbf{R}^n$). In the case $r = n - 1$, one can easily see that for the Fourier-Bessel transform, both the inversion theorem and Plancherel's formula hold. In particular we have

$$f(x) = a(r)^{-1} \int_0^\infty \hat{f}(\lambda) \phi_\lambda(x) d\mu(\lambda)$$

almost everywhere on $\mathbf{R}^+$ providing $f \in L^1(\mu)$ and $\hat{f} \in L^1(\mu)$. For $f \in L^2(\mu)$ we have

$$\int_0^\infty |f(x)|^2 d\mu(x) = a(r)^{-2} \int_0^\infty |\hat{f}(\lambda)|^2 d\mu(\lambda).$$

The above results for general $r > 0$ can be established via a summability argument similar to that in the case of the ordinary Fourier transform. The key idea is that the Gaussian is invariant under the Fourier-Bessel transform when appropriately normalized. We also note that in the case $r = n - 1$, the operator $-L$ is the radial part of the ordinary Laplace operator on $\mathbf{R}^n$. Finally, we mention that C will be used to denote a constant which may vary from line to line.

For any bounded function m on $(0, \infty)$ we define the multiplier operator T_m by $(T_m f)^\wedge = m \hat{f}$. The aim of this note is to prove the following theorem which is a generalization of the celebrated Hörmander-Mihlin multiplier theorem for the Fourier transform (see [3]).

Theorem 1.1. *Let k denote the least even integer $> (r + 1)/2$ and let $m \in C^k(0, \infty)$ be a bounded function which satisfies*

$$(1.2) \quad \left(\int_{R/2}^R |m^{(s)}(\lambda)|^2 d\mu(\lambda) \right)^{1/2} \leq B R^{(r+1)/2-s},$$

where B is a constant independent of $R > 0$ and $s = 0, 1, \dots, k$. Then the operator T_m is of weak-type $(1, 1)$, i.e.

$$(1.3) \quad \mu(\{x \in (0, \infty) : |T_m f(x)| > \alpha\}) \leq C \alpha^{-1} \|f\|_1$$

where C is independent of $\alpha > 0$ and $f \in L^1(\mu)$.

Using the interpolation theorem of Marcinkiewicz and duality we also obtain the following

Corollary 1.2. *Let m satisfy the assumptions of the Theorem 1.1. Then for $1 < p < \infty$, T_m is of strong type (p, p) , i.e. $\|T_m f\|_p \leq C_p \|f\|_p$.*

Remark. The result of the above corollary was obtained previously by the second author by means of a Littlewood-Paley theory for the Fourier-Bessel transform (see [4, 5]). The proof of the above theorem is more direct and is based upon a generalized convolution structure which is natural for the Fourier-Bessel

transform. The novel ingredient of our proof is a Bernstein-type inequality for this convolution structure. We establish this inequality in the following section.

2. A BERNSTEIN-TYPE INEQUALITY FOR GENERALIZED TRANSLATIONS

Since Hirschman and Bochner considered the Banach algebra of radial functions over thirty years ago, it has been known that $L^1(\mu)$ admits a commutative Banach algebra structure for which the Fourier-Bessel transform becomes the Gelfand transform (see [4 and 5] for details and for other references). In order to describe this structure we define the generalized translation operator T^y , $y \geq 0$, for suitable functions f , by

$$(2.1) \quad T^y f(x) = \int_0^\pi f((x, y)_\theta) d\nu(\theta)$$

where $(x, y)_\theta = (x^2 + y^2 - 2xy \cos \theta)^{1/2}$, $0 \leq \theta \leq \pi$, $x, y \geq 0$ and $d\nu(\theta)$ denotes the probability measure $b(r)(\sin \theta)^{r-1} d\theta$ on $[0, \pi]$. Here

$$b(r) = \pi^{-1/2} \Gamma((r+1)/2) \Gamma(r/2)^{-1}.$$

The operator T^y can also be described by

$$(2.2) \quad T^y f(x) = \int_{|x-y|}^{x+y} f(z) dW_{x,y}(z)$$

where the probability measure $dW_{x,y}(z)$ is supported on $[|x-y|, x+y]$ and given by

$$dW_{x,y}(z) = c(r) \frac{\Delta(x, y, z)^{r-2}}{(xyz)^{r-1}} d\mu(z),$$

where $c(r) = 2^{r-2} \Gamma((r+1)/2) \Gamma(r/2)^{-1} \pi^{-1/2}$. In the above formula $x, y, z \geq 0$, $|x-y| \leq z \leq x+y$, and $\Delta(x, y, z)$ denotes the area of a triangle with sides x, y, z . It is quite straightforward to go from (2.2) to (2.1) by a change of variables. Next, it is easy to check that the measure $dW_{x,y}(z) d\mu(y)$ is symmetric with respect to y and z , i.e. $dW_{x,y}(z) d\mu(y) = dW_{x,z}(y) d\mu(z)$. From this it follows that the operators T^y , $y \geq 0$, are selfadjoint on $L^2(\mu)$ and moreover that

$$(2.3) \quad \int_0^\infty f(x) T^y g(x) d\mu(x) = \int_0^\infty T^y f(x) g(x) d\mu(x),$$

for any reasonable pair of functions f and g , e.g. if $f \in L^\infty(\mu)$ and $g \in L^1(\mu)$. It is also easy to see that generalized translations are submarkovian contractions, i.e. $0 \leq f \leq 1$ implies $0 \leq T^y f \leq 1$ and $\|T^y f\|_p \leq \|f\|_p$, for $1 \leq p \leq \infty$. We now define the generalized convolution $f * g$ for $f, g \in L^1(\mu)$ by

$$(2.4) \quad f * g(x) = \int_0^\infty f(y) T^y g(x) d\mu(y).$$

It is well known that the following identity involving the functions ϕ_λ and the generalized translations T^ν is valid

$$(2.5) \quad T^\nu \phi_\lambda(x) = \phi_\lambda(x) \phi_\lambda(y)$$

(see [4, 5] for references). Using (2.3) and (2.5) one can easily verify that for any $f, g \in L^1(\mu)$ we have $(f * g)^\wedge = \hat{f} \hat{g}$ and $(T^\nu f)^\wedge(\lambda) = \phi_\lambda(y) \hat{f}(\lambda)$. Therefore

$$(2.6) \quad T^\nu g * f = g * T^\nu f.$$

It is also easy to check that with the pseudodistance $\rho(x, y) = |x - y|$, the space $(\mathbf{R}^+, \mu, \rho)$ becomes a space of homogeneous type in the sense of [1]. Finally, from (2.2) it is immediate that $T^\nu f(x) = 0$ if f vanishes on the interval $(|x - y|, x + y)$ and that $|T^\nu f(x)| \leq T^\nu(|f|)(x)$ for $x, y \geq 0$.

In the proof of the Hörmander-Mihlin theorem (cf. [3]), Bernstein's inequality plays an important role. We now prove a Bernstein-type inequality for the generalized translations T^ν , $y \geq 0$.

Theorem 2.1. *Suppose $h \in L^1(\mu)$ is a differentiable function on $\mathbf{R}^+$ with $h'(x) \in L^1(\mu)$. Then*

$$(2.7) \quad \|T^{y_1} h - T^{y_2} h\|_1 \leq \|h'\|_1 |y_1 - y_2|.$$

Proof. Using (2.1) we write

$$(2.8) \quad \begin{aligned} \|T^{y_1} h - T^{y_2} h\|_1 &= \int_0^\infty \left| \int_0^\pi [h((x, y_1)_\theta) - h((x, y_2)_\theta)] d\nu(\theta) \right| d\mu(x) \\ &\leq \int_0^\infty \int_0^\pi |h((x, y_1)_\theta) - h((x, y_2)_\theta)| d\nu(\theta) d\mu(x) \\ &= \int_0^\infty \int_0^\pi \left| \int_0^1 \frac{d}{ds} (h((x, y_2 + s(y_1 - y_2))_\theta)) ds \right| d\nu(\theta) d\mu(x). \end{aligned}$$

For fixed $x, y_1, y_2 \geq 0$ and $\theta \in (0, \pi]$, let

$$\Phi(s) \equiv \Phi_{x, \theta}^{y_1, y_2}(s) = (x, y_2 + s(y_1 - y_2))_\theta, \quad s \in [0, 1].$$

Then

$$(2.9) \quad \left| \frac{d}{ds} \Phi(s) \right| \leq |y_1 - y_2|$$

independently of x, y_1, y_2, θ . To verify (2.9) consider the vectors $X = (x, 0)$, $Y_i = (y_i \cos \theta, y_i \sin \theta)$, $i = 1, 2$, in $\mathbf{R}^2$. Letting $\|\cdot\|$ denote the Euclidean norm on $\mathbf{R}^2$ it is easy to see that

$$(x, y_2 + s(y_1 - y_2))_\theta = \|Y_2 + s(Y_1 - Y_2) - X\| \equiv \|\xi_s\|.$$

Since $\nabla(\|\cdot\|)(\xi) = \xi/\|\xi\|$ for $\xi \in \mathbf{R}^2$, Schwarz' inequality implies

$$\begin{aligned} \left| \frac{d}{ds} \Phi(s) \right| &= \left| \frac{d}{ds} (\|Y_2 + s(Y_1 - Y_2) - X\|) \right| = \left| \left\langle \frac{\xi_s}{\|\xi_s\|}, Y_1 - Y_2 \right\rangle \right| \\ &\leq \|Y_1 - Y_2\| = |y_1 - y_2|. \end{aligned}$$

Here $\langle \cdot, \cdot \rangle$ denotes the ordinary Euclidean inner product on $\mathbf{R}^2$. This gives (2.9). Now, using (2.9) and the contraction property of generalized translations, we have from (2.8)

$$\begin{aligned} & \|T^{y_1}h - T^{y_2}h\|_1 \\ & \leq |y_1 - y_2| \int_0^1 \int_0^\infty \int_0^\pi |h'((x, y_2 + s(y_1 - y_2))_\theta)| d\nu(\theta) d\mu(x) ds \\ & \leq |y_1 - y_2| \int_0^1 \|T^{y_1+s(y_1-y_2)}(|h'|)\|_1 ds \\ & \leq |y_1 - y_2| \|h'\|_1. \end{aligned}$$

This completes the proof of Theorem 2.1.

As in the case of the Fourier transform (cf. [2]) Theorem 2.1 implies the following corollary. We include its proof for the sake of completeness.

Corollary 2.2. *There exists a constant $C > 0$ such that for any $\lambda > 0$ and any function $f \in L^1(\mu)$ with $\text{supp } \hat{f} \subset (0, \lambda)$*

$$(2.10) \quad \|T^{y_1}f - T^{y_2}f\|_1 \leq C\lambda|y_1 - y_2| \|f\|_1.$$

Proof. Choose $\chi \in C^\infty(\mathbf{R}^+)$ such that $\chi(\lambda) = 1$ for $0 < \lambda \leq 1$ and $\chi(\lambda) = 0$ for $\lambda \geq 2$. Then by the inversion theorem we have $\chi = \hat{h}$ for some C^∞ function $h \in L^1(\mu)$ with $h' \in L^1(\mu)$. Let $h_\varepsilon(x) = \varepsilon h(\varepsilon x)$ for $\varepsilon > 0$. Then $\hat{h}_\varepsilon(\lambda) = \chi(\lambda/\varepsilon) = 1$ for $0 \leq \lambda \leq \varepsilon$. Thus, for $\varepsilon = \lambda$ we have

$$T^{y_1}f - T^{y_2}f = h_\lambda * (T^{y_1}f - T^{y_2}f) = (T^{y_1}h_\lambda - T^{y_2}h_\lambda) * f$$

and therefore by (2.7) we have

$$\begin{aligned} \|T^{y_1}f - T^{y_2}f\|_1 & \leq \|T^{y_1}h_\lambda - T^{y_2}h_\lambda\|_1 \|f\|_1 = \|T^{\lambda y_1}h - T^{\lambda y_2}h\|_1 \|f\|_1 \\ & \leq \lambda|y_1 - y_2| \|h'\|_1 \|f\|_1. \end{aligned}$$

This establishes (2.10) with $C = \|h'\|_1$ and completes the proof of the corollary.

3. PROOF OF THEOREM 1.1

We closely follow the original proof of Hörmander (see [3, pp. 121–123]). Choose a function $\psi \in C_0^\infty(\mathbf{R}^+)$ with support in $(\frac{1}{2}, 2)$ such that $\sum_{j=-\infty}^\infty \psi(2^{-j}\lambda) = 1$, $\lambda > 0$, and let

$$(3.1) \quad m_j(\lambda) = m(\lambda)\psi(2^{-j}\lambda) \equiv m(\lambda)\psi_j(\lambda)$$

and

$$(3.2) \quad k_j(x) = \check{m}_j(x) = a(r)^{-1} \int_0^\infty m_j(\lambda)\phi_\lambda(x)\lambda^r d\lambda.$$

Then $T_m = \sum_{j=-\infty}^\infty T_{m_j}$ where $T_{m_j}f = k_j * f$ and $k_j \in L^1(\mu)$. In order to prove the weak-type $(1, 1)$ inequality (1.3), it suffices to establish (see [1, p. 75])

$$(3.3) \quad \sum_{j=-\infty}^\infty \int_{|x-y_0|>2|y-y_0|} |T^y k_j(x) - T^{y_0} k_j(x)| d\mu(x) \leq C,$$

where C is independent of y , $y_0 \geq 0$. This will be implied in a standard way by the following estimates:

$$(3.4) \quad \int_{|x-y_0| \geq 2|y-y_0|} |T^y k_j(x) - T^{y_0} k_j(x)| d\mu(x) \leq C(2^j |y - y_0|)^{(r+1)/2-k},$$

which is good when $2^j |y - y_0| \geq 1$, and

$$(3.5) \quad \int_{|x-y_0| \geq 2|y-y_0|} |T^y k_j(x) - T^{y_0} k_j(x)| d\mu(x) \leq C2^j |y - y_0|,$$

which is good when $2^j |y - y_0| < 1$.

To establish (3.4) we first note that conditions (1.2) imply

$$(3.6) \quad \left(\int_0^\infty |L^l m_j(\lambda)|^2 d\mu(\lambda) \right)^{1/2} \leq CB(2^j)^{(r+1)/2-2l}, \quad l = 0, 1, \dots, k/2.$$

In fact, by induction, one can easily verify the following Leibniz rule for L : for $f, g \in C^{2l}(\mathbf{R}^+)$

$$L^l(fg) = \sum_{0 < \alpha + \beta \leq 2l} C_{\alpha, \beta}^l f^{(\alpha)} g^{(\beta)} \lambda^{\alpha + \beta - 2l}.$$

Thus, taking $R = 2^j$ in (1.2), we can estimate the left-hand side of (3.6) by

$$\begin{aligned} & C \sum_{0 < \alpha + \beta \leq 2l} \left(\int_0^\infty |m^{(\alpha)}(\lambda) \psi_j^{(\beta)}(\lambda) \lambda^{\alpha + \beta - 2l}|^2 d\mu(\lambda) \right)^{1/2} \\ & \leq C \sum_{0 < \alpha + \beta \leq 2l} 2^{-\beta j} \left(\int_{2^{j-1}}^{2^{j+1}} |m^{(\alpha)}(\lambda) \lambda^{\alpha + \beta - 2l}|^2 d\mu(\lambda) \right)^{1/2} \\ & \leq C \sum_{0 < \alpha + \beta \leq 2l} (2^j)^{\alpha - 2l} \left(\int_{2^{j-1}}^{2^{j+1}} |m^{(\alpha)}(\lambda)|^2 d\mu(\lambda) \right)^{1/2} \\ & \leq CB(2^j)^{(r+1)/2-2l}. \end{aligned}$$

This gives (3.6). Next, using the Plancherel theorem and (3.6) we obtain

$$\begin{aligned} \|(1 + (2^j x)^2)^{k/2} k_j\|_2 &= \|[(1 + 2^{2j} L)^{k/2} m_j]^\vee\|_2 \\ &= a(r)^{-2} \|(1 + 2^{2j} L)^{k/2} m_j\|_2 \\ &\leq C \sum_{l=0}^{k/2} (2^{2j})^l \|L^l m_j\|_2 \leq CB(2^j)^{(r+1)/2}. \end{aligned}$$

Thus by Schwarz' inequality and the estimate $\|(1 + (2^j x)^2)^{-k/2}\|_2 \leq C(2^j)^{-(r+1)/2}$ we obtain

$$(3.7) \quad \|k_j\|_1 \leq \|(1 + (2^j x)^2)^{k/2} k_j\|_2 \|(1 + (2^j x)^2)^{-k/2}\|_2 \leq CB.$$

Moreover, since $\|(2^j x)^k k_j\|_2 \leq \|(1 + (2^j x)^2)^{k/2} k_j\|_2$ we obtain in the same way

$$(3.8) \quad \int_{x \geq t} |k_j(x)| d\mu(x) \leq \|(2^j x)^k k_j\|_2 \left(\int_{x \geq t} (2^j x)^{-2k} d\mu(x) \right)^{1/2} \\ \leq CB(2^j t)^{(r+1)/2-k}.$$

If χ_A is the characteristic function of a set $A \subset \mathbf{R}^+$, then $0 \leq T^y \chi_A(x) \leq 1$ for $x, y \in \mathbf{R}^+$. We claim that

$$(3.9) \quad T^{y_0} \chi_{\{z: |z-y_0| > 2|y-y_0|\}} \leq \chi_{\{z: z > 2|y-y_0|\}}$$

and

$$(3.10) \quad T^y \chi_{\{z: |z-y_0| > 2|y-y_0|\}} \leq \chi_{\{z: z > |y-y_0|\}}.$$

We only prove (3.9) and note that the proof of (3.10) is similar. Let E denote the set $\{z: |z-y_0| > 2|y-y_0|\}$. To verify (3.9) we need only check that $0 \leq x \leq 2|y-y_0|$ implies $T^{y_0} \chi_E(x) = 0$. Since $0 \leq x \leq 2|y-y_0|$ implies $(|y_0-x|, y_0+x) \cap E = \emptyset$, it follows that $T^{y_0} \chi_E(x) = 0$ (see remark after (2.6)). This proves (3.9).

Using (3.7) through (3.10) and the selfadjointness of generalized translations, cf. (2.3), we find that the left-hand side of (3.4) is majorized by

$$\int_0^\infty \chi_E(x) |T^y k_j(x)| d\mu(x) + \int_0^\infty \chi_E(x) |T^{y_0} k_j(x)| d\mu(x) \\ \leq \int_{|y-y_0|}^\infty |k_j(x)| d\mu(x) + \int_{2|y-y_0|}^\infty |k_j(x)| d\mu(x) \\ \leq CB(2^j |y-y_0|)^{(r+1)/2-k}.$$

Thus (3.4) is established.

Finally, to establish (3.5) we use Bernstein's inequality. Since $\hat{k}_j = m_j$ has support in $(0, 2^{j+1})$ it follows from Corollary (2.2) and (3.7) that

$$\int_{|x-y_0| \geq 2|y-y_0|} |T^y k_j - T^{y_0} k_j| d\mu \leq \|T^y k_j - T^{y_0} k_j\|_1 \\ \leq 2^{j+1} |y-y_0| \|k_j\|_1 \leq CB2^j |y-y_0|.$$

Thus (3.5) is established. Combining (3.4) and (3.5) we obtain (3.3) which completes the proof of the theorem.

4. A FINAL REMARK

In connection with the Bernstein-type inequality (2.7) the following question arises. Consider the "mean value" operators T^y , $y \geq 0$, defined for functions on $\mathbf{R}^n$, $n \geq 2$, by

$$T^y f(\bar{x}) = \int_{S^{n-1}} f(\bar{x} + y\bar{t}) d\sigma(\bar{t}),$$

where $d\sigma$ is normalized surface measure on the sphere S^{n-1} . In other words T^y is the convolution operator with the uniformly distributed probability measure on $\{\bar{x} \in \mathbb{R}^n: \|\bar{x}\| = y\}$. Is it true that for any $h \in C^1(\mathbb{R}^n)$ with, say, $\|\nabla h\| \in L^1(\mathbb{R}^n)$, the following estimate holds

$$\|T^{y_1}h - T^{y_2}h\|_{L^1(\mathbb{R}^n)} \leq C|y_1 - y_2|,$$

where C depends on h but not on y_1 and y_2 ? We note that Theorem 2.1 gives a positive answer to this problem only in the case when h is radial.

But one can rather easily remark that the above problem may be answered affirmatively by writing out $\|T^{y_1}h - T^{y_2}h\|_{L^1(\mathbb{R}^n)}$, interchanging the order of integration and then applying the ordinary Bernstein-type inequality on $\mathbb{R}^n$. This yields an easy proof of Theorem 2.1 in the case of integral values of r .

REFERENCES

1. R. Coifman and G. Weiss, *Analyse harmonique non commutative sur certains espaces homogènes*, Lecture Notes in Math., vol. 242, Springer-Verlag, Berlin and New York, 1971.
2. R. E. Edwards and G. I. Gaudry, *Littlewood-Paley and multiplier theory*, Springer-Verlag, Berlin and New York, 1977.
3. L. Hörmander, *Estimates for translation invariant operators in L^p spaces*, Acta Math. **104** (1960), 93–140.
4. K. Stempak, *La théorie de Littlewood-Paley pour la transformation de Fourier-Bessel*, C. R. Acad. Sci. Paris **303** (1986), 15–18.
5. —, *The Littlewood-Paley theory for the Fourier-Bessel transform*, Univ. of Wrocław, Preprint no. 45, 1985.

Current address (John Gosselin): DEPARTMENT OF MATHEMATICS, UNIVERSITY OF GEORGIA, ATHENS, GEORGIA 30602

Current address (Krzysztof Stempak): INSTITUTE OF MATHEMATICS, UNIVERSITY OF WROCLAW, PL. GRUNWALDZKI 2/4, 50-384 WROCLAW, POLAND