A NEW EXAMPLE IN K -THEORY OF LOOPSPACES

TAHSIN GHAZAL

(Communicated by Frederick R. Cohen)

ABSTRACT. The “Eilenberg-Moore” type spectral sequences which connect $K^*(\Omega X)$ and $K^*(X)$ have well-known bad properties, when, for example, $X = K(\mathbb{Z}/p, 2)$. This paper shows that the result can be as bad when X is a finite complex.

The “Eilenberg-Moore” type spectral sequences which connect $K^*(\Omega X)$ and $K^*(X)$ have well-known bad properties, such as when $X = K(\mathbb{Z}/p, 2)$ [4]. The purpose of this note is to show that the spectral sequence properties can be just as bad when X is a finite complex. Specifically, we prove

Theorem 1. *For any odd prime p , there is a simply connected finite complex X such that*

- (i) $\tilde{K}^*(X; \mathbb{Z}) = 0$ (integral K -theory)
- (ii) $\tilde{K}^*(\Omega X; \mathbb{Z})$ contains a direct summand isomorphic to $\mathbb{Z}/p$.

It follows (informally) that there is no way of deducing $K^*(\Omega X; \mathbb{Z})$ from the limit of the Eilenberg-Moore spectral sequence with $E_2 = \text{Tor}_{K^*(X; \mathbb{Z})}(\mathbb{Z}, \mathbb{Z})$, even when X is finite.

The example X is a very obvious one; the cofibre of the Adams map [1] in a suitable dimension. We describe this space X and prove that it has trivial K -theory in §1; in §2 we show that its loop space has nontrivial K -theory.

1. THE SPACE X

Let $p > 2$ be a prime and let $P^n(p)$ denote the Moore space $S^{n-1}U_p e^n$. In [1] J. F. Adams described stable maps

$$(1) \quad A: P^{n+2p-2}(p) \rightarrow P^n(p)$$

(n large) such that A^* is an isomorphism in K -theory. We define $X(n, p)$ to be the cofibre of A :

$$(2) \quad X(n, p) = P^n(p)U_A C(P^{n+2p-2}(p)).$$

Received by the editors August 25, 1988 and, in revised form, March 1, 1989.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 55N15; Secondary 19L20.

Key words and phrases. Adams map, K -theory.

Lemma 1. $\tilde{K}^*(X(n, p); \mathbb{Z}) = 0$.

Proof. The proof follows immediately from the cofibration exact sequence.

Now choose n so that $\tilde{K}^*(\Omega X; \mathbb{Z})$ is not 0. The construction is only possible when n has its minimum value n_0 , since for $n > n_0$, X is a suspension ΣY , with $\tilde{K}^*(Y; \mathbb{Z}) = 0$. By the theory of the James construction [5], $\Omega X = \Omega \Sigma Y$ also has trivial K -theory in this case.

Cohen and Neisendorfer [3] have computed the minimum value for n to be 3. We therefore set $X = X(3, p) = P^3(p)U_A C(P^{2p+1}(p))$. We have $H^*(X; \mathbb{Z})$ given by

$$(3) \quad \begin{aligned} H^3(X; \mathbb{Z}) &= H^{2p+2}(X; \mathbb{Z}) = \mathbb{Z}/p, \\ H^j(X; \mathbb{Z}) &= 0 \quad \text{otherwise.} \end{aligned}$$

2. THE K -THEORY OF ΩX

Proving that the K -theory of ΩX is nontrivial is simple. Using (3) and the transgression exact sequence of [6] gives

Lemma 2. $H^1(\Omega X; \mathbb{Z}) = 0$; $H^2(\Omega X; \mathbb{Z}) = \mathbb{Z}/p$.

But $H^2(\Omega X; \mathbb{Z})$ is a direct summand in $\tilde{K}^0(\Omega X; \mathbb{Z})$, as is well known. (The mapping $K(\mathbb{Z}, 2) = BU(1) \rightarrow BU$ is split by the determinant map.) This completes the proof of Theorem 1.

We can prove the analogue to Theorem 1 for K -theory mod p using the same space X ; a careful use of the universal coefficient theorem [2] will provide a $\mathbb{Z}/p$ summand in $K^0(\Omega X; \mathbb{Z}/p)$.

Of course the Adams maps are defined (in suitable degrees) if p is replaced by p^r :

$$A_r: P^{n+2(p-1)p^{r-1}}(p^r) \rightarrow P^n(p^r).$$

If we know the exact level of desuspension of A_r , we can expect a similar result for K -theory of the cofibre and its loop space.

REFERENCES

1. J. F. Adams, *On the groups $J(X)$* IV, *Topology* **5** (1966), 21–71.
2. D. W. Anderson, *Universal coefficient theorems for K -theory*, Unpublished (1963).
3. F. R. Cohen and J. A. Neisendorfer, *Note on desuspending the Adams map*, *Math. Proc. Cambridge Philos. Soc.*, **99** (1986), 59–64.
4. L. H. Hodgkin, *The equivariant Künneth theorem in K -theory*, *Lecture Notes in Math.*, no. 496, Springer, Berlin 1975, pp. 1–101.
5. I. M. James, *Reduced product spaces*, *Ann. of Math.* **62** (2) (1955), 170–197.
6. J. P. Serre, *Homologie singulière des espaces fibres*, *Ann. of Math.* **54** (2) (1951), 425–505.

MATHEMATICS DEPARTMENT, KING SAUD UNIVERSITY, P.O. BOX 2455, RIYADH 11451, SAUDI ARABIA