TWO CLASSES OF FRÉCHET-URYSOHN SPACES

ALAN DOW

(Communicated by Dennis Burke)

ABSTRACT. Arhangel'skii introduced five classes of spaces, α_i -spaces ($i < 5$), which are important in the study of products of Fréchet-Urysohn spaces. For each $i < 5$, each α_i -space is an α_{i+1} -space and it follows from the continuum hypothesis that there are countable α_{i+1} -spaces which are not α_i -spaces. A v -space (w -space) is a Fréchet-Urysohn α_1 -space (α_2 -space). We show that there is a model of set theory in which each α_2 -space (w -space) is an α_1 -space (v -space).

INTRODUCTION

Arhangel'skii defines a point $x \in X$ to be an α_1 -point (α_2 -point) if whenever F_n is a sequence converging to x , for each $n < \omega$, there is a sequence F converging to x such that $F_n - F$ is finite ($F_n \cap F$ is infinite) for each $n < \omega$. Furthermore a point is called an α_0 -point if it has a countable neighbourhood base. A space is called an α_1 -space if each point is an α_1 -point. A space is *Fréchet-Urysohn* if whenever a point is in the closure of a set there is a sequence from it converging to the point.

Nyikos has shown that there is a countable w -space which is not first countable [Ny1, Ny2]. In [Ny2] Nyikos asks if there is a countable w -space which is not a v -space and a countable v -space which is not first-countable. Nyikos [Ny2] produces examples of countable w -spaces which are not v -spaces from a special set-theoretic assumption following from, for example, Martin's Axiom (even $\underline{b} < \underline{d}$). Nogura [N] has shown that MA (even $\underline{b} = \underline{d}$) implies there is an example of a countable v -space which is not first countable. It is shown in [DS] that it is consistent that each countable v -space is first-countable.

Gruenhage [G] introduced w -spaces and Sharma [Sh] obtained their characterization in terms of α_2 -spaces. The term v -space seems to be due to Nyikos.

Received by the editors July 18, 1988 and, in revised form, December 2, 1988.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 54E35.

Key words and phrases. Fréchet-Urysohn space, v -space, w -space.

This research was supported by NSERC of Canada.

ω -SPLITTING FAMILIES FROM α_2 -POINTS

1. Definition. We say that $\mathfrak{X} \subset [\omega]^\omega$ is ω -splitting if for each countable family $\{A_n : n \in \omega\} \subset [\omega]^\omega$ there is an $X \in \mathfrak{X}$ such that $|A_n \cap X| = |A_n - X|$ for each $n \in \omega$. We shall say that X is M -splitting if $|A \cap X| = |A - X|$ for each $A \in M \cap [\omega]^\omega$.

Let x be a point in a space Y and suppose $F_n \in [Y]^\omega$ is a sequence converging to x for each $n \in \omega$. Identify $\bigcup F_n$ with ω and define $\mathfrak{X}$ to be the set of X which "think" that x is an α_1 -point. That is, define $\mathfrak{X}(x, \langle F_n \rangle) = \{X \in [\omega]^\omega : \exists F \subset Y \text{ } (F \text{ converges to } x \text{ and } (X \cap F_n) - F \text{ is finite for each } n \in \omega)\}$. The definition of $\mathfrak{X}(x, \langle F_n \rangle)$ of course depends on the identification of $\bigcup F_n$ with ω , but only up to a permutation on ω and this will never matter to us.

2. Lemma. If $F_n \in [Y]^\omega$ ($n \in \omega$) converges to an α_2 -point x , then $\mathfrak{X} = \mathfrak{X}(x, \langle F_n \rangle)$ is an ω -splitting family.

Proof. Assume that $\bigcup F_n$ is identified with ω and let $\{A_k : k \in \omega\} \subset [\omega]^\omega$. Let $I = \{j \in \omega : A_j \cap F_n \text{ is finite for each } n\}$. Choose for each n , a finite subset H_n of $F_n - \bigcup_{m < n} F_m$ so that $A_j - H$ is finite for each $j \in I$ where $H = \bigcup H_n$. Now for each $j \in \omega - I$, choose $n_j \in \omega$ so that $A_j \cap F_{n_j}$ is infinite. Next choose an infinite $B_j \subset A_j \cap F_{n_j}$ so that $B_j \cap B_k = \emptyset$ for $j \neq k$ in $\omega - I$. Now since x is an α_2 -point, there is a sequence F converging to x such that $F \cap B_j$ is infinite for each $j \in \omega - I$. It follows that $F \cup H$ hits each A_j in an infinite set and that $F \cup H \in \mathfrak{X}$. Finally any infinite subset of $F \cup H$ is a member of $\mathfrak{X}$ hence there is an $X \in \mathfrak{X}$ which splits $\{A_j : j \in \omega\}$.

3. Lemma. If every ω -splitting family contains an ω -splitting family of cardinality less than $\underline{b}$, then each α_2 -point is an α_1 -point.

Proof. Let x be an α_2 -point of a space Y and assume F_n is a sequence converging to x for each $n \in \omega$. Since we wish to show that x is an α_1 -point we may assume that the F_n 's are pairwise disjoint and that $\bigcup F_n = \omega$. Let $\mathfrak{X} = \mathfrak{X}(x, \langle F_n \rangle)$ be defined as above. By Lemma 2 and the hypothesis of this lemma, there is an ω -splitting family $\mathfrak{X}' \in [\mathfrak{X}]^{<\underline{b}}$. For each $A \in \mathfrak{X}'$, choose (by the definition of $\mathfrak{X}$) a sequence F_A converging to x and $f_A \in {}^\omega \omega$ so that $(A \cap F_n) - F_A \subset f_A(n)$ for each $n \in \omega$. Now choose $f \in {}^\omega \omega$ so that $f_A <^* f$ for each $A \in \mathfrak{X}'$ which we may do since $|\mathfrak{X}'| < \underline{b}$. We claim that $F = \bigcup F_n - f(n)$ converges to x , which would show that x is an α_1 -point. Indeed, assume F does not converge to x and choose $F' \in [F]^\omega$ such that x is not a limit point of F' . Since $\mathfrak{X}'$ is splitting, choose $A \in \mathfrak{X}'$ so that $A \cap F'$ is infinite. However this contradicts that F_A converges to x since $A \cap F' - F_A$ is finite.

4. *Remark.* We could replace the notion of ω -splitting in Lemma 3 by what one might call “ ω -hitting”.

ω -SPLITTING FAMILIES AND LAVER FORCING

5. **Theorem.** *In a model obtained by adding ω_2 -Laver reals to a model of CH, every ω -splitting family contains an ω -splitting family of cardinality ω_1 . Hence in this model each α_2 -space (w -space) is an α_1 -space (v -space).*

The theorem follows from the following four results. Proposition 6 is a collection of standard facts about Laver forcing, Lemma 7 is a standard reflection argument, Lemma 8 is a special case of a general preservation scheme proven in [S2] and Lemma 9 is new. Recall that $T \in \mathbb{L}$ (the Laver poset defined in [L]) if $T \subset {}^{<\omega}\omega$ has a root $t_0 = \text{root}(T)$ and for $t_0 \leq t \in T$ $\{n: t \hat{\ } n \in T\}$ is infinite. $\mathbb{L}$ is ordered by inclusion. For $\lambda \leq \omega_2$ let $\mathbb{P}_\lambda$ be the countable support λ -stage iteration of the forcing notion $\mathbb{L}$.

6. **Proposition.** [CH] $\mathbb{P}_{\omega_2}$ is an ω_2 -cc proper poset such that $1 \Vdash_{\mathbb{P}_\lambda} \underline{b} = \underline{c} = |\lambda \cdot \omega_1|^V$. Furthermore if $\mu < \lambda$ then $\mathbb{P}_\lambda$ is forcing isomorphic to $\mathbb{P}_\mu * \mathbb{P}_\lambda$.

For proofs of the various assertions in Proposition 6 we refer the reader to [L] and [S1].

7. **Lemma.** [CH] Let $\{X_\alpha: \alpha < \omega_2\}$ be $\mathbb{P}_{\omega_2}$ -names such that $1 \Vdash \{X_\alpha: \alpha < \omega_2\} \subset [\omega]^\omega$ is ω -splitting, then there is a $\lambda < \omega_2$ such that $1 \Vdash_{\mathbb{P}_\lambda} \{X_\alpha: \alpha < \lambda\}$ is ω -splitting.

Notation. If p is a member of a poset P and M is a set, “ $p \Vdash_P X$ is M -splitting” will abbreviate $p \Vdash |A \cap X| = |A - X|$ for each P -name $A \in M$ such that $p \Vdash A \in [\omega]^\omega$.

We shall say that a poset P is ω -splitting if the following are satisfied: whenever $P \in M$, where M is a countable elementary submodel of $H(\theta)$ for any sufficiently large θ , $p \in M \cap P$ and X is M -splitting then there is some $q < p$ which is (M, P) -generic and such that $q \Vdash X$ is M -splitting.

Note that an iteration of finitely many ω -splitting posets is again ω -splitting; hence proper.

8. **Lemma.** If $P_\delta = \langle (P_\alpha, Q_\alpha): \alpha < \delta \rangle \in M$ is a countable support iteration of ω -splitting (hence proper) posets then P_δ is also ω -splitting.

9. **Lemma.** Let M be a countable elementary submodel of $H(\omega_3)$ and let $T \in \mathbb{L} \cap M$, then if X is M -splitting there is an $(M, \mathbb{L})$ -generic $T' < T$ such that $T' \Vdash X$ is M -splitting. Therefore $\mathbb{L}$ is ω -splitting.

It may be worthwhile to record the following corollary to the above results.

10. **Corollary.** If a family is ω -splitting then it will still be ω -splitting after forcing with the countable support iteration of Laver forcing. Furthermore in any

model obtained by adding iteratively ω_2 -Laver reals the splitting number, $\underline{s}$, will be ω_1 .

Before proving Lemmas 7–9 let us indicate how Theorem 5 now follows. Let G be $\mathbb{P}_{\omega_2}$ -generic over V (a model of CH). Let $\mathfrak{X} = \{X_\alpha : \alpha < \omega_2 = \underline{c}\}$ be an ω -splitting family. Choose $\mathbb{P}_{\omega_2}$ -names $\{X_\alpha : \alpha < \omega_2\} \in V$ so that $1 \Vdash \mathfrak{X} = \{X_\alpha : \alpha < \omega_2\}$. By Lemma 7, there is a $\lambda < \omega_2$ so that $1 \Vdash_{P_\lambda} \{X_\alpha : \alpha < \lambda\}$ is ω -splitting. Let $G_\lambda = G \cap P_\lambda$; hence $V[G_\lambda] \models \{X_\alpha : \alpha < \lambda\}$ is ω -splitting. Since $\mathbb{P}_{\omega_2} \cong \mathbb{P}_\lambda * \mathbb{P}_{\omega_2}$, $V[G] \models \{X_\alpha : \alpha < \lambda\}$ is ω -splitting by Lemmas 8 and 9.

It remains to prove the Lemmas.

Proof of Lemma 7. By Proposition 6, we may assume that for each $\alpha < \omega_2$, there is an $f(\alpha) < \omega_2$ such that X_α is a $\mathbb{P}_{f(\alpha)}$ -name. Also if G is $\mathbb{P}_{\omega_2}$ -generic, $V[G] \models$ for each $\alpha < \omega_2$, there is a $g(\alpha) < \omega_2$ such that for each set $M \in [[\omega]^\omega]^\omega \cap V[G_\alpha]$ there is an $X \in \{\text{val}(X_\beta, G) : \beta < g(\alpha)\}$ which is M -splitting (since $V[G_\alpha] \models \underline{c} = \omega_1$). Since $\mathbb{P}_{\omega_2}$ is ω_2 -cc we may assume that $g \in V$. Now let h be a continuous strictly increasing function from ω_2 into ω_2 such that $f(\alpha) + g(\alpha) < h(\alpha + 1)$ for all $\alpha < \omega_2$. Choose $\lambda < \omega_2$ such that $h(\lambda) = \lambda$; it follows that $1 \Vdash_{P_\lambda} \{X_\alpha : \alpha < \lambda\}$ is ω -splitting.

Proof of Lemma 8. Technically we make the inductive assumption that for each $\beta < \alpha < \delta$ and each $p \in P_\beta$ we have that $p \Vdash P_\alpha/P_\beta$ is ω -splitting; where, as usual, P_α/P_β denotes the P_β -name satisfying the equation $P_\alpha = P_\beta * (P_\alpha/P_\beta)$. However in proving the inductive step we can just force with P_β and work in the extension. Therefore we may make the above inductive assumption and complete the proof by showing that P_δ is ω -splitting. As remarked when we defined the notion of an ω -splitting poset the proof is trivial in case δ is a successor ordinal. Now let θ be a large enough cardinal (i.e. $|\mathcal{P}(P_\delta)| < \theta$) and let $P_\delta \in M$ where M is a countable elementary submodel of $H(\theta)$. Suppose further that $p \in M \cap P_\delta$ and that X is M -splitting. Let $\{A_n : n \in \omega\}$ index the set of P_δ -names of subsets of ω which are in M . We may as well assume that $\text{cf}(\delta) = \omega$ since we will be choosing q to be M -generic and this q will force that the A_n 's are essentially $P_{M \cap \delta}$ -names. We may therefore choose a countable increasing sequence of ordinals cofinal in δ and by our inductive assumption we may as well assume that $\delta = \omega$.

As in [S2], we shall choose sequences p_n , q_n , k_n and m_n , by induction on n , so that:

- (1) $p_n \in P_\omega$, $q_n \in P_n$ and k_n , m_n are P_ω -names of integers,
- (2) $p_{n+1}|n = p_n|n$, $p_{n+1} < p_n$, $q_{n+1}|n = q_n$ and $q_n < p_n|n$,
- (3) for $n > 0$, $q_n \wedge p_n \Vdash A_n$ is finite or $m_n \in X \cap A_n$ and $k_n \in A_n - X$,
($q_n \wedge p_n$ is meant to denote the element $(q_n \wedge p_n|n)^\wedge p|n, \omega \in P_\omega$)
- (4) q_n is (M, P_n) -generic and $q_n \Vdash_{P_n} X$ is M -splitting, and
- (5) $q_n \Vdash_{P_n} \in M$.

Before we begin the induction, let us comment on what (5) means. It is not the case that p_n will be in M but $q_n \Vdash (\exists p' \in M)$ such that $p'/P_n = p_n/P_n$;

where p/P_n denotes the P_ω/P_n -name corresponding to p in the extension by P_n . It is not even the case that we can bring the “ $\exists$ ” sign outside the forcing statement. However condition 5 is essential in order for the induction to continue. Let $p_1 = p$ where $p \in P_\omega \cap M$ is as chosen above. Since P_1 is proper (and we have chosen θ large enough) we may choose $q_1 < p_1 \upharpoonright 1$ to be (M, P_1) -generic such that $q_1 \Vdash X$ is M -splitting. Suppose that p_n and q_n have been chosen. Let G_n be P_n -generic such that q_n and $p_n \upharpoonright n$ are in G_n . In $V[G_n]$, let p'_n be the element of M so that $p'_n/P_n = p_n/P_n$, and let $B_{n+1} = \{m : (\exists p \in P_\omega/P_n) p < p'_n/P_n \text{ and } p \Vdash m \in A_{n+1}\}$. Here we are assuming that A_n/P_n is the P_ω/P_n -name which results from evaluating the P_ω -name A_n and similarly for p'_n . Now $M[G_n]$ is an elementary submodel of $H(\theta)^{V[G_n]}$ which is a model of ZF-P (see [S1]) and $p'_n \in M$, hence $B_{n+1} \in M[G_n]$. Since X is $M[G_n]$ -splitting we may choose $m_{n+1} \in X \cap B_{n+1}$ and $p'/P_n < p'_n/P_n$, $p' \in M$ (by elementarity) such that $p' \Vdash m_{n+1} \in A_{n+1}$. Similarly we may choose $p'_{n+1}/P_n < p'/P_n$ and $k_{n+1} \notin X$ such that $p'_{n+1} \in M$, $p'_{n+1} \upharpoonright n \in G_n$ and $p'_{n+1} \Vdash k_{n+1} \in A_{n+1}$. By assumption Q_n is ω -splitting (in $V[G_n]$) hence we may choose $q'_{n+1} < p'_{n+1}(n)$, $q'_{n+1} \in Q_n$ so that q'_{n+1} is $(M[G_n], Q_n)$ -generic and so that $q'_{n+1} \Vdash X$ is $M[G_n]$ -splitting. Now we use the maximality principle to choose q_{n+1} , p_{n+1} and the names m_{n+1} and k_{n+1} so as to satisfy (1)–(5). That is, $p_n \upharpoonright n \Vdash$ if X is $M[G_n]$ -splitting and $M[G_n]$ is an elementary submodel of $H(\theta)^{V[G_n]}$ then there are $p'_{n+1}/P_n \in M[G_n]$, k_{n+1} and m_{n+1} as above. So we may choose a P_n -name, say p''_{n+1} , of an element of P_ω/P_n and P_ω -names k_{n+1} and m_{n+1} so that $p \upharpoonright n \Vdash$ if X is $M[G_n]$ -splitting and $M[G_n]$ is an elementary submodel of $H(\theta)^{V[G_n]}$ then $p''_{n+1} \in M[G_n]$ and $p''_{n+1} \Vdash k_{n+1}$ and m_{n+1} are as above. Next we choose a P_n -name q_{n+1} for q'_{n+1} . We let $p_{n+1} = p_n \wedge p''_{n+1}$ and $q_{n+1} = q_n \wedge q_{n+1}$. It is clear that (1), (2) are satisfied. By [S1], q_{n+1} is (M, P_{n+1}) -generic and clearly $q_{n+1} \Vdash X$ is M -splitting; hence (4) holds. The reason that (3), (5) are satisfied is that $q_{n+1} \Vdash M[G_{n+1}]$ is an elementary submodel of $H(\theta)^{V[G_{n+1}]}$, hence p''_{n+1} has the desired properties.

Now if $q \in P_\omega$ is such that $q \upharpoonright n = q_n$ for each $n \in \omega$ then q is (M, P_ω) -generic (see [S1]) and $q < p_n$ for each $n \in \omega$. It follows that $q \Vdash X$ is M -splitting, since for each $i < j < \omega$ there is an $n < \omega$ such that $1 \Vdash A_n = A_i - j$ and so $q < q \upharpoonright n \wedge p_n \Vdash m_n \in X \cap A_i - j$ and $k_n \in A_i - (X \cup j)$.

Proof of Lemma 9. Let $T \in L \cap M$ where M is a countable elementary submodel of $H(\omega_s)$ and assume X is M -splitting. Fix indexings $\{A_n \mid n \in \omega\}$ of $M \cap \{A \mid A \text{ is an } L\text{-name and } 1 \models A \in [\omega]^\omega\}$, and $\{D_n \mid n \in \omega\}$ of $M \cap \{D \subseteq L \mid D \text{ is dense open}\}$. We shall inductively define a descending sequence $\{T_n \mid n \in \omega\} \subset L$ (with $T_0 = T$) so that, for each $n \in \omega$:

1. $T_n \cap^n \omega = T_{n+1} \cap^n \omega$, and
2. If $T' < T_n$ then there is a $t \in T'$ such that
 - (a) $(T_n)_t \in M \cap D_n$ and

(b) $(T_n)_t \models X \cap A_n \neq \emptyset$ and $A_n - X \neq \emptyset$.

If we accomplish this, then condition 1 guarantees that $T' = \cap_n T_n \in L$. It is easy to see that condition 2(b) guarantees that $T' \models X$ is M -splitting. Condition 2(a) actually has the double role of ensuring that T' is M -generic and allowing the induction to continue.

Following [L], if $S', S \in L$ then we use $S' <^0 S$ to denote the situation where $S' < S$ and they have the same root. One of the key facts about Laver forcing from [L] is that if $S \in L$ and φ is any sentence of the forcing language then there is an $S' <^0 X$ such that either $S' \models \varphi$ or $S' \models \neg\varphi$. Therefore

$$(*) = \begin{cases} \text{if } A \text{ is an } L\text{-name and } F \text{ is a finite set such that} \\ S \models F \cap A \neq \emptyset, \text{ then there is an } S' <^0 S \text{ and an } x \in F \\ \text{such that } S \models x \in A. \end{cases}$$

Let us assume that $0 < n \in \omega$ and that T_{n-1} has been chosen as above. Let I be the set of members of $T_{n-1} - \leq^n \omega$ which are minimal with respect to the property that $(T_{n-1})_t \in M$. Note that the minimality of the members of I and condition 2(a) guarantee that the collection $\{(T_{n-1})_t | t \in I\}$ is an antichain in L which is maximal below T_{n-1} . Now if we find, for each $t \in I$, a condition $T'_t <^0 (T_{n-1})_t$ satisfying condition 2, then we can define T_n to be $\bigcup \{T'_t | t \in I\}$. This works since $\{T'_t | t \in I\}$ is a maximal-below- T_n antichain. For the same reason, repeated uses of Facts 1 to 3 finish the proof.

Fact 1. If $S \in L \cap M$ and $n \in \omega$, then there is an $S' <^0 X$ such that the collection $(S')_t | (S')_t \in D_n \cap M$ is predense below S' .

Fact 2. If $S \in L \cap M$ and $n \in \omega$ there is an $S' <^0 X$ such that the collection $\{(S')_t : (S')_t \in M \text{ and } (S')_t \models X \cap A_n \neq \emptyset\}$ is predense below S' .

Fact 3. If $S \in L \cap M$ and $n \in \omega$ there is an $S' <^0 S$ such that the collection $\{(S')_t | (S')_t \text{ and } (S')_t \models A_n - X \neq \emptyset\}$ is predense below S' .

Fact 1 is, of course, a well-known property of L and its proof is similar to the proof of Fact 2. In the proof of Fact 2 we are just using that X is M -splitting. Since $\omega - X$ is also M -splitting, Fact 3 follows from Fact 2. Now let us prove Fact 2.

Let $B_n = \{k \in \omega | (\exists S' <^0 S) S' \models k \in A_n\}$. Let us first suppose that B_n is infinite. In this case, $B_n \in [\omega]^\omega \cap M$, hence we have that $X \cap B_n \neq \emptyset$. It follows that we may choose $S' <^0 S$, $S' \in M$ and $k \in X \cap B_n$ so that $S' \models k \in A_n$ —which certainly suffices.

Now let us suppose that $\max(B_n) < m$. Let I be the set of minimal elements of $\{t \in S | (\exists m_t \geq m)(\exists S'_t <^0 X_t) S'_t \models m_t \in A_n\}$. Since $1 \models (\exists k \geq m) k \in A_n$ and the members of I are minimal, $\{S'_t | t \in I\}$ is a maximal-below- S antichain of L . For each $t \in I$, fix a minimal m_t and an $S'_t \in M$ as in the description of I ; hence $\{m_t | t \in I\} \in M$. We shall show that $S' = \bigcup \{S'_t | t \in I \text{ and } m_t \in X\}$ works. That is, we prove that $S' \in L$, $S' <^0 S$ and simply note that $\{S'_t | t \in I \text{ and } m_t \in X\}$ is a maximal-below- S' antichain. Therefore it suffices to show

that if $s \in S'$ is such that $\text{root}(S) \geq s$, then s has infinitely many immediate successors in S' . First suppose that there is a $t \in I$ with $t \leq s$. Since members of I are minimal, it follows that $(S')_s = S' \cap S_s$ —hence $t \in S'$ and $m_t \in X$. Therefore $S'_t \subseteq S'$ and s has infinitely many immediate successors in S' . If there is no such $t \in I$, then S_s has no $<^0$ -extension which decides a value of A_n above m . Suppose now that $k \in \omega$ is such that $\max(\{i | s^{\wedge} i \in S'\}) < k$. Let S'' be the $<^0$ -extension of S_s obtained by removing $\{t | (\exists i < k) s^{\wedge} i \leq t\}$. We claim that $\{m_t | t \in I \cap S''\}$ is infinite. Indeed, since $\{S_t | t \in I \cap S''\}$ is predense below S'' , $S'' \models A_n \cap \{m_t | t \in I \cap S''\} \neq \emptyset$. If the set was finite then, by (*), S'' (hence S_s) would have $<^0$ -extension picking one of the values. But now $X \cap \{m_t | t \in I \cap S''\}$ is nonempty, hence any $t \in I \cap S''$ with $m_t \in X$ is an extension of s in S' —a contradiction to the choice of k .

REFERENCES

- [A1] A. V. Arhangel'skii, *The frequency spectrum of a topological space and the classification of spaces*, Dokl. Akad. Nauk. SSSR **206** (1972), 265–268 = Sov. Math. Dokl. **13** (1972), 265–268.
- [A2] —, *The frequency spectrum of a topological space and the product operation*, Trudy Mosk. Mat. Obs. **40** (1979) = Transl. Moscow Math. Soc. (1981), Issue 2, 163–200.
- [DS] A. Dow and J. Steprāns.
- [G] G. Gruenhage, *Infinite games and generalizations of first countable spaces*, Gen. Top. Appl. **6** (1976), 339–352.
- [L] R. Laver, *On the consistency of Borel's conjecture*, Acta Math. **137** (1976), 151–169.
- [LNy] R. Levy and P. Nyikos, *Families in $\beta\omega$ whose union is regular open*, preprint.
- [N] T. Nogura, *A counterexample for a problem of Arhangel'skii concerning the product of Fréchet spaces*, Top. Appl. **25** (1987), 75–80.
- [Ny1] P. Nyikos, *$^{\omega}\omega$ and the Fréchet-Urysohn property*, in preparation.
- [Ny2] —, *The Cantor tree and the Fréchet-Urysohn property*, preprint, 1987.
- [O] R. C. Olson, *Bi-quotient maps and countably bisequential spaces*, Gen. Top. Appl. **4** (1974), 1–28.
- [S1] S. Shelah, *Proper forcing*, Springer Lecture Notes, 1982.
- [S2] —, *Cardinal invariants of the continuum*, Axiomatic Set Theory, Eds. J. Baumgartner, D. A. Martin, S. Shelah, Contemp. Math. Amer. Math. Soc., Providence, R.I., 1986.
- [Sh] Sharma, *Some characterizations of W -spaces and w -spaces*, Gen. Top. Appl. **9** (1978) 289–293.

DEPARTMENT OF MATHEMATICS, YORK UNIVERSITY, NORTH YORK, ONTARIO, M3J 1P3, CANADA