SPLITTING THEOREM FOR HOMOLOGY OF $GL(R)$

STANISLAW BETLEY

(Communicated by Frederick R. Cohen)

ABSTRACT. It is proved that if $\{M_n\}$ is a stable system of coefficients for $GL_n(R)$, and $H_0(GL(R), \lim(M_n))$ contains $\mathbb{Z}$, then for any j , the group $H_j(GL(R), \lim(M_n))$ contains $H_j(GL(R), \mathbb{Z})$ as a direct summand. Now let $GL(\mathbb{Z})$ act on $M(\mathbb{Z})$ (matrices over $\mathbb{Z}$) by conjugation. Then our theorem implies that the trace map $\text{tr}: M(\mathbb{Z}) \rightarrow \mathbb{Z}$ is a split epimorphism on homology.

0. INTRODUCTION

In the late seventies the development of algebraic K -theory forced mathematicians to study the following question: does the inclusion of $GL_n(R)$ into $GL_{n+1}(R)$ induce the isomorphism on homology with trivial or twisted coefficient systems? (R is any ring). The most general answer to this question was given by W. van der Kallen in [6]. He solved that problem in affirmative for rings which satisfy Bass's stable range condition (see [1] for the definition). In [2] we used the homological stability theorems to get information about the homology of $GL_n(R)$. More precisely, we gave there a description of coefficient systems for $GL_n(R)$ which give stably trivial homology. In the following note we study the opposite situation and describe systems of coefficients which give nontrivial homology of $GL(R) = \varinjlim_n GL_n(R)$.

§1

1.1. Definition. Let R be any ring. Let M_n be a $GL_n(R)$ -module for any natural n . Assume that we have homomorphisms $F_n: M_n \rightarrow M_{n+1}$ which are equivariant under the inclusion of $GL_n(R)$ into $GL_{n+1}(R)$ given by $B \rightarrow \begin{vmatrix} B & 0 \\ 0 & 1 \end{vmatrix}$. Then we say that the set $\{M_n\}$ forms a system of coefficients for $GL_n(R)$.

Received by the editors October 10, 1988, and in revised form, February 20, 1989.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 20G10; Secondary 18G99, 55N99.

Key words and phrases. Homology of a group, stable system of coefficients.

©1990 American Mathematical Society
0002-9939/90 \$1.00 + \$.25 per page

In this section we will study only systems $\{M_n\}$ of coefficients for $\mathrm{Gl}_n(R)$ which satisfy the following conditions:

- (a) $\{M_n\}$ is a stable system of coefficients for $\mathrm{Gl}_n(R)$ (i.e. for any i and sufficiently large n the map $H_i(\mathrm{Gl}_n(R); M_n) \rightarrow H_i(\mathrm{Gl}_{n+1}(R); M_{n+1})$ is an isomorphism);
- (b) for any n and k , the image of M_n in M_{n+k} is fixed pointwise by the subgroup $\begin{vmatrix} I_n & 0 \\ 0 & \mathrm{Gl}_k \end{vmatrix}$ of $\mathrm{Gl}_{n+k}(R)$.

For convenience we will assume that for any n , the module M_n is finitely generated as an abelian group, and R always denotes a ring with unit which satisfies Bass's stable range condition. We will denote $\varinjlim_n \mathrm{Gl}_n(R)$ as $\mathrm{Gl} R$ and $\varinjlim_n M_n$ as M .

1.2. Theorem. Assume that $\varinjlim H_0(\mathrm{Gl}_n(R); M_n) = A \neq 0$. Let $f: \mathbf{Z} \rightarrow A$ be a homomorphism and $f_*: H_j(\mathrm{Gl} R; \mathbf{Z}) \rightarrow H_j(\mathrm{Gl} R; A)$ the induced homomorphism. Then, for any $j \geq 0$, f_* can be factored through $H_j(\mathrm{Gl} R; M)$. In particular, if f_* is nontrivial, then $H_j(\mathrm{Gl} R; M) \neq 0$.

Proof. Let n be such a large natural number that $H_j(\mathrm{Gl}_n(R); M_n) = H_j(\mathrm{Gl} R; M)$ and $H_j(\mathrm{Gl}_n(R); \mathbf{Z}) = H_j(\mathrm{Gl} R; \mathbf{Z})$. Then by [5] $H_0(\mathrm{Gl}_n(R); M_n) = A$ and the quotient map $M_n \xrightarrow{j_n} A$ is a $\mathrm{Gl}_n(R)$ -homomorphism where $\mathrm{Gl}_n(R)$ acts trivially on A . We have the following commutative diagram of $\mathrm{Gl}_n(R)$ -modules:

$$\begin{array}{ccccc} M_n & \xrightarrow{F_n} & \dots & \xrightarrow{F_{2n-1}} & M_{2n} \\ \downarrow j_n & & & & \downarrow j_{2n} \\ A & = & \dots & = & A \end{array}$$

We have the following commutative diagram of groups and modules:

$$(*) \quad \begin{array}{ccc} & & (\mathrm{Gl}_n(R), \mathbf{Z}) \\ & \searrow G & \downarrow F \\ & & (\mathrm{Gl}_{2n}(R), M_{2n}) \\ & & \downarrow (\mathrm{id}, j_{2n}) \\ (\mathrm{Gl}_{2n}(R), \mathbf{Z}) & \xrightarrow{(\mathrm{id}, f)} & (\mathrm{Gl}_{2n}(R), A) \end{array}$$

where G is defined as the map which takes a matrix $B \in \mathrm{Gl}_n(R)$ to $\begin{vmatrix} I_n & 0 \\ 0 & B \end{vmatrix}$ and is equal to identity on coefficients. To define F we use condition (b) about

$\{M_n\}$. Let h be a map in the category of abelian groups given by the following commutative diagram:

$$\begin{array}{ccc} & & Z \\ & \nearrow h & \downarrow f \\ M_n & \xrightarrow{j_n} & A \end{array}$$

We can then define F as the same map as G on the groups and as $F_{2n-1} \circ \dots \circ F_n \circ h$ on coefficients. By condition (b), F is well defined.

Looking now at diagram (*) after taking the j th homology group, we see that the map $((id, f) \circ G)_*: H_j(Gl_n(R); \mathbb{Z}) \rightarrow H_j(Gl_{2n}(R); A)$ can be factored through the group $H_j(Gl_{2n}(R); M_{2n})$. On the other hand, it is obvious that $((id, f) \circ G)_*$ agrees with $f_*: H_j(Gl R; \mathbb{Z}) \rightarrow H_j(Gl R; A)$ — we have only to use the fact that the lower and upper inclusions of $Gl_n(R)$ into $Gl_{2n}(R)$ give the same map on homology with trivial coefficients. Thus f_* factors through $H_j(Gl_{2n}(R); M_{2n})$, and by the assumption on n this last group is isomorphic to $H_j(Gl R; M)$.

1.3. *Remark.* The following two conditions are equivalent for any group G and any G -module M which is finitely generated as an abelian group:

- (1) $H_0(G; M) \neq 0$;
- (2) there exists a G -epimorphism f , from M to the trivial G -module $\mathbb{Z}$ or $\mathbb{Z}_p$.

Proof. If $H_0(G; M) \neq 0$, take as f the natural projection $M \rightarrow H_0(G; M)$ composed with a projection from $H_0(G; M)$ onto any of its factors.

If there exists f as in condition (2), then for any $g \in G$ and any $m \in M$, $f(gm) = gf(m)$; so $f((g-1)m) = 0$, and hence f induces an epimorphism $H_0(G; M) \rightarrow \mathbb{Z}$ or $\mathbb{Z}_p$.

1.4. **Corollary.** Let R be any commutative ring with unit satisfying the hypothesis of Theorem 1.2. Let M_n denotes the $n \times n$ matrices with entries in R on which the group $Gl_n(R)$ acts by conjugation. Then the trace map $tr: M \rightarrow R$ induces a nontrivial map on homology provided that $H_*(Gl R; \mathbb{Z}) \rightarrow H_*(Gl R; R)$ is nontrivial (this map is induced by the obvious homomorphism $\mathbb{Z} \rightarrow R$).

1.5. *Remark.* Now let R denote any ring with unit satisfying Bass's stable range conditions and let P denote the R -bimodule. Let $H_*^{\text{Hoch}}(R, P)$ denote the Hochschild homology of R with coefficients in P . Let $M(P)$ denote the full group of matrices over P on which $Gl R$ acts by conjugation. It was shown by Kassel in [7] that

$$H_0(Gl R, M(P)) = H_0^{\text{Hoch}}(R, P)$$

and

$$H_1(Gl R, M(P)) = H_1(Gl R, \mathbb{Z}) \otimes H_0^{\text{Hoch}}(R, P) \oplus H_1^{\text{Hoch}}(R, P).$$

Our Theorem 1.2 implies that $H_j(\mathrm{Gl} R, \mathbf{Z}) \otimes H_0^{\mathrm{Hoch}}(R, P)$ is a direct summand of $H_j(\mathrm{Gl} R, M(P))$ provided that $H_0^{\mathrm{Hoch}}(R, P)$ is $\mathbf{Z}$ -free (see also [5]).

1.6. Corollary. *Now let $R = \mathbf{Z}$ and M be as in 1.4. Then $tr: M \rightarrow \mathbf{Z}$ gives a split epimorphism on homology.*

It is known that in 1.6, tr_* is an isomorphism when we tensor homology groups with Q ([4]); but such splitting as in 1.6 is obvious. Using Kassel's results from [7], it is easy to calculate that tr_* from 1.6 is an isomorphism on H_0 and H_1 .

1.7. Remark. Condition (b) is not very restrictive. The main examples of stable systems of coefficients for $\mathrm{Gl}_n(R)$ are obtained by applying any functor of finite degree to R^n considered in an obvious way as a $\mathrm{Gl}_n(R)$ module (see [3] or [6]). But it is easy to see that in such a way we can obtain only systems which satisfy (b).

1.8. Remark. Theorem 1.2 remains true when we replace the groups $\mathrm{Gl}_n R$ by any family of subgroups $G_n \subseteq \mathrm{Gl}_n R$ such that

(1) the following diagram is commutative:

$$\begin{array}{ccc} \mathrm{Gl}_n R & \longrightarrow & \mathrm{Gl}_{n+1} R \text{ (the standard "upper" inclusion)} \\ \cup & & \cup \\ G_n & \longrightarrow & G_{n+1} \end{array}$$

(2) the "lower" and "upper" inclusions $G_n \rightarrow G_{2n}$ are defined and conjugated, and they agree with the "lower" and "upper" inclusions of $\mathrm{Gl}_n R$ into $\mathrm{Gl}_{2n} R$ for some sufficiently large n (depending on j).

In particular, it is true for $\mathrm{Sl}_n R$ in a commutative case and more generally for $E_n(R)$.

§2

In this section we drop the assumption that our system of coefficients is stable. We call $\varinjlim_n H_0(\mathrm{Gl}_n(R); M_n)$ as A .

2.1. Theorem. *Assume that there is a $\mathbf{Z}$ -homomorphism $f: \mathbf{Z} \rightarrow A$ which gives us a nontrivial map*

$$f_*: \varinjlim_n H_j(\mathrm{Gl}_n(R); \mathbf{Z}) \rightarrow \varinjlim_n H_j(\mathrm{Gl}_n(R); A).$$

Then $\varinjlim_n H_j(\mathrm{Gl}_n(R); M_n)$ is nontrivial.

Proof. Let $M = \varinjlim_n M_n$. There is a natural epimorphism $g: M \rightarrow A$. Let f' be a map obtained from the following diagram

$$\begin{array}{ccc} & \mathbf{Z} & \\ f' \swarrow & & \downarrow f \\ M & \xrightarrow{g} & A \end{array}$$

For sufficiently large n we can easily find a map H which fits into the following commutative diagram:

$$\begin{array}{ccccc} & & \mathbf{Z} & & \\ & & \downarrow f & & \\ & H \swarrow & & & \\ & M_n & \xrightarrow{\quad} & M & \\ & \downarrow j_n & & \downarrow g & \\ H_0(Gl_n(R); M_n) & \xrightarrow{\quad} & A & & \end{array}$$

We can now apply precisely the same method as in the proof of 1.2, using our new map H instead of h , replacing j_{2n} by j_{2n} composed with the standard homomorphism $H_0(Gl_{2n} R; M_{2n}) \rightarrow A$, and taking sufficiently large n . We then obtain the following commutative diagram:

$$\begin{array}{ccc} \varinjlim_n H_j(Gl_n(R); \mathbf{Z}) & \xrightarrow{f_*} & \varinjlim_n H_j(Gl_n(R); A) \\ & \searrow & \nearrow \\ & \varinjlim_n H_j(Gl_n(R); M_n) & \end{array}$$

The existence of such a diagram immediately implies our Theorem 2.1.

REFERENCES

1. H. Bass, *Algebraic K-theory*, Benjamin, New York, 1968.
2. S. Betley, *Vanishing theorems for homology of $GL R$* , J. Pure and Appl. Algebra, 58 (1989) 213–226.
3. W. G. Dwyer, *Twisted homological stability for general linear groups*, Ann. of Math. (2) **111** (1980), 239–251.
4. F. T. Farrell and W. C. Hsiang, *On the rational homotopy groups of the diffeomorphism groups of discs, spheres and aspherical manifolds*, Proc. Sympos. Pure Math. **32** (1978), 325–337.
5. T. G. Goodwillie, *On the general linear group and Hochschild homology*, Ann. of Math. (2) **121** (1985), 383–407.

6. W. van der Kallen, *Homology stability for linear group*, Invent. Math. **60** (1980), 269–295.
7. C. Kassel, *Calcul algébrique de l'homologie de certains groupes de matrices*, J. Algebra **80** (1983), 235–260.

WYDZIAŁ MATEMATYKI UW, INSTYTUT MATEMATYKI, PKiN IXp., 00-901 WARSAW, POLAND