$\mathcal{E}_1$ IS UNIFORMLY KADEC-KLEE

CHRIS LENNARD

(Communicated by William J. Davis)

ABSTRACT. A dual Banach space X is Kadec-Klee in the weak* topology if weak* and norm convergence of sequences coincide in the unit sphere of X . We shall consider a stronger, uniform version of this property. A dual Banach space X is uniformly Kadec-Klee in the weak* topology (UKK^*) if for each $\varepsilon > 0$ we can find a δ in $(0, 1)$ such that every weak*-compact, convex subset C of the unit ball of X whose measure of norm compactness exceeds ε must meet the $(1 - \delta)$ -ball of X . We show in this paper that $C_1(\mathcal{H})$, the space of trace class operators on an arbitrary infinite-dimensional Hilbert space $\mathcal{H}$ is UKK^* . Consequently $C_1(\mathcal{H})$ has weak*-normal structure. This answers affirmatively a question of A. T. Lau and P. F. Mah. From this it follows that $C_1(\mathcal{H})$ has the weak*-fixed point property.

0. INTRODUCTION

We establish in this paper that the trace class $(\mathcal{E}_1, \|\cdot\|_1)$ has the weak* uniform Kadec-Klee property (UKK^*). This strengthens a result of Arazy [1] and of Simon [11] who show that $(\mathcal{E}_1, \|\cdot\|_1)$ has the weak* Kadec-Klee property (KK^*). They in fact show more—a unitary matrix space (minimal symmetrically normed ideal) $\mathcal{E}_E$ is KK^* if the symmetric sequence space E corresponding to $\mathcal{E}_E$ under the Calkin correspondence is KK^* . The analogous question for the UKK^* property remains open.

The fact that $(\mathcal{E}_1, \|\cdot\|_1)$ is UKK^* has a number of interesting consequences $(\mathcal{E}_1, \|\cdot\|_1)$ must have weak* normal structure which implies the weak* fixed point property, by van Dulst and Sims [2, Theorem 3]. This answers affirmatively a question of Lau and Mah [8], who show that $(\mathcal{E}_1, \|\cdot\|_1)$ has quasi-weak* normal structure (which is weaker than weak* normal structure).

We note that a Banach space $(X, \|\cdot\|)$ having ε - UKK^* for some $\varepsilon \in (0, 1)$ is sufficient to give us that X has weak* normal structure. That $(\mathcal{E}_1, \|\cdot\|_1)$ enjoys the stronger UKK^* property shows us, by van Dulst and Sims [2, Theorem 4], that the Chebychev centres of weak*-compact, convex sets in $\mathcal{E}_1$ are compact.

Received by the editors December 12, 1987.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 46B20.

The financial assistance of a Fellowship from Kent State University is gratefully acknowledged.

©1990 American Mathematical Society
 0002-9939/90 \$1.00 + \$.25 per page

For more information on the concepts discussed above, all of which can be related to fixed point properties for nonexpansive mappings, see van Dulst and Sims [2] or Sims [12, 13]. Also we remark that van Dulst and de Valk [3] have recently investigated Kadec-Klee properties in the sequence spaces l_M and h_M , where M is an Orlicz function.

I wish to thank Brailey Sims for suggesting this problem and for many stimulating conversations concerning it.

1. PRELIMINARIES

Recall that if $(X, \|\cdot\|)$ is a Banach space, a standard measure of compactness of a nonempty subset S of X is given by

$$\gamma(S) := \sup\{\text{sep}(\{x_n\}_{n \in N}) : \{x_n\}_{n \in N} \subseteq S\},$$

where $\text{sep}(\{x_n\}_{n \in N}) := \inf_{n \neq m} \|x_n - x_m\|$. We will refer to γ as “the measure of compactness”.

Define now, for a dual space X , and for $S \subseteq X$,

$$\beta(S) := \sup \left\{ \text{sep}(\{x_n\}_{n \in N}) : \{x_n\}_{n \in N} \subseteq S \right. \\ \left. \text{and weak}^* - \lim_{n \in N} x_n = x \text{ for some } x \in X \right\}.$$

Clearly $\beta(S) \leq \gamma(S)$. In some cases more can be said. The proof of the following lemma is straightforward and is therefore omitted.

1.1. Lemma. *Let X be a dual Banach space with a weak*-sequentially compact unit ball. Then $\beta(S) = \gamma(S)$ for every nonempty set $S \subseteq X$.*

A dual Banach space $(X, \|\cdot\|)$ is said to be Kadec-Klee in the weak*-topology (KK^*) if weak* and norm convergence of sequences coincide in the unit sphere of X . We remark that the analogous Kadec-Klee property for the weak topology of a general Banach space (denoted by KK) is often called the Radon-Riesz property or property H .

It is easily seen that a Banach space (dual Banach space) $(X, \|\cdot\|)$ is KK (KK^*) if whenever $x_n \rightarrow x$ as $n \rightarrow \infty$ in the weak (weak*) topology on X and $\lim_{n \in N} \|x_n\| = \|x\|$, it follows that $\lim_{n \in N} \|x_n - x\| = 0$.

We introduce below the properties ε - UKK^* and UKK^* in a manner due to Sims [13].

1.2. Definition. Suppose that X is a dual Banach space and $\varepsilon > 0$. X is ε -uniformly Kadec-Klee in the weak* topology (ε - UKK^*) if there exists $\delta \in (0, 1)$ such that whenever C is a weak* compact, convex subset of B_X with $\gamma(C) \geq \varepsilon$ it follows that $C \cap (1 - \delta)B_X \neq \emptyset$.

X is uniformly Kadec-Klee in the weak* topology (UKK^*) if it is ε - UKK^* for all $\varepsilon > 0$.

By substituting the weak topology on an arbitrary Banach space for the weak* topology on a dual Banach space in the above definitions we get the definitions

of the properties ε - UKK (also called $WUKK$) and UKK . UKK was introduced by Huff [5] and ε - UKK , ε - UKK^* and UKK^* are due to van Dulst and Sims [2].

It is not hard to see that $UKK \Rightarrow KK$ and, for dual spaces, $UKK^* \Rightarrow KK^*$. Moreover, both implications are strict. Indeed set $X = (l_2 \oplus l_3 \oplus \cdots \oplus l_n \oplus \cdots)_2$. Then, as noted by Huff [5], X is reflexive, X is KK but X is not UKK . Further, X does not have an equivalent norm for which $(X, \|\cdot\|)$ is UKK . In contrast to this, from Kadec [6] and Klee [7], we have that every separable Banach space (dual Banach space) admits an equivalent norm (dual norm) such that X becomes KK (KK^*). This result may also be found in Lindenstrauss and Tzafriri [9].

If a Banach space $(X, \|\cdot\|)$ is uniformly convex (which forces X to be reflexive) then X must be UKK^* . The converse fails. $(l_1, \|\cdot\|_1)$ and $(\mathcal{E}_1, \|\cdot\|_1)$ are both UKK^* but fail to be uniformly convex (or even uniformly convexifiable).

We present below a simple characterization of dual Banach spaces with ε - UKK^* (and consequently also those with UKK^*) due to Sims [13].

1.3. Proposition. *Let $\varepsilon > 0$. A dual Banach space $(X, \|\cdot\|)$ over $\mathbf{K}$ ($= \mathbf{R}$ or $\mathbf{C}$) is ε - UKK^* if and only if there exists $k \in (0, 1)$ such that for all weak* continuous linear functionals f with $\|f\| = 1$ we have $\gamma(S[f, k]) \leq \varepsilon$, where*

$$S[f, k] := \{x \in B_X : \operatorname{Re} f(x) \geq k\}.$$

Throughout this paper $(\mathcal{H}, (\cdot, \cdot))$ will denote an arbitrary infinite-dimensional real or complex Hilbert space. We will denote the field by $\mathbf{K}$ i.e. $\mathbf{K} = \mathbf{R}$ or $\mathbf{C}$. $(\mathcal{B}(\mathcal{H}), \|\cdot\|_\infty)$ is the Banach algebra of all bounded linear operators on $\mathcal{H}$, with the usual supremum norm. Any $B \in \mathcal{B}(\mathcal{H})$ has a polar decomposition $B = U[B]$, where $[B] := (B^*B)^{1/2}$ is the absolute value of B and U is that partial isometry with initial set the closure of the range of $[B]$, and final set the closure of the range of B .

$\mathcal{C}_\infty(\mathcal{H})$ is the ideal of compact operators in $\mathcal{B}(\mathcal{H})$. Any $C \in \mathcal{C}_\infty$ has a Schmidt decomposition

$$C = \sum_{j=1}^{\infty} s_j(C) \phi_j \otimes \overline{\psi_j},$$

where $s(C) := \{s_j(C)\}_j \subseteq c_0$ is the sequence of singular values of C , $\{\phi_j\}_{j \in \mathbf{N}}$ and $\{\psi_j\}_{j \in \mathbf{N}}$ are orthonormal sequences in $\mathcal{H}$ and $f \otimes g := (\cdot, g)f$ for all $f, g \in \mathcal{H}$.

$\mathcal{E}_1(\mathcal{H})$ is the ideal of all trace class operators in $\mathcal{B}(\mathcal{H})$ and $\|\cdot\|_1$ is given by

$$\|A\|_1 = \operatorname{tr}[A],$$

for all $A \in \mathcal{E}_1$. $(\mathcal{E}_1, \|\cdot\|_1)$ is isometrically isomorphic to the dual of the Banach space $(\mathcal{C}_\infty, \|\cdot\|_\infty)$, and the duality is given by the trace functional i.e. each $\varphi \in \mathcal{C}_\infty^*$ is of the form

$$\varphi : C \mapsto \operatorname{tr}(AC) : \mathcal{C}_\infty \rightarrow \mathbf{K}.$$

For more information on $\mathcal{B}(\mathcal{H})$, the trace class and trace duality see Schatten [10] or Gohberg and Krein [4].

$\mathcal{B}_{\text{s.a.}}(\mathcal{H})$ is the class of all selfadjoint operators in $\mathcal{B}(\mathcal{H})$, and it is partially ordered by the quadratic form ordering $\leq$ i.e. for $B, B' \in \mathcal{B}_{\text{s.a.}}$, $B \leq B'$ if for all $f \in \mathcal{H}$

$$(Bf, f) \leq (B'f, f).$$

For a family $\{B_\tau\}_\tau$ in $\mathcal{B}_{\text{s.a.}}$ and $B \in \mathcal{B}_{\text{s.a.}}$, we denote by $B_\tau \uparrow_\tau B$ ($B_\tau \downarrow_\tau B$) the fact that $\{B_\tau\}_\tau$ is upwards directed (downwards directed) to B .

2. $\mathcal{E}_1$ is UKK^*

We will need the following lemma. The proof, using the Schmidt decomposition mentioned above, is straightforward and is therefore omitted.

2.1. Lemma. *Let $\{Q_\tau\}_\tau$ be a family of orthogonal projections in $\mathcal{B}(\mathcal{H})$ with $Q_\tau \downarrow_\tau 0$ and $C \in \mathcal{E}_\infty$. Then*

$$\lim_\tau \|CQ_\tau\|_\infty = 0.$$

The proposition below, upon which our main result depends, is due to Arazy [1]. We include a proof for the sake of completeness.

2.2. Proposition. *Let $A \in \mathcal{E}_1$, let P be an orthogonal projection in $\mathcal{B}(\mathcal{H})$ and $Q = I - P$. Then*

$$(a) \quad \|A\|_1^2 \geq \|PAP\|_1^2 + \|PAQ\|_1^2 + \|QAP\|_1^2 + \|QAQ\|_1^2$$

and

$$(b) \quad \|PAQ\|_1 + \|QAP\|_1 + \|QAQ\|_1 \leq \sqrt{3}(\|A\|_1^2 - \|PAP\|_1^2)^{1/2}.$$

Proof. Clearly (b) follows from (a). To prove (a), we begin by letting $A = U[A]$ be the polar decomposition of A . Note that $[A^*] = U[A]U^*$.

We remark that

$$(1) \quad \text{if } 0 \leq X \in \mathcal{E}_1 \text{ and } 0 \leq Y \leq Z \in \mathcal{B}(\mathcal{H}), \text{ then } \text{tr}(XY) \leq \text{tr}(XZ).$$

Now consider PAQ . Let $PAQ = V[PAQ]$ be its polar decomposition. We have that

$$\begin{aligned} \|PAQ\|_1^2 &= (\text{tr}[PAQ])^2 = (\text{tr}(V^*PAQ))^2, \\ &= (\text{tr}(V^*PU[A]Q))^2 = (\text{tr}(S^*T))^2, \\ &\quad \text{where } S = [A]^{1/2}U^*PV \text{ and } T = [A]^{1/2}Q, \\ &\leq \text{tr}(S^*S) \text{tr}(T^*T), \text{ by the Cauchy-Schwarz inequality,} \\ &= \text{tr}(V^*PU[A]U^*PV) \text{tr}(Q[A]Q) \\ &= \text{tr}(P[A^*]PVV^*) \text{tr}(Q[A]Q) \\ &\leq \text{tr}(P[A^*]P) \text{tr}(Q[A]Q), \end{aligned}$$

by (1) above, as VV^* is an orthogonal projection and so $VV^* \leq I$. So we have

$$\|PAQ\|_1^2 \leq \text{tr}(P[A^*]P) \text{tr}(Q[A]Q),$$

and of course we also have three analogous upper estimates for each of $\|QAP\|_1^2$, $\|PAP\|_1^2$ and $\|QAQ\|_1^2$.

Consequently,

$$\begin{aligned} & \|PAP\|_1^2 + \|PAQ\|_1^2 + \|QAP\|_1^2 + \|QAQ\|_1^2 \\ & \leq (\text{tr}(P[A]P) + \text{tr}(Q[A]Q))(\text{tr}(P[A^*]P) + \text{tr}(Q[A^*]Q)) \\ & = (\text{tr}[A])(\text{tr}[A^*]) = \|A\|_1 \|A^*\|_1 \\ & = \|A\|_1^2. \end{aligned}$$

2.3. Lemma. $\mathcal{E}_1$ has a weak* sequentially compact unit ball.

Proof. Any sequence $\{A_n\}_{n \in \mathbb{N}}$ in $\mathcal{E}_1 = \mathcal{E}_1(\mathcal{H})$ lives essentially in $\mathcal{E}_1(\mathcal{H}_0)$, where $\mathcal{H}_0$ is the separable Hilbert space

$$\text{closure} \left(\text{span} \left(\bigcup_{n=1}^{\infty} (\text{range}(A_n) \cup \text{range}(A_n^*)) \right) \right);$$

while the unit ball of $\mathcal{E}_1(\mathcal{H}_0)$ is weak* sequentially compact since $\mathcal{E}_1(\mathcal{H}_0) = \mathcal{E}_{\infty}^*(\mathcal{H}_0)$ and $\mathcal{E}_{\infty}(\mathcal{H}_0)$ is separable. From this, using the trace functional, which (as noted above) gives the duality between $\mathcal{E}_1(\mathcal{H})$ and $\mathcal{E}_{\infty}(\mathcal{H})$, the result follows. $\square$

We now present the main result. The argument is similar to that of Brailey Sims in [13] for the case of $(I_1, \|\cdot\|_1)$. We use the characterization of UKK^* that comes from Proposition 1.3.

2.4. Theorem. $(\mathcal{E}_1, \|\cdot\|_1)$ is UKK^* .

Proof. Fix $k \in (0, 1)$ and then fix $C \in \mathcal{E}_{\infty}$ with $\|C\|_{\infty} = 1$. Choose a family $\{P_{\tau}\}_{\tau}$ of finite-dimensional orthogonal projections with $P_{\tau} \uparrow I$. Let $Q_{\tau} = I - P_{\tau}$ for each τ and note that $Q_{\tau} \downarrow 0$.

Consider $E \in S[C, k]$ and define for each τ ,

$$\alpha_{\tau}(E) = \|P_{\tau}EQ_{\tau}\|_1 + \|Q_{\tau}EP_{\tau}\|_1 + \|Q_{\tau}EQ_{\tau}\|_1.$$

By Proposition 2.2, for each τ ,

$$(1) \quad \alpha_{\tau}(E) \leq \sqrt{3}(1 - \|P_{\tau}EP_{\tau}\|_1^2)^{1/2}.$$

Now,

$$\begin{aligned} \|P_{\tau}EP_{\tau}\|_1 & \geq |\text{tr}(P_{\tau}EP_{\tau}C)| \geq \text{Retr}(EP_{\tau}CP_{\tau}) \\ & = \text{Retr}(EC) - \text{Retr}(EP_{\tau}CQ_{\tau}) - \text{Retr}(EQ_{\tau}CP_{\tau}) - \text{Retr}(EQ_{\tau}CQ_{\tau}) \\ & \geq k - \|E\|_1 \|P_{\tau}CQ_{\tau}\|_{\infty} - \|E\|_1 \|Q_{\tau}CP_{\tau}\|_{\infty} - \|E\|_1 \|Q_{\tau}CQ_{\tau}\|_{\infty} \\ & \geq k - \|CQ_{\tau}\|_{\infty} - \|Q_{\tau}C\|_{\infty} - \|CQ_{\tau}\|_{\infty} \\ & \geq k - 3 \max\{\|CQ_{\tau}\|_{\infty}, \|C^*Q_{\tau}\|_{\infty}\}. \end{aligned}$$

Let us fix $\eta \in (0, k)$. As C and C^* are compact operators, by Lemma 2.1 we have that there exists τ_0 such that for all τ with $Q_\tau \leq Q_{\tau_0}$,

$$3 \max\{\|CQ_\tau\|_\infty, \|C^*Q_\tau\|_\infty\} < \eta.$$

From inequality (1), for each $E \in S[C, k]$ and for every τ with $Q_\tau \leq Q_{\tau_0}$,

$$(2) \quad \alpha_\tau(E) \leq \sqrt{3}(1 - (k - \eta)^2)^{1/2}.$$

Consider an arbitrary sequence $\{A_n\}_{n \in \mathbb{N}} \subseteq S[C, k]$. From Lemma 2.3 we know that $\mathcal{B}_{\mathcal{E}_1}$ is weak* sequentially compact. We wish to estimate $\gamma(S[C, k])$ and so by Lemma 1.1 there is no loss of generality in assuming that $\{A_n\}_{n \in \mathbb{N}}$ is weak* convergent.

Note that since weak* and norm convergence coincide on finite-dimensional subspaces of a dual Banach space, we have

$$(3) \quad \|P_{\tau_0}(A_n - A_m)P_{\tau_0}\|_1 \rightarrow 0 \quad \text{as } n, m \rightarrow \infty.$$

Consequently, for each $n, m \in \mathbb{N}$,

$$\begin{aligned} \|A_n - A_m\|_1 &\leq \|P_{\tau_0}(A_n - A_m)P_{\tau_0}\|_1 + \alpha_{\tau_0}(A_n) + \alpha_{\tau_0}(A_m) \\ &\leq \|P_{\tau_0}(A_n - A_m)P_{\tau_0}\|_1 + 2\sqrt{3}(1 - (k - \eta)^2)^{1/2}, \end{aligned}$$

from inequality (2); and hence using (3) we see that

$$\inf_{n \neq m} \|A_n - A_m\|_1 \leq 2\sqrt{3}(1 - (k - \eta)^2)^{1/2}.$$

It is clear now that

$$\gamma(S[C, k]) \leq 2\sqrt{3}(1 - (k - \eta)^2)^{1/2},$$

and since $\eta \in (0, k)$ is arbitrary, we have

$$(4) \quad \gamma(S[C, k]) \leq 2\sqrt{3}(1 - k^2)^{1/2}.$$

Finally, if $\varepsilon \in (0, 2\sqrt{3})$ is given, choose

$$k = (1 - (\varepsilon/2\sqrt{3})^2)^{1/2}.$$

$k \in (0, 1)$ and inequality (4) gives us that for each $C \in S_{\mathcal{E}_\infty}$,

$$\gamma(S[C, k]) \leq \varepsilon. \quad \square$$

ACKNOWLEDGMENTS

My thanks to Ray Ryan and Paddy Dowling for their constructive suggestions during the preparation of this paper, and I thank Catherine for her careful typing.

REFERENCES

1. J. Arazy, *More on convergence in unitary matrix spaces*, Proc. Amer. Math. Soc. **83** (1981), 44–48.
2. D. van Dulst and Brailey Sims, *Fixed points of nonexpansive mappings and Chebyshev centers in Banach spaces with norms of type (KK)* , Banach Space Theory and its Applications (Proc. Bucharest 1981), Lecture Notes in Math., vol. 991, Springer-Verlag, Berlin, Heidelberg and New York.
3. D. van Dulst and V. de Valk (KK) , *properties, normal structure and fixed points of nonexpansive mappings in Orlicz sequence spaces*, Canad. J. Math. **38** (1986), 728–750.
4. I. C. Gohberg and M. G. Krein, *Introduction to the theory of linear nonselfadjoint operators*, Transl. Math. Mono., vol. 18 Amer. Math. Soc., Providence, RI, (1969).
5. R. Huff, *Banach spaces which are nearly uniformly convex*, Rocky Mountain J. Math. **10** (1980), 743–749.
6. M. I. Kadec, *On the connection between weak and strong convergences*, Dopovidі Akad. Nauk Ukrain **9** (1959), 949–952. (Ukrainian)
7. V. L. Klee, *Mappings into normed linear spaces*, Fund. Math. **49** (1960/61), 25–34.
8. A. T. Lau and P. F. Mah, *Quasi-normal structures for certain spaces of operators on a Hilbert space*, Pacific J. Math. **121** (1986), 109–118.
9. J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces. I*, A Series of Modern Surveys in Mathematics, vol. 92, Springer-Verlag, Berlin, Heidelberg, New York, 1977.
10. R. Schatten, *Norm ideals of completely continuous operators*, Springer-Verlag, Berlin, 1960.
11. B. Simon, *Convergence in trace ideals*, Proc. Amer. Math. Soc. **83** (1981), 39–43.
12. Brailey Sims, *Fixed points of nonexpansive maps on weak and weak* compact sets*, Lecture Notes, Queen's Univ., Kingston, 1982.
13. —, *The existence question for fixed points of nonexpansive maps*, Lecture Notes, Kent State Univ., 1986.

DEPARTMENT OF MATHEMATICS, KENT STATE UNIVERSITY, KENT, OHIO 44242