THE EXISTENCE OF ABSOLUTELY CONTINUOUS INVARIANT MEASURES FOR $C^{1+\varepsilon}$ JABLONSKI TRANSFORMATION IN R^n

YOU-SHI LOU

(Communicated by Kenneth R. Meyer)

ABSTRACT. Using a result of Rychlik, we present a sufficient condition for the existence of an absolutely continuous invariant measure for $C^{1+\varepsilon}$ Jablonski transformation in R^n .

1. INTRODUCTION

In 1973, Lasota and Yorke [1] proved a general sufficient condition for the existence of an absolutely continuous invariant measure for expanding piecewise C^2 transformation τ on the interval. Their proof uses the fact that the Frobenius–Perron operator induced by τ is quasicompact in the space of function of bounded variation.

For $I^n = [0, 1]^n$, let $\beta = \{D_1, D_2, \dots, D_p\}$ be a partition of I^n such that

$$\bigcup_{j=1}^p D_j = I^n, \quad D_j \cap D_k = \emptyset \quad \text{for } j \neq k.$$

A partition β is called a Jablonski partition if for any $1 \leq j \leq p$, D_j is a rectangle. A transformation $\tau: I^n \rightarrow I^n$ is called a Jablonski transformation if it is given by the formula

$$\begin{aligned} \tau(x_1, \dots, x_n) &= (\varphi_{1j}(x_1), \dots, \varphi_{nj}(x_n)), \\ \text{for } (x_1, \dots, x_n) &\in D_j, \quad j = 1, \dots, p, \quad p < \infty, \end{aligned}$$

where $D_j = \prod_{i=1}^n [a_{ij}, b_{ij})$, $\varphi_{ij}: [a_{ij}, b_{ij}) \rightarrow [0, 1]$ and we write

$$[a_{ij}, b_{ij}) = \begin{cases} [a_{ij}, b_{ij}), & \text{if } b_{ij} < 1 \\ [a_{ij}, b_{ij}], & \text{if } b_{ij} = 1. \end{cases}$$

In 1983, Jablonski [2] proved the existence of an absolutely continuous invariant measure for the above τ under the conditions that for any $1 \leq i \leq n$,

Received by the editors December 18, 1988.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 28A65; Secondary 58F11.

©1990 American Mathematical Society
0002-9939/90 \$1.00 + \$.25 per page

$1 \leq j \leq p$, φ_{ij} is a C^2 -function and

$$\lambda = \inf_{i,j} \left\{ \inf_{[a_{ij}, b_{ij}]} |\varphi'_{ij}| \right\} > 1.$$

The proof in [2] uses bounded variation, which is a complicated notion in higher dimension.

In this paper we present a sufficient condition for the existence of an absolutely continuous invariant measure for Jablonski transformations, which does not use bounded variation in R^n . The proof is much easier and allows us to weaken the $\varphi_{ij} \in C^2$ condition to $\varphi_{ij} \in C^{1+\varepsilon}$ for some $0 < \varepsilon < 1$. There is, however, a new condition $\bar{r}^{(t)}/\lambda^{nt} < 1$ for some integer $t \geq 1$ in this paper, where $\bar{r}^{(t)}$ is a number which depends on the defining partition of the transformation.

2. RYCHLIK'S SUFFICIENT CONDITION

Let (X, Σ, m) be a Lebesgue space with a σ -algebra Σ and a probability measure m . Let $\tau: X \rightarrow X$ be a measurable, nonsingular transformation and g_τ be the absolute value of the reciprocal of the Jacobian of τ . Now we can express the Frobenius–Perron operator in the following way:

$$P_\tau f(x) = \sum_{y \in \tau^{-1}(x)} g_\tau(y) f(y), \quad x \in X.$$

Let β be a partition of X which is a generator for τ , i.e. $V_{k=0}^\infty \tau^{-k}(\beta) = \varepsilon$, where ε is a partition into points. For any positive integer l , let β_l be a partition of X which is a generator for τ^l . Let $g = g_1 = g_\tau$ and $g_l = g_{\tau^l}$. For every $A \in \Sigma$, we define $\beta(A) = \{B \in \beta: m(B \cap A) > 0\}$.

Condition 1 (Distortion condition). *There exists a positive number b such that for any $l \geq 1$ and any $B \in \beta_l$, we have $\sup_B g_l \leq b \inf_B g_l$.*

Condition 2 (Localization condition). *There exist $\varepsilon > 0$ and $0 < \gamma < 1$ such that for any $l \geq 1$ and $B \in \beta_l$, $m(\tau^l B) < \varepsilon$ implies $\sum_{B' \in \beta(\tau^l B)} \sup_{B'} g \leq \gamma$.*

Condition 3 (Boundedness condition). $\sum_{B \in \beta} \sup_B g < \infty$.

Under these three conditions Rychlik [3] proved the existence of fixed point for P_τ .

Theorem 1. *Assume Conditions 1, 2, and 3 are satisfied. Then the sequence $\{P_\tau^l 1\}_{l=1}^\infty$ is bounded in $L^\infty(m)$, and the averages $(1/l) \sum_{j=0}^{l-1} P_\tau^j 1$ converge in $L^1(m)$ to some $\varphi \in L^\infty(m)$ such that $P_\tau \varphi = \varphi$.*

3. MAIN RESULT

Let $X = I^n$. For any $x, y \in I^n$, we define

$$d(x, y) = \left(\sum_{i=1}^n (x_i - y_i)^2 \right)^{1/2},$$

where $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_n)$.

Lemma 1. *If there exists a positive number M such that for any $l \geq 1$ $B \in \beta_l$ and $x, y \in B$ we have $g_l(x)/g_l(y) \leq M$, then Condition 1 is satisfied.*

Lemma 2. *Let $\tau: I^n \rightarrow I^n$ be a nonsingular piecewise $C^{1+\varepsilon}$ transformation, i.e. there exists a constant $\bar{c}$ such that for any $x, y \in D_j$, $j = 1, \dots, p$, $|J(x) - J(y)| \leq \bar{c}d(x, y)^\varepsilon$, where J is the absolute value of the Jacobian of τ . If there exist constants $\lambda > 1$ and $c > 0$ such that for $l \geq 1$, $B \in \beta_l$ and $x, y \in B$ we have $d(\tau^l x, \tau^l y) \geq c\lambda^l d(x, y)$, then there exists a constant M such that $g_l(x)/g_l(y) \leq M$ and Condition 1 is satisfied.*

Proof. For $x, y \in B \in \beta_l$, $k = 0, 1, \dots, l-1$, we have

$$d(\tau^k x, \tau^k y) \leq c^{-1} \lambda^{-(l-k)} d(\tau^l x, \tau^l y) \leq c^{-1} \lambda^{-(l-k)} \sqrt{n}.$$

Then, using the fact that τ is piecewise $C^{1+\varepsilon}$, we have

$$\begin{aligned} \frac{J(\tau^k y)}{J(\tau^k x)} &= \left| 1 + \frac{J(\tau^k y) - J(\tau^k x)}{J(\tau^k x)} \right| \leq 1 + \frac{|J(\tau^k y) - J(\tau^k x)|}{J(\tau^k x)} \\ &\leq 1 + \frac{\bar{c}d(\tau^k x, \tau^k y)^\varepsilon}{c^n \lambda^n} \leq 1 + \frac{\bar{c}\sqrt{n}^\varepsilon}{c^{n+\varepsilon} \lambda^n} \lambda^{-(l-k)\varepsilon}, \quad k = 0, 1, \dots, l-1, \end{aligned}$$

and

$$\begin{aligned} \frac{g_l(x)}{g_l(y)} &= \frac{J(\tau^{l-1} y) J(\tau^{l-2} y) \cdots J(\tau y) J(y)}{J(\tau^{l-1} x) J(\tau^{l-2} x) \cdots J(\tau x) J(x)} \\ &\leq \prod_{k=0}^{l-1} \left(1 + \frac{\bar{c}\sqrt{n}^\varepsilon}{c^{n+\varepsilon} \lambda^n} \lambda^{-(l-k)\varepsilon} \right) = \prod_{i=1}^l \left(1 + \frac{\bar{c}\sqrt{n}^\varepsilon}{c^{n+\varepsilon} \lambda^n} \lambda^{-\varepsilon i} \right) \\ &\leq \prod_{i=1}^{\infty} \left(1 + \frac{\bar{c}\sqrt{n}^\varepsilon}{c^{n+\varepsilon} \lambda^n} \lambda^{-\varepsilon i} \right) = M. \quad \square \end{aligned}$$

Let $\tau: I^n \rightarrow I^n$ be a Jablonski transformation:

$$\tau(x) = (\varphi_{1j}(x_1), \dots, \varphi_{nj}(x_n)), \quad x \in D_j,$$

where for any $1 \leq j \leq p$, D_j is a rectangle, $D_j \cap D_k = \emptyset$ for $j \neq k$ and $\bigcup_{j=1}^p D_j = I^n$. $\beta = \{D_1, \dots, D_p\}$. For any $l \geq 1$, let $\beta_l = \{D_1^{(l)}, \dots, D_p^{(l)}\}$.

Then every $D_j^{(l)}$ is a rectangle and τ^l is a Jablonski transformation.

Let $L_i(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$ be the straight line parallel to the i th axis and having all the coordinates but the i th one fixed and equal to $x_1, \dots, x_{i-1}, \dots, x_n$. Let $r_i^{(l)}(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n)$ be the number of $D_j^{(l)}$'s

FIGURE 1.

for which $D_j^{(l)} \cap L_i(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n) \neq \emptyset$. For any $1 \leq i \leq n$ let

$$r_i^{(l)} = \sup_{0 \leq x_k \leq 1, 1 \leq k \leq n, k \neq i} r_i^{(l)}(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_n), \quad r_i = r_i^{(1)},$$

$$r^{(l)} = r_1^{(l)} r_2^{(l)} \cdots r_n^{(l)} = \prod_{i=1}^n r_i^{(l)}, \quad r = r^{(1)},$$

$$\bar{r}_i^{(l)} = r^{(l)} / r_i^{(l)} = r_1^{(l)} r_2^{(l)} \cdots r_{i-1}^{(l)} r_{i+1}^{(l)} \cdots r_n^{(l)} = \prod_{j=1, j \neq i}^n r_j^{(l)}, \quad \bar{r}_i = \bar{r}_i^{(1)},$$

$$\bar{r}^{(l)} = \max_{1 \leq i \leq n} \bar{r}_i^{(l)}, \quad \bar{r} = \bar{r}^{(1)},$$

i.e. $r_i^{(l)}$ is the maximal number of parts of β_l in x_i direction. For example $n = 2$, β_l is shown in Figure 1 and we have: $r_1^{(l)} = 4$, $r_2^{(l)} = 5$, $r^{(l)} = r_1^{(l)} r_2^{(l)} = 20$, $\bar{r}_1^{(l)} = r^{(l)} / r_1^{(l)} = 5$, $\bar{r}_2^{(l)} = r^{(l)} / r_2^{(l)} = 4$, $\bar{r}^{(l)} = \max(\bar{r}_1^{(l)}, \bar{r}_2^{(l)}) = 5$.

Lemma 3. Let $\tau: I^n \rightarrow I^n$ be a Jablonski transformation and assume that $\lambda = \inf_{i,j} \inf_{0 \leq x_i \leq 1} |\phi'_{ij}| > 1$. If $\bar{r} \lambda^{-n} < 2^{-1}$, then Condition 2 is satisfied.

Proof. Take $\varepsilon > 0$ such that $\varepsilon^{1/n} < d = \min_{1 \leq j \leq p} d_j$, where d_j is the minimal width of D_j , and $0 < \gamma < 1$ such that $0 < 2\bar{r} \lambda^{-n} < \gamma < 1$. Since $|\phi'_{ij}| \geq \lambda$, we have $g \leq \lambda^{-n}$. Since $d > \varepsilon^{1/n}$, for any $B \in \beta_l$, $m(\tau^l B) < \varepsilon$ means at least one of the widths of $\tau^l B$ is smaller than $\varepsilon^{1/n} < d$ and the number of $\beta(\tau^l B)$ is at most $2\bar{r}$. Hence

$$\sum_{B' \in \beta(\tau^l B)} \sup_{B'} g < \frac{\text{the number of } \beta(\tau^l B)}{\lambda^n} \leq \frac{2\bar{r}}{\lambda^n} < \gamma. \quad \square$$

Lemma 4. Let $\beta = \{D_1, \dots, D_p\}$, $p < \infty$. If for any D_j , $\sup_{D_j} g < \infty$, then Condition 3 is satisfied.

Proof. Let $M = \max_{1 \leq j \leq p} \sup_{D_j} g$. Then we have $\sum_{B \in \beta} \sup_B g \leq pM < \infty$, which is Condition 3. $\square$

By Lemma 2, 3, and 4, and Theorem 1, we have:

Theorem 2. Let $\tau: I^n \rightarrow I^n$ be a Jablonski transformation with a finite partition $\beta = \{D_1, \dots, D_p\}$, and let $\lambda = \inf_{i,j} \inf_{0 \leq x_i \leq 1} |\phi'_{ij}| > 1$. If for any i and j , $\phi_{ij} \in C^{1+\varepsilon}$, $0 < \varepsilon < 1$ and $\bar{r}/\lambda^n < \frac{1}{2}$ then the sequence $\{P_\tau^l 1\}_{l=1}^\infty$ is bounded in $L^\infty(m)$ and the averages $(1/l) \sum_{j=0}^{l-1} P_\tau^j$ converge in $L^1(m)$ to some $\varphi \in L^\infty(m)$ such that $P_\tau \varphi = \varphi$.

Lemma 5. If φ is a fixed point of $P_{\tau^t} = P_\tau^t$ for some positive integer t then $\psi = (1/t)(\varphi + P_\tau \varphi + \dots + P_\tau^{t-1} \varphi)$ is a fixed point of P_τ .

For a Jablonski transformation τ and any $l \geq 1$ we have $r_i^{(l)} \leq r_i^l$, $r^{(l)} \leq r^l$, $\bar{r}_i^{(l)} \leq \bar{r}_i^l$, $\bar{r}^{(l)} \leq \bar{r}^l$, $g_l < \lambda^{-nl}$. We therefore have:

Theorem 3. Let $\tau: I^n \rightarrow I^n$ be a Jablonski transformation with a finite partition $\beta = \{D_1, \dots, D_p\}$, and assume $\lambda = \inf_{i,j} \inf_{0 \leq x_i \leq 1} |\phi'_{ij}| > 1$. If for any i and j $\phi_{ij} \in C^{1+\varepsilon}$, with $0 < \varepsilon < 1$ and $\bar{r} \cdot \lambda^{-n} < 1$, then there exists a function $\psi \in L^\infty(m)$ such that $P_\tau \psi = \psi$.

Proof. Since $\bar{r} \cdot \lambda^{-n} < 1$, we can take an integer $t \geq 1$ such that $(\bar{r} \cdot \lambda^{-n})^t < \frac{1}{2}$. Since τ^t satisfies all the conditions of Theorem 2, there exists a function $\varphi \in L^\infty(m)$ such that $P_{\tau^t} \varphi = P_\tau^t \varphi = \varphi$. By Lemma 5

$$\psi = \frac{1}{t} \sum_{s=0}^{t-1} P_\tau^s \varphi$$

satisfies $P_\tau \psi = \psi$. $\square$

For $n = 1$, we have $\bar{r} = 1$. By virtue of Theorem 3, we have:

Theorem 4. Let $\tau: I \rightarrow I$ be a piecewise expanding and piecewise $C^{1+\varepsilon}$ transformation with $0 < \varepsilon < 1$. Then there exists a function $\varphi \in L^\infty(m)$ such that $P_\tau \varphi = \varphi$.

Theorem 5. Let $\tau: I^n \rightarrow I^n$ be a Jablonski transformation with a finite partition $\beta = \{D_1, \dots, D_p\}$, and $\lambda = \inf_{i,j} \inf_{0 \leq x_i \leq 1} |\phi'_{ij}| > 1$. If for any i and j , $\phi_{ij} \in C^{1+\varepsilon}$, $0 < \varepsilon < 1$, and there exists a positive integer $t \geq 1$ such that $\bar{r}^{(t)}/\lambda^{nt} < 1$ then there exists a function $\psi \in L^\infty(m)$ such that $P_\tau \psi = \psi$.

Proof. Since τ^t satisfies all the conditions of Theorem 3, there exists a function $\varphi \in L^\infty(m)$ such that $P_{\tau^t} \varphi = P_\tau^t \varphi = \varphi$. By Lemma 5

$$\psi = \frac{1}{t} \sum_{s=0}^{t-1} P_\tau^s \varphi \quad \text{satisfies } P_\tau \psi = \psi. \quad \square$$

4. GENERALIZED-JABLONSKI TRANSFORMATIONS

A transformation $\tau: I^n \rightarrow I^n$ is called a generalized-Jablonski transformation if it is given by the formula

$$\begin{aligned} \tau(x_1, \dots, x_n) &= (\varphi_{1j}(x_1), \varphi_{2j}(x_1, x_2), \dots, \varphi_{nj}(x_1, x_2, \dots, x_n)), \\ (x_1, \dots, x_n) &\in D_j, \quad j = 1, \dots, p, \quad p < \infty, \end{aligned}$$

the image of any hyperplane in I^n is a collection of segments of hyperplanes in I^n and the inverse image of any hyperplane in I^n is a collection of segments of hyperplanes in I^n , where D_j is the same as for a Jablonski transformation and $\varphi_{ij}(x_1, \dots, x_i): \prod_{k=1}^i [a_{kj}, b_{kj}] \rightarrow [0, 1]$.

Taking $c = 1$ in Lemma 2, we have:

Lemma 6. *Let $\tau: I^n \rightarrow I^n$ be a nonsingular piecewise $C^{1+\varepsilon}$ transformation. If there exists a constant $\lambda > 1$ such that for any $B \in \beta$ and $x, y \in B$ we have*

$$(*) \quad d(\tau x, \tau y) \geq \lambda d(x, y),$$

then Condition 1 holds.

Lemma 7. *Let $\tau: I^n \rightarrow I^n$ be a generalized-Jablonski transformation. If $(*)$ is satisfied with $\lambda > 1$ and $2\bar{r}/\lambda^n < 1$, then Condition 2 holds.*

Proof. Let $\varepsilon > 0$ such that $\varepsilon^{1/n} < d/2^{n-1}$, where $d = \min_{1 \leq j \leq p} d_j$ and d_j is the minimal width of D_j , and $0 < \gamma < 1$ such that $0 < 2\bar{r}\lambda^{-n} < \gamma < 1$. Since for any $x, y \in B \in \beta$, $d(\tau x, \tau y) \geq \lambda d(x, y)$, we have $g \leq \lambda^{-n}$. Since $d > 2^{n-1} \cdot \varepsilon^{1/n}$, for any $B \in \beta_l$, $m(\tau^l B) < \varepsilon$ means at least one of the maximal length of $\tau^l B$ on x_i direction is smaller than $2^{n-1} \cdot \varepsilon^{1/n} < d$ and the number of $\beta(\tau^l B)$ is at most $2\bar{r}$. Hence

$$\sum_{B' \in \beta(\tau^l B)} \sup_{B'} g \leq \frac{(\text{the number of } \beta(\tau^l B))}{\lambda^n} \leq \frac{2\bar{r}}{\lambda^n} < \gamma. \quad \square$$

By Lemmas 4, 6, 7, and Theorem 1, we have:

Theorem 6. *Let $\tau: I^n \rightarrow I^n$ be a nonsingular, piecewise $C^{1+\varepsilon}$ generalized-Jablonski transformation with a finite partition $\beta = \{D_1, \dots, D_p\}$. If for any $x, y \in B \in \beta$, $d(\tau x, \tau y) \geq \lambda d(x, y)$ for some constant $\lambda > 1$ and $2\bar{r}/\lambda^n < 1$ then the sequence $\{P_\tau^l 1\}_{l=1}^\infty$ is bounded in $L^\infty(m)$ and the averages $(1/l) \sum_{j=0}^{l-1} P_\tau^j 1$ converge in $L^1(m)$ to some $\varphi \in L^\infty(m)$ such that $P_\tau \varphi = \varphi$.*

5. EXAMPLE

In Theorem 5 we need a condition $\bar{r}^{(t)}/\lambda^{nt} < 1$ for some integer $t \geq 1$. Now we show that, in general, $\lambda > 1$ does not imply this condition.

(1) Consider the partition $\beta = \{D_1, D_2, \dots, D_9\}$ shown in Figure 2 for that unit square in two-dimensions. Let τ be a Jablonski transformation defined as

Figure 2.

follows. For $j = 1, 2, 4$ and $i = 1, 2$, φ_{ij} is a linear function which is onto $[0, 1]$ with $|\varphi'_{ij}| = 3$. For $j = 3, 5, 6, 7, 8, 9$ and $i = 1, 2$, $\varphi_{ij} \in C^{1+\varepsilon}$ with $\inf_{x_i} |\varphi'_{ij}| = 1.1$. Therefore, $\lambda = 1.1$, $\lambda^2 = 1.21$, $\bar{r} = 3$, $\bar{r}^{(2)} \geq 2\bar{r} + 1 > 2\bar{r}$. For $l > 1$, $\bar{r}^{(l)} > 2^{l-1}\bar{r} = 3 \cdot 2^{l-1} = 1.5 \cdot 2^l$. This means that for any positive integer $t \geq 1$,

$$\frac{\bar{r}^{(t)}}{\lambda^{nt}} = \frac{\bar{r}^{(t)}}{\lambda^{2t}} = \frac{\bar{r}^{(t)}}{1.21^t} > 1.5 \cdot \left(\frac{2}{1.21}\right)^t > 1.$$

(2) For $n = 2$, and the partition $\beta = \{D_1, \dots, D_9\}$ is the same as in (1), assume that for every i and j , $\varphi_{ij} \in C^{1+\varepsilon}$, $\lambda = \inf_{i,j} \inf_{x_i} |\varphi'_{ij}| > \sqrt{3}$. Then we have

$$\frac{\bar{r}}{\lambda^n} < \frac{3}{(\sqrt{3})^2} = 1.$$

For example, let $\varphi(x) = x^{3/2} + 1.74x$. Then

$$\varphi'(x) = \frac{3}{2}x^{1/2} + 1.74 \geq 1.74 > \sqrt{3}$$

for $x \geq 0$ and $\varphi(x)$ maps $[0, \frac{1}{3}]$ onto $[0, 0.77245]$. We let

$$\varphi_{ij}(x_i) = \begin{cases} \varphi(x_i), & \text{if } [a_{ij}, b_{ij}] = [0, \frac{1}{3}], \\ 1.2\varphi(x_i - \frac{1}{3}), & \text{if } [a_{ij}, b_{ij}] = [\frac{1}{3}, \frac{2}{3}], \\ \varphi(x_i - \frac{2}{3}) + .2, & \text{if } [a_{ij}, b_{ij}] = [\frac{2}{3}, 1], \end{cases}$$

and we have $\varphi_{ij}(x_i) \in C^{1+1/2}$, $\lambda = 1.74 > \sqrt{3}$.

By Theorem 3, there exists a function $\psi \in L^\infty(m)$ such that $P_\tau \psi = \psi$.

REFERENCES

1. A. Lasota and J. A. Yorke, *On the existence of invariant measures for piecewise monotonic transformation*, Trans. Amer. Math. Soc. **186** (1973), 481–488.
2. M. Jablonski, *On invariant measures for piecewise C^2 -transformation of the n -dimensional cube*, Ann. Polon. Math. **XLIII** (1983), 185–195.

3. Marek Ryszard Rychlik, *Invariant measures and variational principle for Lozi mappings* (preprint).
4. Ya. G. Sinai, *Dopolnenie k knige G. M. Zaslavskogo. Statisticheskaya neobratimost' v nelineinykh sistemakh*, addition to the book by G. M. Zaslavskii, Statistical irreversibility in non-linear systems, Nauka (1970), 124–143.

DEPARTMENT OF MATHEMATICS, CONCORDIA UNIVERSITY, 7141 SHERBROOKE STREET WEST,
MONTREAL H4B 1R6 CANADA