TOPOLOGICAL RATIONALIZATION OF A CLASS OF MEROMORPHIC FUNCTIONS

A. MEZIANI

(Communicated by Clifford J. Earle, Jr.)

ABSTRACT. We prove that a class of germs at $0 \in \mathbb{C}^2$ of meromorphic functions can be transformed, via homeomorphisms, into rational functions.

0. INTRODUCTION

We prove in this note that if $M(x, y)$ is the germ at $0 \in \mathbb{C}^2$ of a pure meromorphic function whose blow-up $\tilde{M} = M \circ E$, where E is the blow-up of $\mathbb{C}^2$ at 0 , has no essential singularities on $E^{-1}(0)$, then M is topologically conjugate to a rational function $R(x, y)$. That is, there is a homeomorphism Φ of a neighborhood of $0 \in \mathbb{C}^2$ such that $M \circ \Phi = R$.

D. Cerveau and J. F. Mattei have studied [1] finite determination for multiform functions. In particular, they prove rationalization for germs of meromorphic functions with critical set reduced to $\{0\}$. They also show that germs at $0 \in \mathbb{C}^2$ of holomorphic functions are conjugate to polynomials.

We begin this note by recalling briefly the blow-up of $\mathbb{C}^2$ at 0 , and then we define the class of meromorphic functions that we will be dealing with. Next, we prove an intermediate result: a particular conjugation of germs of analytic varieties. Finally, we state and prove the main result.

1. DEFINITIONS

In this section, we recall very briefly the blow-up of $\mathbb{C}^2$ at 0 (see [4]) and define a class of meromorphic functions, called here class A. Some examples will follow.

The blow-up of $\mathbb{C}^2$ at 0 is the subset of $\mathbb{C}^2 \times \mathbb{CP}^1$, where $\mathbb{CP}^1$ is the complex projective space of dimension 1, given as

$$\mathbb{C}_0^2 = \{(p, [p]) \in (\mathbb{C}^2 - \{0\}) \times \mathbb{CP}^1\} \cup (\{0\} \times \mathbb{CP}^1),$$

where $[\]: \mathbb{C}^2 - \{0\} \rightarrow \mathbb{CP}^1$ is the quotient map. It is shown, in [4] for example, that $\mathbb{C}_0^2$ is a two-dimensional complex manifold. Moreover, $\mathbb{C}_0^2$ is covered by

Received by the editors April 3, 1989.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 32A20.

two charts (x, t) , (s, y) (t, s are the canonical coordinates of $\mathbf{CP}^1$) related by

$$s = \frac{1}{t} \quad \text{and} \quad y = xt.$$

The first projection $E: \mathbf{C}_0^2 \mapsto \mathbf{C}^2$, called the blow-up map, is given in the charts (x, t) , (s, y) by

$$E(x, t) = (x, xt) \quad \text{and} \quad E(s, y) = (sy, y).$$

E is a biholomorphism from $\mathbf{C}_0^2 - E^{-1}(0)$ onto $\mathbf{C}^2 - \{0\}$. The exceptional divisor $E^{-1}(0)$ is identified with $\mathbf{CP}^1$.

A meromorphic function M on a two-dimensional complex manifold $\mathcal{M}$ is said to have an *essential singularity* at $p_0 \in \mathcal{M}$ if, near p_0 , the function M is expressed as a quotient α/β , with α, β germs at p_0 of coprime holomorphic functions and $\alpha(p_0) = \beta(p_0) = 0$. Equivalently, M has an essential singularity at p_0 if, in a neighborhood of p_0 , all the level sets of M pass through p_0 . In this case, as in [1], we say that M is a *pure meromorphic* function at p_0 .

Definition 1.1. A germ at $0 \in \mathbf{C}^2$ of a pure meromorphic function $M(x, y)$ is said to be in the class A if the map $\widetilde{M} = M \circ E: \mathbf{C}_0^2 \mapsto \overline{\mathbf{C}}$, defined in a neighborhood of $E^{-1}(0) = \mathbf{CP}^1 \subset \mathbf{C}_0^2$, has no essential singularities.

Remark 1.1. It is easy to see, as a consequence of the desingularization of foliations defined by level sets of meromorphic functions (see [3] for example), that if M is the germ at $0 \in \mathbf{C}^2$ of a pure meromorphic function, then there is a proper holomorphic map $\Pi: \mathcal{M} \mapsto \mathbf{C}^2$, obtained by composition of blow-ups, such that the lift $M \circ \Pi$ of M to $\mathcal{M}$ has no essential singularities. Hence, the class A is the set of all meromorphic functions for which $\mathcal{M} = \mathbf{C}_0^2$.

Example 1. The function

$$M(x, y) = \frac{x^3 + xy^2 + \alpha(x, y)}{y^3 + \beta(x, y)},$$

where α, β are holomorphic functions of order ≥ 4 , is in the class A . The blow-up of M is given in the charts (x, t) , (s, y) respectively by

$$\widetilde{M}(x, t) = \frac{1 + t^2 + (\alpha(x, tx)/x^3)}{y^3 + (\beta(x, tx)/x^3)}, \quad \widetilde{M}(s, y) = \frac{s^3 + s + (\alpha(sy, y)/y^3)}{1 + (\beta(sy, y)/y^3)}.$$

So $\widetilde{M}$ has no essential singularities on $\mathbf{CP}^1 = \{x = 0\} \cup \{y = 0\}$.

Example 2. The meromorphic function $M(x, y) = (y^2 + x^3)/xy$ is not in the class A . Its blow-up $\widetilde{M}(x, t) = (t^2 + x)/t$ has an essential singularity at $(x = 0, t = 0)$. A blow-up of $\mathbf{C}_0^2$ at $(x = 0, t = 0)$ will produce a meromorphic function with no essential singularities.

Remark 1.2. The functions of the form x^r/y^s , where r, s are positive integers and $r > s$, are not in the class A . However, we characterized in [3] all meromorphic functions that are conjugate to the rational functions of the form $R(x^r/y^s)$, where R is a rational function on $\overline{\mathbb{C}}$.

The following proposition characterizes the class A .

Proposition 1.1. Let $M(x, y) = \alpha(x, y)/\beta(x, y)$ be the germ at $0 \in \mathbb{C}^2$ of a pure meromorphic function and let

$$\alpha(x, y) = \sum_{i \geq n} P_i(x, y) \quad \text{and} \quad \beta(x, y) = \sum_{j \geq m} Q_j(x, y)$$

be the Taylor expansions of α and β (P_i and Q_j are homogeneous polynomials with respective degrees i and j). Then M is in the class A if and only if $n = m$ and P_n, Q_n are coprime.

Proof. We first show that if $n = m$ and P_n, Q_n are coprime, then M is in the class A . The blow-up $\widetilde{M} = M \circ E$ of M is given in the charts $(x, t), (s, y)$ by

$$\begin{aligned} \widetilde{M}(x, t) &= \frac{P_n(1, t) + xP_{n+1}(1, t) + \dots}{Q_n(1, t) + xQ_{n+1}(1, t) + \dots}, \\ \widetilde{M}(s, y) &= \frac{P_n(s, 1) + yP_{n+1}(s, 1) + \dots}{Q_n(s, 1) + yQ_{n+1}(s, 1) + \dots}. \end{aligned}$$

Since $P_n(1, t)$ and $Q_n(1, t)$ (resp. $P_n(s, 1)$ and $Q_n(s, 1)$) have no common root, then $\widetilde{M}$ has no essential singularities on $E^{-1}(0)$.

Conversely, let us show that if M is in the class A , then $n = m$ and P_n, Q_n are coprime. By contradiction, first if $n = m + k > m$, then

$$\begin{aligned} \widetilde{M}(x, t) &= \frac{x^k (P_{m+k}(1, t) + xP_{m+k+1}(1, t) + \dots)}{Q_m(1, t) + xQ_{m+1}(1, t) + \dots}, \\ \widetilde{M}(s, y) &= \frac{y^k (P_{m+k}(s, 1) + yP_{m+k+1}(s, 1) + \dots)}{Q_m(s, 1) + yQ_{m+1}(s, 1) + \dots}. \end{aligned}$$

So, if t_0 (resp. s_0) is a root of $Q_m(1, t)$ (resp. $Q_m(s, 1)$), then the point $(x = 0, t = t_0)$ (resp. $(s = s_0, y = 0)$) would be an essential singularity of $\widetilde{M}$. The second eventuality, $n = m$ and P_n, Q_n are not coprime, also cannot occur. Indeed, $P_n(1, t)$ and $Q_n(1, t)$ would have a common root t_0 , and then

$$\begin{aligned} \widetilde{M}(x, t) &= \frac{P_n(1, t) + xP_{n+1}(1, t) + \dots}{Q_n(1, t) + xQ_{n+1}(1, t) + \dots} \\ &= \frac{(t - t_0)P_n^*(1, t) + xP_{n+1}(1, t) + \dots}{(t - t_0)Q_n^*(1, t) + xQ_{n+1}(1, t) + \dots} \end{aligned}$$

would have an essential singularity at $(x = 0, t = t_0)$.

2. AN INTERMEDIATE RESULT

We prove in this section the existence of a particular homeomorphism between two germs of analytic varieties that coincide to high order at $0 \in \mathbb{C}^2$. We will denote by C_c the complex line in $\mathbb{C}^2$ given by $y = cx$, where c is a complex number.

Proposition 2.1. *Let $f(x, y)$ be the germ at $0 \in \mathbb{C}^2$ of a holomorphic function and let $f_k(x, y)$ be its k th jet. Define*

$$V(f) = \{(x, y) ; f(x, y) = 0\} \quad \text{and} \quad V(f_k) = \{(x, y) ; f_k(x, y) = 0\}.$$

Then, for k large enough, there is a germ of a homeomorphism Φ at $0 \in \mathbb{C}^2$ such that

$$\Phi(V(f)) = V(f_k) \quad \text{and} \quad \Phi(C_c) = C_c$$

for every $c \in \mathbb{C}$.

The proof of this proposition is based upon some lemmas.

Lemma 2.1. *Consider a family $D(p_i, r_i)$, $i = 1, \dots, N$, of N disjoint discs in the plane $\mathbb{R}^2$, with respective centers p_i and radii r_i . For every i , let q_i be a point in $D(p_i, r_i)$. Then there is a homeomorphism $\phi: \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that:*

$$\begin{aligned} \phi(p) &= p \quad \text{if } p \notin \bigcup_{i=1}^N D(p_i, r_i) \\ \phi(p_i) &= q_i \quad \text{for } i = 1, \dots, N. \end{aligned}$$

We leave the elementary proof of this result to the reader.

Lemma 2.2. *Let $f(x, y)$ be the germ at $0 \in \mathbb{C}^2$ of an irreducible holomorphic function of order n . Then, we can assume, after a linear change of coordinates, that*

$$f(x, y) = y^n + P_{n+1}(x, y) + \dots + P_{n+k}(x, y) + \dots,$$

where the P_{n+j} 's are homogeneous polynomials of degree $n+j$. Furthermore, if $n > 1$, then there is a k such that $P_{n+k}(x, 0)$ is not identically zero.

Proof. The Taylor expansion of f is $f(x, y) = \sum_{j \geq n} P_j(x, y)$, where the P_j 's are homogeneous polynomials of degree j . Recall from [4] that the irreducibility of f implies that the tangent cone of the variety $\{(x, y) ; f(x, y) = 0\}$ is reduced to a point. That is, the set of points on $\mathbb{CP}^1$ on which $P_n(1, t)$ or $P_n(s, 1) = 0$ is reduced to one point. We can assume, after a linear change of coordinates if necessary, that $s = 0$ is not a root of $P_n(s, 1)$. So, the irreducibility of f implies that $P_n(1, t) = (t - t_0)^i$ for some $t_0 \in \mathbb{C}$ and $i \leq n$. By a linear change of coordinates, we can assume $t_0 = 0$. Now, if $n > i$, then $P_n(s, 1) = s^{n-i}$ would have a root at $s = 0$, and this is impossible. So $i = n$ and $P_n(x, y) = y^n$. Finally, if $n > 1$ and $P_{n+j}(x, 0) = 0$ for every

$j \geq 1$, then y would be a factor of f and f would be reducible. The lemma is proved. $\square$

Lemma 2.3. *Let $f(x, y) = y^n + P_{n+1}(x, y) + \dots + P_{n+k}(x, y) + \dots$, be an irreducible holomorphic function and k be the first integer such that $P_{n+k}(x, 0) \neq 0$. Then there is a positive constant K such that, for every $c \in \mathbb{C}$, $0 < |c| < K$, the variety $V(f) = \{f = 0\}$ intersects the line $\mathbb{C}_c$ at 0 and k other points $p^1(c), \dots, p^k(c)$ which vary continuously with c and $p^1(0) = \dots = p^k(0) = 0$.*

Proof. Let c be fixed in $\mathbb{C}$. Then

$$V(f) \cap \mathbb{C}_c = \{(x, cx) \in \mathbb{C}^2; f(x, cx) = 0\},$$

i.e., $V(f) \cap \mathbb{C}_c$ is given by the equation

$$x^n (c^n + P_{n+1}(1, c)x + \dots + P_{n+k}(1, c)x^k + \dots) = 0.$$

Since $P_{n+j}(1, 0) = 0$ if $j < k$ and $P_{n+k}(1, 0) \neq 0$, then similar arguments as that used in the proof of Theorem 4.13 of Chapter 2 in [2] and the irreducibility of f show that the equation

$$c^n + P_{n+1}(1, c)x + \dots + P_{n+k}(1, c)x^k + \dots = 0$$

has k distinct and continuous roots $p^1(c), \dots, p^k(c)$, provided that $c \neq 0$ is sufficiently small and $p^1(0) = \dots = p^k(0) = 0$. $\square$

Proof of the proposition. We start by proving the proposition in the case where the tangent cone of $V(f)$ is reduced to one point. Let $K > 0$ be such that, for every $c \in \mathbb{C}$, $0 < |c| < K$, we have

$$V(f) \cap \mathbb{C}_c = \{0, p^1(c), \dots, p^k(c)\}$$

(see Lemma 2.3). Consider the homotopy

$$g_t = f + t(f_l - f), \quad 0 \leq t \leq 1,$$

where f_l is the l th jet of f . Since f and g_t coincide to order l at 0, then, for l sufficiently large, we also have

$$V(g_t) \cap \mathbb{C}_c = \{0, p_t^1(c), \dots, p_t^k(c)\},$$

and the functions $p_t^j(c)$'s are continuous in c and t . So, for every j , we can find a continuous function $r_j(c)$, $r_j(c) > 0$ and $r_j(0) = 0$ such that

$$\{p_t^j(c), 0 \leq t \leq 1\} \subset D(p^j(c), r_j(c))$$

and

$$D(p^j(c), r_j(c)) \cap D(p^i(c), r_i(c)) = \emptyset \quad \text{if } j \neq i.$$

Next, for c fixed, $|c| < K$, consider, as in Lemma 2.1, a homeomorphism ϕ_c of the complex line $\mathbb{C}_c$ such that

$$\begin{aligned}\phi_c(p) &= p \quad \text{if } p \in \mathbb{C}_c - \bigcup_{i=1}^k D(p^i(c), r_i(c)), \\ \phi_c(p^i(c)) &= p_1^i(c) \quad \text{for } i = 1, \dots, k.\end{aligned}$$

Note that if U is a small neighborhood of $0 \in \mathbb{C}^2$ and c is such that $K/2 < |c| < K$, then $\phi_c(p) = p$ if $p \in U \cap \mathbb{C}_c$. So, we can define a homeomorphism $\Phi: (\mathbb{C}^2, 0) \rightarrow (\mathbb{C}^2, 0)$ by

$$\begin{aligned}\Phi(p) &= p \quad \text{if } p \notin \{(x, y); y = cx \text{ and } |c| < K\} \\ \Phi(p) &= \phi_c(p) \quad \text{if } p \in \mathbb{C}_c \text{ and } |c| < K.\end{aligned}$$

In this case Φ is the desired homeomorphism of the proposition.

In the general case, the tangent cone of $V(f)$ is not necessarily reduced to a point; we decompose

$$V(f) = V^1 \cup \dots \cup V^N,$$

where each V^j has a tangent cone reduced to a point a_j . Similarly, we decompose

$$V(f_k) = V_k^1 \cup \dots \cup V_k^N,$$

where the tangent cone of V_k^j is the point a_j . By reducing the neighborhood U of $0 \in \mathbb{C}^2$, we can assume that, for every j , there is a K_j such that $V^j \cap \mathbb{C}_c = \{0\}$ if $|c - a_j| \geq K_j$ and the discs $D(a_j, K_j)$ are mutually disjoint. Now we can define a homeomorphism Φ as follows:

$$\begin{aligned}\Phi(p) &= p \quad \text{if } p \notin \bigcup_{j=1}^N \{(x, y); y = cx \text{ and } |c - a_j| \geq K\} \\ \Phi(p) &= \Phi_j(p) \quad \text{if } p \in \{(x, y); y = cx \text{ and } |c - a_j| < K_j\},\end{aligned}$$

where Φ_j is a homeomorphism, as in the previous case, corresponding to V^j . The proposition is therefore proved. $\square$

3. STATEMENT AND PROOF OF THE MAIN RESULT

Theorem. Let $M(x, y)$ be the germ at $0 \in \mathbb{C}^2$ of a meromorphic function in the class A . Then M is topologically conjugate to a rational function.

The proof of this result needs two lemmas. First, we adopt the following notation: if $V(f)$ is the germ of the variety $\{f = 0\}$, we will denote by $\tilde{V}(f)$ the closure in $\mathbb{C}_0^2$ of $E^{-1}(V(f) - \{0\})$. Also, we will say that two varieties $V(f)$ and $V(g)$ coincide to order k if their defining functions f and g coincide to order k at 0 .

Lemma 3.1. *Let f and g be germs of irreducible holomorphic functions of order j at $0 \in \mathbb{C}^2$. Suppose that f and g coincide to order k ($k > j$) at 0 . Then $\tilde{V}(f)$ and $\tilde{V}(g)$ coincide to order $k - j$ on $E^{-1}(0)$.*

Proof. Without loss of generality, we can assume that the tangent cone of $V(f)$ or $V(g)$ is not the point $(s = 0, y = 0)$. So the varieties $\tilde{V}(f)$ and $\tilde{V}(g)$ admit, respectively, $\tilde{f}(x, t) = f(x, tx)/x^j$ and $\tilde{g}(x, t) = g(x, tx)/x^j$ as defining functions. Therefore, it follows from the fact that f and g coincide to order k at 0 , that $\tilde{f}(x, t)$ and $\tilde{g}(x, t)$ coincide to order $k - j$ on $E^{-1}(0)$.

Lemma 3.2. *Suppose that A, B are germs of holomorphic functions that coincide to order k_1 and $V(f), V(g)$ are germs of analytic varieties that coincide to order k_2 at $0 \in \mathbb{C}^2$. Then the images via the maps $F(x, y) = (x, A(x, y))$ and $G(x, y) = (x, B(x, y))$ of, respectively, $V(f)$ and $V(g)$ coincide to an order $\nu(k_1, k_2)$, where ν increases with k_1, k_2 .*

Proof. Let p be the order of f and g at 0 . Then a parametrization of $V(f)$ (resp. $V(g)$) is (see [4])

$$(x = t^p, y = a(t)) \quad \left(\text{resp. } (x = t^p, y = b(t)) \right),$$

where a, b are holomorphic functions of order $\geq p + 1$. Moreover, it follows from the fact that f and g coincide to order k_2 that a and b coincide to an order $\eta(k_2)$, where η increases with k_2 . The parametrization of $F(V(f))$ (resp. $G(V(g))$) is

$$(x = t^p, y = A(t^p, a(t))) \quad \left(\text{resp. } (x = t^p, y = B(t^p, b(t))) \right).$$

The conclusion follows from the fact that A, B coincide to order k_1 and a, b coincide to order $\eta(k_2)$. $\square$

Proof of the theorem. We write

$$M(x, y) = \frac{\alpha(x, y)}{\beta(x, y)} = \frac{\sum_{j \geq n} P_j(x, y)}{\sum_{j \geq n} Q_j(x, y)},$$

where P_j, Q_j are homogeneous polynomials of degree j and P_n, Q_n coprime. For $k > n$, define the rational function R_k by

$$R_k(x, y) = \frac{\alpha_k(x, y)}{\beta_k(x, y)} = \frac{\sum_{j=n}^k P_j(x, y)}{\sum_{j=n}^k Q_j(x, y)}.$$

Note that the blow-up $\tilde{M}$ of M coincides on $E^{-1}(0)$ with the blow-up $\tilde{R}_k$ of R_k , it is

$$\tilde{M}(0, t) = \tilde{R}_k(0, t) = \frac{P_n(1, t)}{Q_n(1, t)}.$$

Consider the maps $\mathcal{P}: \mathbb{C}_0^2 \mapsto \mathbb{C}_0^2$ and $\mathcal{P}_k: \mathbb{C}_0^2 \mapsto \mathbb{C}_0^2$, defined in the charts (x, t) , (s, y) by

$$\mathcal{P}(x, t) = \left(x, \widetilde{M}(x, t)\right) \quad \text{and} \quad \mathcal{P}(s, y) = \left(\frac{1}{\widetilde{M}(sy, 1/s)}, sy\widetilde{M}\left(sy, \frac{1}{s}\right)\right),$$

$$\mathcal{P}_k(x, t) = \left(x, \widetilde{R}_k(x, t)\right) \quad \text{and} \quad \mathcal{P}_k(s, y) = \left(\frac{1}{\widetilde{R}_k(sy, 1/s)}, sy\widetilde{R}_k\left(sy, \frac{1}{s}\right)\right).$$

Note that the meromorphic map $\mathcal{P}$ (resp. $\mathcal{P}_k$) is holomorphic outside the variety $\{\widetilde{M} = \infty\}$ (resp. $\{\widetilde{R}_k = \infty\}$). Since the jacobian of $\mathcal{P}$ (resp. $\mathcal{P}_k$), in the chart (x, t) , vanishes only on the variety

$$X = \left\{(x, t); \frac{\partial \widetilde{M}}{\partial t} = 0\right\} \quad \left(\text{resp. } X_k = \left\{(x, t); \frac{\partial \widetilde{R}_k}{\partial t} = 0\right\}\right),$$

then $\mathcal{P}$ (resp. $\mathcal{P}_k$) is a covering outside X (resp. X_k) whose number of sheets is the multiplicity of $\widetilde{M}(0, t)$. Since

$$E(X) = \left\{(x, y); \beta \frac{\partial \alpha}{\partial y} - \alpha \frac{\partial \beta}{\partial y} = 0\right\}$$

and

$$E(X_k) = \left\{(x, y); \beta_k \frac{\partial \alpha_k}{\partial y} - \alpha_k \frac{\partial \beta_k}{\partial y} = 0\right\}$$

and the functions $\beta \frac{\partial \alpha}{\partial y} - \alpha \frac{\partial \beta}{\partial y}$, $\beta_k \frac{\partial \alpha_k}{\partial y} - \alpha_k \frac{\partial \beta_k}{\partial y}$ coincide to order $2k - 1$, then X and X_k coincide to an order $\eta(k)$ (see Lemma 3.1). Therefore the varieties $W = \mathcal{P}(X)$ and $W_k = \mathcal{P}_k(X_k)$ coincide to a certain order $\mu(k)$, and μ increases with k (see Lemma 3.2).

Now, it follows from Proposition 2.1 that if k is large enough, then there is a homeomorphism $\Psi: (\mathbb{C}^2, 0) \mapsto (\mathbb{C}^2, 0)$ such that

$$\widetilde{\Psi}(W) = W_k \quad \text{and} \quad \widetilde{\Psi}(\{(x, t); t = t_0\}) = \{(x, t); t = t_0\},$$

for every constant t_0 . Next, since $\mathcal{P}$ (resp. $\mathcal{P}_k$) maps the analytic varieties $\widetilde{M}^{-1}(\widetilde{M}(x_0, t_0))$ (resp. $\widetilde{R}_k^{-1}(\widetilde{R}_k(x_0, t_0))$) into the complex line $t = t_0$, there is a homeomorphism $\widetilde{\Phi}$ such that the diagram

$$\begin{array}{ccc} (\mathbb{C}_0^2, X) & \xrightarrow{\widetilde{\Phi}} & (\mathbb{C}_0^2, X_k) \\ \downarrow \mathcal{P} & & \downarrow \mathcal{P}_k \\ (\mathbb{C}_0^2, W) & \xrightarrow{\widetilde{\Psi}} & (\mathbb{C}_0^2, W_k) \end{array}$$

commutes, i.e., $\widetilde{\Phi}$ conjugates the foliation defined by the level sets of $\widetilde{M}$ to that defined by the level sets of $\widetilde{R}_k$. Since $\widetilde{M} = \widetilde{R}_k$ on $E^{-1}(0)$ and $\widetilde{\Psi}$ is the identity on $E^{-1}(0)$, then $\widetilde{\Phi}$ can be chosen to be the identity on $E^{-1}(0)$. Finally, the homeomorphism $\Phi = E \circ \widetilde{\Phi} \circ E^{-1}$, defined in a neighborhood of $0 \in \mathbb{C}^2$, conjugates R_k with M , i.e., $R_k \circ \Phi = M$. The theorem is proved. $\square$

REFERENCES

1. D. Cerveau and J. F. Mattei, *Formes integrables holomorphes singulieres*, Asterisque **97** (1982).
2. K. Kendig, *Elementary algebraic geometry*, Springer-Verlag, New York, 1977.
3. A. Meziani, *On the integrability of singular differential forms in two complex variables*, Ph.D. Thesis, Rutgers, 1988; Trans. Amer. Math. Soc. (to appear).
4. H. Whitney, *Complex analytic varieties*, Addison Wesley, Reading, MA, 1972.

DEPARTMENT OF MATHEMATICS, UNIVERSITY PARK, FIU, MIAMI, FLORIDA 33199