A RESULT ABOUT THE HILBERT TRANSFORM ALONG CURVES

LINDA V. SAAL

(Communicated by J. Marshall Ash)

ABSTRACT. Let G be a connected and simply connected, nilpotent Lie group and let $\gamma: (-1, 1) \rightarrow G$ be a (connected) analytic curve such that $\gamma(0) = 0$. Then the Hilbert transform along γ ,

$$Tf(x) = p.v. \int_{0 < |t| < 1} f(x\gamma(t)^{-1}) dt/t,$$

is bounded on $L^p(G)$, $1 < p < \infty$.

INTRODUCTION

Let G be a simply connected, nilpotent Lie group. From [Ch] we know that if γ is a homogeneous curve with respect to a group of dilations of G , $\gamma(0) = 0$, then the Hilbert transform along γ is a bounded operator on $L^p(G)$, for $1 < p < \infty$; moreover $Tf(x)$ exists a.e. for all $f \in L^p(G)$. In [RS2] singular integral operators whose kernels are homogeneous distributions of critical degree with mean value zero and supported on a homogeneous, analytic submanifold of G are considered.

The purpose of this note is to prove that if $\gamma: (-1, 1) \rightarrow G$ is a connected, analytic curve such that $\gamma(0) = 0$ then the

$$Tf(x) = p.v. \int_{0 < |t| < 1} f(x\gamma(t)^{-1}) dt/t$$

is a bounded operator on $L^p(G)$, $1 < p < \infty$.

For the proof we first assume that a certain group of dilations associated to γ are automorphisms of G and also that γ generates G . We use the iteration argument of [Ch]. The general case follows from a transference theorem [RS2].

Finally the boundedness of the maximal operator

$$T^*f(x) = \sup_{r>0} \left| \int_{r<|t|<1} f(x\gamma(t)^{-1}) dt/t \right|$$

on $L^p(G)$, implies that $Tf(x)$ exists a.e. for all $f \in L^p$.

Received by the editors January 10, 1989 and, in revised form, July 1, 1989.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 42B20.

This research was supported by the Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina.

The proof of the theorem. Let G be a connected and simply connected, nilpotent Lie group. Let $\gamma: (-1, 1) \rightarrow G$ be a (connected) analytic curve that generates G in the sense that γ is not contained in any proper closed subgroup of G . Also assume that $\gamma(0)$ is the identity of G . Denote by $\exp: \mathcal{G} \rightarrow G$ the exponential map. First of all we remark the following facts:

(i) γ generates G if and only if $\{d^j \bar{\gamma}/dt^j|_{t=0}\}_{j \in \mathbb{N}}$ generate $\mathcal{G}$ as a Lie algebra, with $\bar{\gamma} = \gamma \circ \exp^{-1}$. Indeed, let $\mathcal{L}$ be the Lie subalgebra of $\mathcal{G}$ generated by $\{d^j \bar{\gamma}/dt^j|_{t=0}\}$ and $H = \exp \mathcal{L}$. Since $\bar{\gamma}$ is analytic, $\bar{\gamma} \subset \mathcal{L}$.

(ii) We can choose a coordinate system on G such that

$$\gamma(t) = \left(\frac{t^{a_1}}{a_1!} \varphi_1(t), \dots, \frac{t^{a_k}}{a_k!} \varphi_k(t), 0, \dots, 0 \right)$$

with $1 \leq a_1 < a_2 < \dots < a_k$, φ_i are analytic functions and $\varphi_i(0) = 1$. Indeed, following [SW], we choose $a_1 = \inf\{j | d^j \bar{\gamma}/dt^j|_{t=0} \neq 0\}$. Given $a_1, \dots, a_i$, define

$$a_{i+1} = \inf \left\{ j | j > a_i \text{ and } \left\{ \frac{d^{a_1} \bar{\gamma}}{dt^{a_1}} \Big|_{t=0}, \dots, \frac{d^{a_i} \bar{\gamma}}{dt^{a_i}} \Big|_{t=0}, \frac{d^j \bar{\gamma}}{dt^j} \Big|_{t=0} \right\} \right. \\ \left. \text{are linearly independent} \right\}.$$

Define $e_i = d^{a_i} \bar{\gamma}/dt^{a_i}|_{t=0}$. We thus obtain a set $\{e_1, \dots, e_k\}$, $k \leq n$, maximal with respect to be L.I. such that

$$\bar{\gamma}(t) = \frac{t^{a_1}}{a_1!} \varphi_1(t) e_1 + \dots + \frac{t^{a_k}}{a_k!} \varphi_k(t) e_k.$$

We extend $\{e_1, \dots, e_k\}$ to a basis of $\mathcal{G}$, $\{e_1, \dots, e_k, e_{k+1}, \dots, e_n\}$. If A denotes the change of basis matrix, the coordinates of $\gamma(t)$ with respect to $\psi = \exp \circ A$ are $((t^{a_1}/a_1!) \varphi_1(t), \dots, (t^{a_k}/a_k!) \varphi_k(t), 0, \dots, 0)$. From now on we fix this coordinate system on G .

(iii) We now associate to γ a group of dilations D_r , $r > 0$, by

$$D_r(x_1, \dots, x_n) = (r^{a_1} x_1, \dots, r^{a_k} x_k, r^{a_{k+1}} x_{k+1}, \dots, r^{a_n} x_n),$$

where $a_1, \dots, a_k$ are as in (ii) and $a_{k+1}, \dots, a_n$ are arbitrary integers. Let $Q = a_1 + \dots + a_n$.

(iv) Let γ_0 be the curve in G with coordinates

$$(t^{a_1}/a_1!, \dots, t^{a_k}/a_k!, 0, \dots, 0).$$

Then γ_0 is analytic, homogeneous with respect to D_r and generates G . Indeed, if $\bar{\gamma} = \psi^{-1} \circ \gamma$ and $\bar{\gamma}_0 = \psi^{-1} \circ \gamma$, then $d^{a_i} \gamma_0/dt^{a_i}|_{t=0} = d^{a_i} \gamma/dt^{a_i}|_{t=0} = e_i$, $1 \leq i \leq k$.

We have associated to γ a group of dilations $\{D_r\}_{r>0}$ and a curve γ_0 that is homogeneous with respect to D_r and also generates G . For the moment we also assume that $\{D_r\}$ are automorphisms of G and fix a homogeneous norm

with respect to D_r , e.g., $|x| = \max_{1 \leq i \leq n} \{|x_i|^{1/a_i}\}$. Denote by $\bar{B}$ the closure of the unit ball in $\mathbf{R}^n$.

Lemma 1. Let ϕ, ϕ_0 be the functions on $\bar{B}$ defined by $\phi(t_1, \dots, t_n) = \gamma(t_1) \dots \gamma(t_n)$; $\phi_0(t_1, \dots, t_n) = \gamma_0(t_1) \dots \gamma_0(t_n)$. For each $j \in \mathbf{N}$, $j \geq 0$, define ϕ_j on $\{t \in \mathbf{R}^n | \frac{1}{2} \leq |t_i| \leq 1\}$ by $\phi_j(t) = D_{2^j} \phi(2^{-j}t)$. If $J_j(t) = \det d\phi_j(t)$ and $J_0(t) = \det d\phi_0(t)$, then $J_j(t) \xrightarrow{j \rightarrow +\infty} J_0(t)$ uniformly on $\{t \in \mathbf{R}^n | \frac{1}{2} \leq |t_i| \leq 1\}$.

Proof. Let $\bar{\phi}_j(t) = \exp^{-1} \phi_j(t)$, $\bar{\phi}_0(t) = \exp^{-1} \phi_0(t)$.

$$\begin{aligned} \bar{\phi}_j(t) &= \exp^{-1}(D_{2^j} \phi(2^{-j}t)) = D_{2^j} \exp^{-1} \phi(2^{-j}t) \\ &= D_{2^j} \exp^{-1}(\gamma(2^{-j}t_1) \dots \gamma(2^{-j}t_n)). \end{aligned}$$

Then

$$(*) \quad \bar{\phi}_j(t) = D_{2^j}(\bar{\gamma}(2^{-j}t_1) + \dots + \bar{\gamma}(2^{-j}t_n)) + D_{2^j}(P_1(\dots), \dots, P_n(\dots))$$

where $P_j(\dots)$ denotes $P_j(\bar{\gamma}(2^{-j}t_1), \dots, \bar{\gamma}(2^{-j}t_n))$ and P_j are polynomial functions on the coordinates of $\gamma(t_1), \dots, \gamma(t_n)$.

If γ is homogeneous, $\bar{\phi}_j(t) = \bar{\phi}(t)$ and the first term of the right side of (*) is also independent of j . Thus either $P_l \equiv 0$ or $P_l(t_1, \dots, t_n)$ is homogeneous of degree a_l , $l = 1, \dots, n$,

$$\begin{aligned} P_l(t_1, \dots, t_n) &= \sum \beta_{j_1, \dots, j_k}^{i_1, \dots, i_k} \varphi_{j_1}(t_{i_1}) \dots \varphi_{j_k}(t_{i_k}) t_{i_1}^{a_{j_1}} \dots t_{i_k}^{a_{j_k}} \\ &= \sum \beta_{j_1, \dots, j_k}^{i_1, \dots, i_k} F_{j_1, \dots, j_k}(t_{i_1}, \dots, t_{i_k}) t_{i_1}^{a_{j_1}} \dots t_{i_k}^{a_{j_k}}, \end{aligned}$$

with $a_{j_1} + \dots + a_{j_k} = a_l$, $\beta_{j_1, \dots, j_k}^{i_1, \dots, i_k} \in \mathbf{C}$ and $F_{j_1, \dots, j_k}(0) = 1$. Then

$$\begin{aligned} \bar{\phi}_j(t) &= \left(\dots, \sum_{i=1}^n \varphi_i(2^{-j}t_i) t_i^{a_i}, \dots \right) \\ &\quad + \left(\dots, \sum \beta^{i_s}_{j_1, \dots, j_k} F_{j_1, \dots, j_k}(2^{-j}t_{i_1}, \dots, 2^{-j}t_{i_k}) t_{i_1}^{a_{j_1}} \dots t_{i_k}^{a_{j_k}}, \dots \right). \end{aligned}$$

By the mean value theorem applied to $\varphi_i^{i_s}$ and $F_{j_1, \dots, j_k}^{i_s}$, we have that

$$\begin{aligned} \bar{\phi}_j(t) &= \left(\dots, \sum_{i=1}^n t_i^{a_i} + \sum \beta^{i_s}_{j_1, \dots, j_k} t_{i_1}^{a_{j_1}} \dots t_{i_k}^{a_{j_k}}, \dots \right) \\ &\quad + 2^{-j}(G_1(t_1, \dots, t_n), \dots, G_n(t_1, \dots, t_n)) \\ &= \bar{\phi}_0(t) + 2^{-j}(G_1(t_1, \dots, t_n), \dots, G_n(t_1, \dots, t_n)), \end{aligned}$$

with G_k bounded on $\{t | \frac{1}{2} \leq |t_i| \leq 1\}$, $k = 1, \dots, n$. This proves the lemma since $\det d_x \exp = 1$.

Remark. (v) If γ is an analytic curve that generates G then $\phi: \bar{B} \rightarrow G$ given by $\phi(t_1, \dots, t_n) = \gamma(t_1) \dots \gamma(t_n)$ has rank n almost everywhere. Indeed, since γ_0 is a homogeneous curve that generates G , $J_0 \not\equiv 0$ [Ch]. Since γ_0 is analytic,

$J_0(t) \neq 0$ almost everywhere. Let $J(t) = \det \phi(t)$. Since $\phi_j(t) = D_{2^j} \phi(2^{-j}t)$, $2^{j(Q-n)} J(2^{-j}t) = J_j(t) \xrightarrow{j \rightarrow \infty} J_0(t)$ on $\{t | \frac{1}{2} \leq |t_i| \leq 1\}$. Then $J \neq 0$. (Furthermore $J(t) \neq 0$ almost everywhere).

By a result of [RS2] we know that if $\phi: \bar{B} \rightarrow \mathbf{R}^n$ is an analytic function having rank n at almost everywhere point and ψ is a C^1 -function supported on $\bar{B}$ then the transported measure by ϕ of $\psi(t) dt$ is absolutely continuous and its density ρ satisfies and L^1 -Hölder condition $\int |\rho(x+y) - \rho(x)| dx \leq c \|y\|^\delta$, for some $\delta > 0$ depending on ϕ , $c > 0$ that depends also on ψ .

In the next lemma we say a little more about c .

Lemma 2. Let $J(t) = \det d\phi(t)$. Then

$$\int |\rho(x+y) - \rho(x)| dx \leq \tilde{C} \|y\|^\delta \left(\int_{\bar{B}} |\nabla \psi| + |\psi| \right)^\delta \left(\int_{\bar{B}} |\psi| / 2\delta / |J|(1-\delta) \right)^{1-\delta},$$

for some $\delta > 0$, $\tilde{C} > 0$, depending only on ϕ .

Proof. Following the proof of [RS2] we denote by Z the set of zeros of $J(t)$ and consider a covering $\{B_j(z_j, r_j)\}_j$ of $\bar{B} - Z$, satisfying,

(1) $r_j = c_0 |J(z_j)|$;

(2) the balls $B_j^* = B(z_j, 2r_j)$ are disjoint from Z and have the bounded overlapping property; i.e. there is an integer N such that no point belongs to more than N of the B_j^* .

Let $\{n_j\}$ be a smooth partition of unity on $\bar{B} - Z$ subordinated to $\{B_j\}$ and such that $\|\nabla n_j\|_\infty \leq \tilde{C} r_j^{-1}$. Let $\psi_j = \psi n_j$ and let ρ_j be the transported measure by ϕ of ψ_j . Since $\phi|_{B_j}$ is invertible [RS2] $\rho_j(x) = |J\phi^{-1}(x)|^{-1} \psi_j(\phi^{-1}(x))$ and thus

$$\int |\rho_j(x)| dx = \|\psi_j\|_1 = \int |\psi_j|$$

and

$$\begin{aligned} & \int |\nabla \rho_j(x)| dx \\ & \leq \tilde{C} \left(\int |\nabla \psi(t)| n_j(t) |J(t)|^{-1} dt + \int |\psi| |\nabla n_j| |J(t)|^{-1} dt + \int |J(t)|^{-2} |\psi| dt \right) \\ & \leq \tilde{C} r_j^{-2} \left(\int_{B_j} |\nabla \psi| + \int_{B_j} |\psi| \right). \end{aligned}$$

This implies that if $0 < \delta \leq 1$,

$$\int |\rho_j(x+y) - \rho_j(x)| dx \leq \tilde{C} \|y\|^\delta r_j^{-2\delta} \left(\int_{B_j} |\nabla \psi| + |\psi| \right)^\delta \left(\int_{B_j} |\psi| \right)^{1-\delta}.$$

Now by Hölder's inequality

$$\begin{aligned} & \sum_j \left(\int_{B_j} |\nabla \psi| + |\psi| \right)^\delta r_j^{-2\delta} \left(\int_{B_j} |\psi| \right)^{1-\delta} \\ & \leq \left(\sum_j \left(\int_{B_j} |\nabla \psi| + |\psi| \right)^\delta \right) \left(\sum_j r_j^{-2\delta/(1-\delta)} \int_{B_j} |\psi| \right)^{1-\delta}. \end{aligned}$$

The lemma follows from the fact that $J|_{B_j^*}$ is comparable with r_j and from the bounded overlapping property.

Theorem. Let γ be an analytic curve that generates G and such that the group of dilations associated to γ are automorphisms of G . Then

$$Tf(x) = p.v. \int_{0 < |t| < 1} f(x\gamma(t)^{-1}) dt/t$$

defines a bounded operator on $L^p(G)$, $1 < p < \infty$.

Proof. (We follow [RS2] for the proof.) Denoted by μ the distribution given by $\mu(f) = p.v. \int_{0 < |t| \leq 1} f(\gamma(t)) dt/t$. Let $\tilde{\varphi}$ be in $\mathcal{C}^\infty(\frac{1}{2}, 2)$ such that $\sum_{j=0} \tilde{\varphi}(2^j |t|) = 1$. Let $\varphi(x) = \varphi(|x|)$ where $|\cdot|$ denote the homogeneous norm defined in Remark (iv). Let $\mu_j = \tilde{\varphi}(2^j |t|) \mu$.

Let $\varphi_j(x) = 2^{jQ} \varphi(D_{2^j} x)$ and $\eta_j = \frac{1}{c}(\varphi_{j+1} - \varphi_j)$ with $c = \int \varphi$.

Since $\{c^{-1} \varphi_j\}$ is an approximate identity in G we can write

$$\delta = c^{-1} \lim_{j \rightarrow +\infty} \varphi_j = \sum_{j \geq j_0} \eta_j + c^{-1} \varphi_{j_0}$$

for each j_0 fixed. Also $\eta = \sum_{j \geq 0} \mu_j$. Then

$$\mu = \delta * \sum_j \mu_j = \sum_j \left(\sum_{k \geq j_0} \eta_k * \mu_j \right) + c^{-1} \sum_j \varphi_{j_0} * \mu_j.$$

For each j fixed, take $j_0 = j + 1$. Then

$$\begin{aligned} \mu &= \sum_{k \geq 1} \left(\sum_{j \geq 0} \eta_{k+j} * \mu_j \right) + c^{-1} \sum_j (\varphi_{j+1} * \mu_j) \\ &= \sum_{k \geq 1} M_k + N. \end{aligned}$$

If we prove

(1) For $0 < \epsilon \leq 1$, $\|M_k\|_{p,p} \leq C_{\epsilon,p} 2^{\epsilon k}$, $\|N\|_{p,p} \leq C$.

(2) $\exists \sigma > 0$ such that $\|M_k\|_{2,2} \leq C 2^{-\sigma k}$.

Then the theorem follows by interpolation and duality. Indeed, fix $1 < p < 2$ and take $1 < p_0 < p$. Then $\frac{1}{p} = \frac{1}{2}s + (1-s)\frac{1}{p_0}$ for some $0 < s < 1$. Thus

$\|M_k\|_{p,p} \leq C_\epsilon 2^{-\sigma k s} 2^{\epsilon k(1-s)}$ for $0 < \epsilon \leq 1$ and some $\sigma > 0$. Choosing ϵ small enough we obtain $\sum_k \|M_k\|_{p,p} < +\infty$.

(1) We observe that $M_k = \sum_{j \geq 0} \eta_{k+j} * \mu_j = \sum_{j \geq 0} f_j^k$ where f_j^k has the following properties

- (a) $f_j^k = 0$,
- (b) $\int |f_j^k(xy) - f_j^k(x)| dx \leq C 2^{(k+j)\epsilon} |y|^\epsilon$,
- (c) $\text{supp } f_j^k \subset \{x | |x| \leq C 2^{-j}\}$,
- (d) $\int |f_j^k| \leq C$.

Then Cotlar's lemma implies that $\|M_k\|_{2,2} \leq C 2^{\epsilon k}$ and the weak type (1.1) of M_k follows by checking that $\int_{|x| > 2|y|} |M_k(xy) - M_k(x)| dx \leq C 2^{k\epsilon}$. The same argument holds for N . For a proof see [ChNSW].

(2) By Cotlar's lemma it is enough to prove that for some $\sigma > 0$ $\|f_j^k * f_l^k\|_{2,2} \leq C 2^{-\sigma k} 2^{-|j-l|\sigma}$. We check this for $j > l$, by using the iteration argument of [Ch].

$$\begin{aligned} \|f_j^k * f_l^k\|_{2,2} &\leq C \|\eta_{k+j} * \mu_j * \mu_l^*\|_{2,2} \\ &\leq C \|\eta_{k+j} * \mu_j\|^{1/2} \|\mu_l^* * \mu_l * \mu_j^* * \eta_{k+j}\|^{1/2} \\ &\leq \dots \leq C \|\eta_{k+j} * \mu_j\|^{1-2^{-r}} \|(\mu_l^* * \mu_l)^{2^{r-1}} * \mu_j^* * \eta_{k+j}\|^{2^{-r}}. \end{aligned}$$

Let $\psi_l(t_1, \dots, t_n) = \prod_{i=1}^n \frac{1}{t_i} \tilde{\phi}(2^l t_i)$ and $\phi(t_1, \dots, t_n) = \gamma(t_1) \cdots \gamma(t_n)$ as in Lemma 1.

Since $\mu_l * \dots * \mu_l$ is the transported measure by ϕ of $\psi_l(t) dt$ and since ϕ is analytic with rank n almost everywhere, $\mu_l * \dots * \mu_l$ is absolutely continuous [RS2].

If ρ_l denotes its density, we have to prove that

$$\|\rho_l * \mu_j^* * \eta_{k+j}\|_1 \leq C 2^{-\sigma k} 2^{-|j-l|\sigma}$$

for some $\sigma > 0$. (Recall that we have assumed $l < j$.)

Let $\tilde{\rho}_l(x) = 2^{-lQ} \rho_l(D_2^{-l} x)$. Then $\tilde{\rho}_l$ is the transported measure by $\phi_l(t) = D_2^l \phi(2^{-l} t)$ of $\tilde{\psi}(t) = 2^{-ln} \psi_l(2^{-l} t)$ ($\text{supp } \tilde{\psi} \subset \{t | \frac{1}{2} \leq |t| \leq 2\}$).

If we prove that

$$(A) \quad \int_G |\tilde{\rho}_l(xy) - \tilde{\rho}_l(x)| dx \leq C |y|^\epsilon, \quad \text{for some } C, \epsilon > 0, \text{ independent of } l$$

then $\int_G |\rho_l(xy) - \rho_l(x)| dx \leq C 2^{l\epsilon} |y|^\epsilon$. From this and the fact that η_{k+j} has mean value zero and $\text{supp } \eta_{k+j} \subset \{x | |x| \leq C 2^{-(k+j)}\}$ we obtain the inequality desired.

But if $\bar{\rho}_l = \tilde{\rho}_l \circ \exp$ and $\bar{\phi}_l = \exp^{-1} \circ \phi_l$ then $\bar{\rho}_l$ is the transported measure of $\tilde{\psi}(t) dt$ by $\bar{\phi}_l$ and (A) is equivalent to

$$(B) \quad \int_{\mathbf{R}^n} |\bar{\rho}_l(x+y) - \bar{\rho}_l(x)| dx \leq C \|y\|^\delta, \quad C, \delta > 0, \text{ independent of } l \text{ (see [RS2])}.$$

Finally we prove (B). Since by Lemma 2,

$$\int_{\mathbb{R}^n} |\bar{\rho}_l(x+y) - \bar{\rho}_l(x)| dx \leq C \|y\|^\delta \left(\int |\nabla \tilde{\psi}| + |\tilde{\psi}| \right)^\delta \left(\int |\tilde{\psi}|/|J_l| 2\delta/1 - \delta \right)^{1-\delta}$$

for some $0 < \delta < 1$, we only have to check that $\int_{\sup \tilde{\psi}} |\bar{J}_l^{(t)}|^{-\alpha} dt \leq C$, with C independent of l , for some $0 < \alpha < 1$. By Lemma 1, $2^{l(Q-n)} J(2^{-l}t) = \bar{J}_l(t)$ converges uniformly to $\bar{J}_0(t)$, which is a homogeneous polynomial of degree $Q - n$. Thus if we develop $\bar{J}(t)$ about 0 in the Taylor expansion, $\bar{J}(t) = P(t) + R(t)$, where P is a homogeneous polynomial of degree $Q - n$ and $\frac{R(t)}{|t|^{Q-n}} \rightarrow_{|t| \rightarrow 0} 0$. By a linear change of variables we can normalize P in the t_n -direction, i.e., $P(t) = t_n^{Q-n} + \sum_{j=0}^{Q-n-1} b_j(t_1, \dots, t_{n-1}) t_n^j$ with $b_j(0) = 0$. The Weierstrass preparation theorem now implies that

$$\bar{J}(t) = (t_n^{Q-n} + a_1(t') t_n^{Q-n-1} + \dots + a_{Q-n}(t')) h(t), \quad t' = (t_1, \dots, t_{n-1}),$$

where $a_1, \dots, a_{Q-n}$ and h are analytic functions in a neighborhood of 0, and $h(0) \neq 0$. (See [H].) Then

$$\begin{aligned} \int_{\sup \tilde{\psi}} |\bar{J}_l(t)|^{-\alpha} dt &= 2^{-l(Q-n)\alpha} \int_{\sup \tilde{\psi}} |\bar{J}(2^{-l}t)|^{-\alpha} dt \\ &\leq C 2^{-l(Q-n)\alpha} \int_{\frac{1}{2} \leq |t| \leq 2} 2^{l(n-Q)\alpha} |t_n^{Q-n} + 2^l a_1(t') t_n^{Q-n-1} \\ &\quad + \dots + 2^{l(Q-n)} a_{Q-n}(t')|^{-\alpha} dt_n dt' \\ &\leq C A_\alpha \int_{1 \leq |t'| \leq 2} (1 + 2^l |a_1(t')| + \dots + 2^{l(Q-n)} |a_{Q-n}(t')|)^{-\alpha} dt' \end{aligned}$$

which is bounded independent of l . The last inequality follows from [RS1] and the proof of the theorem is complete.

The general case where γ is any connected analytic curve in G , follows by transference [CW, RS2] as is shown in the following remarks.

Remark. (vi) Assume now that γ generates a proper subgroup of G . As in Remark (ii) we choose $\{e_1, \dots, e_l\}$ that generate a Lie subalgebra $\mathcal{L}$ of $\mathcal{G}$. Extend it to a basis $\{e_1, \dots, e_l, \dots, e_n\}$ of $\mathcal{G}$ and define for $r > 0$, $D_r(x_1, \dots, x_l, \dots, x_n) = (r^{a_1} x_1, \dots, r^{a_l} x_l, \dots, r^{a_n} x_n)$ where $a_1, \dots, a_l$ are determined by γ and $a_{l+1}, \dots, a_n$ are arbitrary integers. Assume that D_r are automorphisms of G , let γ_0 be the homogeneous curve (w.r.t. D_n) associated to γ , and let H be the subgroup of G generated by γ_0 .

Then H is homogeneous with the dilations induced and γ generates H . By transference, $Tf = f * \mu$ is bounded on $L^p(G)$.

Remark. (vii) Let γ be an analytic curve that generates a connected and simply connected, nilpotent Lie Group N . Let D_r be a group of dilation associated to

γ , which are not necessarily automorphisms of N and let γ_0 be as in Remark (ii). As in [RS2] let $\mathcal{g}$ be the step m free Lie algebra generated by $\mathfrak{n}$, for m large enough and let G be the simply connected group with $\mathcal{L}(G) = \mathcal{g}$. Then there exists H a normal subgroup of G such that G/H is isomorphic to N . Denote by $\sigma: \mathcal{g} \rightarrow \mathfrak{n}$ the quotient morphism. By construction of $\mathcal{g}$ the dilations D_r on $\mathfrak{n}$ extend to automorphisms $\tilde{D}_r$ of $\mathcal{g}$.

Let $\bar{\gamma} = \exp_N^{-1} \circ \gamma$, $\tilde{\gamma} = \exp_G \circ \bar{\gamma}$ and $\bar{\gamma}_0 = \exp_N^{-1} \circ \gamma_0$, $\tilde{\gamma}_0 = \exp_G \bar{\gamma}_0$. Then $\tilde{\gamma}_0$ is a homogeneous curve w.r.t. $\tilde{D}_r$ and the coordinates of $\tilde{\gamma}_0$ (resp. $\tilde{\gamma}$) with respect to $\exp_G$ are those of γ_0 (resp. γ). Thus $Tf = f * \mu$ is bounded on $L^p(G)$, $1 < p < \infty$ and by transference on $L^p(N)$, $1 < p < \infty$.

Remark. (viii) The maximal function along γ ,

$$M_\gamma f(x) = \sup_{r \leq 1} \frac{1}{r} \left| \int_{0 \leq |t| \leq r} |f(x\gamma(t)^{-1})| dt \right|$$

is bounded on $L^p(G)$, $1 < p \leq \infty$. Also

$$T^* f(x) = \sup_r \left| \int_{r \leq |t| \leq 1} f(x\gamma(t)^{-1}) dt/t \right|$$

is bounded on $L^p(G)$, $1 < p < \infty$ and hence $Tf(x)$ exists a.e. for all $f \in L^p(G)$, $1 < p < \infty$.

To see this we first remark that if $Hf = \sup_j |f * f_j|$ with f_j satisfying

- (i) $\|f_j\|_1 \leq C_1$,
- (ii) $\int |f_j(x\gamma) - f_j(x)| \leq C_1 2^{j\epsilon} |y|^\epsilon \forall \epsilon > 0$,
- (iii) $\text{supp } f_j \subset \{|x| \leq C 2^{-j}\}$.

Then H is bounded on L^∞ and of weak type (1.1) with constant $C_1 C_\epsilon$. Indeed, we can assume that $f \geq 0$. For $\lambda > 0$ we decompose $f = g + b$ where $\|g\|_\infty \leq \lambda$, $b = \sum_j b_j$, $\text{supp } b_j \subset B_j = B(a_j, r_j)$, $\int b_j = 0$, $\int |b_j| \leq A\lambda |B_j^*|$, and $\sum |B_j^*| \leq \frac{A}{\lambda} \|f\|_1$, $B_j^* = B(a_j, 2r_j)$. Then it is enough to estimate $|\{x | Hb(x) > \lambda\}|$. But

$$\{x | Hb(x) > \lambda\} \subset \left(\bigcup_j B_j^* \right) \cup \left\{ x \in {}^c \left(\bigcup_j B_j^* \right) \mid Hb(x) > \lambda \right\}$$

and

$$\begin{aligned}
& \left| \left\{ x \in {}^c \left(\bigcup B_j^* \right) \mid Hb(x) > \lambda \right\} \right| \\
& \leq \frac{1}{\lambda} \int_{{}^c(\cup B_j^*)} Hb(x) \leq \frac{1}{\lambda} \sum_i \int_{{}^c B_i^*} Hb_i(x) \\
& = \frac{1}{\lambda} \sum_i \int_{{}^c B_i^*} \sup_j \left| \int_{B_i} b_i(y) f_j(y^{-1}x) dy \right| dx \\
& = \frac{1}{\lambda} \sum_i \int_{{}^c B_i^*} \sup_j \left| \int_{B_i} b_i(y) (f_j(y^{-1}x) - f_j(a_i^{-1}x)) dy \right| dx \\
& \leq \frac{1}{\lambda} \sum_i \int_{B_i} |b_i(y)| \sum_j \int_{{}^c B_i^*} |f_j(y^{-1}x) - f_j(a_i^{-1}x)| dx dy \\
& \leq \frac{1}{\lambda} \sum_i \int_{B_i} |b_i(y)| \sum_j \int_{|z| > 2|y|} |f_j(\tilde{y}^{-1}z) - f_j(z)| dz \\
& \leq C_\epsilon \cdot \frac{1}{\lambda} \sum_i \int_{B_i} |b_i(y)| dy \quad \text{by (ii) and (iii), } \tilde{y} = y^{-1}a_i
\end{aligned}$$

Now

$$M_\gamma f(x) \leq C \sup(f * |\mu_j|)(x)$$

and

$$\sup_j (f * |\mu_j|) \leq \sum_{k \geq 1} \sup_j |f * \eta_{k+j} * |\mu_j|| + \sup_j (f * \varphi_j * |\mu_j|) = \sum_k \widetilde{M}_k + \widetilde{N}.$$

By the above $\widetilde{M}_k$ is bounded on L^∞ and for each $\epsilon > 0$ there exists a constant C_ϵ such that $\widetilde{M}_k$ is of weak type (1.1) with constant $C_\epsilon 2^{k\epsilon}$. The same holds for $\widetilde{N}$. We also have that $\|\widetilde{M}_k\|_{2,2} \leq C 2^{-\sigma k}$ for some $\sigma > 0$, by using the technique of the square functions (see [Ch]). Indeed we have to check that the operators $S_k f(x) = \sum_j a_j f * \eta_{k+j} * |\mu_j|$, $a_j = \pm 1 \forall j$, have L^2 boundedness $C 2^{-\sigma k}$ but this follows as in the proof of the theorem since n -times $|\mu_j| * \dots * |\mu_j|$ is absolutely continuous.

The boundedness of T^* on L^p , $1 < p < \infty$, reduces to prove that the operators

$$T_1 f(x) = \sup_{i \geq 0} \left| f * \sum_{j \geq i} \mu_j * \varphi_i \right|$$

and

$$T_2 f(x) = \sup_i \left| \sum_{j=0}^i f * \mu_j * (\delta - \varphi_i) \right|$$

are bounded on L^p for $1 < p < \infty$ (see [Ch]). $T_1 f(x)$ is bounded pointwise by the maximal operator on G since the functions $\psi_i = \sum_{j > i} \mu_j * \varphi_i$ satisfy $\sup \psi_i \subset \{x \mid |x| \leq C 2^{-i}\}$ and $|\psi_i(x)| \leq C 2^{iQ}$.

Now $T_2 f(x) \leq \sum_k \sup_j |f * \mu_j * (\delta - \varphi_{k+j})| = \sum_k \widetilde{M}_k$ and $\widetilde{M}_k f \leq M_\gamma f + \sup_j |f * \mu_j * \varphi_{k+j}|$. Since $\sup_j |f * \mu_j * \varphi_{k+j}|$ is bounded on L^∞ and of weak type (1.1), it is bounded on L^p for $1 < p < \infty$ and so is $\widetilde{M}_k$. To see that $\|\widetilde{M}_k\|_{2,2} \leq C2^{-\sigma k}$, for some C , $\sigma > 0$, we can argue as in the theorem since n -times $\mu_j * \dots * \mu_j$ is absolutely continuous with density ρ_j satisfying the equation $\int |\rho_j(xy) - \rho_j(x)| \leq C2^{j\epsilon} |y|^\epsilon$ and $(\delta - \varphi_{k+j})$ is a measure with bounded L^1 -norm, $\text{supp}(\delta - \varphi_{k+j}) \subset \{x | |x| \leq C2^{-(k+j)}\}$ and it has mean value zero.

ACKNOWLEDGMENTS

I am deeply indebted to Fulvio Ricci with whom I had the privilege to work at the Politecnico di Torino. He suggested this problem and I wish to express my gratitude for his generous help during its preparation.

Added in proof. I am very grateful to the referee for informing me of a recent preprint by M. Christ, A. Nagel, E. Stein, and S. Wainger [ChNSW] where more general results are presented, and from which our main theorem follows.

REFERENCES

- [Ch] M. Christ, *Hilbert transform along curves*, I, Ann. of Math. **122** (1985).
- [ChNSW] M. Christ, A. Nagel, E. Stein, and S. Wainger, *Singular and maximal Radon transforms*, preprint.
- [CW] R. Coifman and G. Weiss, *Analyse harmonique non-commutative sur certain espaces homogenes*, Lecture Notes in Math., vol 242, Springer-Verlag, Berlin and New York, 1971.
- [H] L. Hormander, *An introduction to complex analysis in several variables*, 2nd ed. North-Holland Mathematical Library, vol. 7, 1973.
- [RS1] F. Ricci and E. Stein, *Harmonic analysis on nilpotent groups and singular integrals*, Internat. J. Funct. Anal. **73** (1987).
- [RS2] —, *Harmonic analysis on nilpotent groups and singular integrals*, II. J. Funct. Anal. **77** (1988).
- [SW] E. Stein and S. Wainger, *Problems in harmonic analysis related to curvature*, Bull. Amer. Math. Soc. **84** (1978).

FA. M.A.F., VALPARAÍSO Y ROGELIO MARTÍNEZ, CIUDAD UNIVERSITARIA 5000, CORDOBA, ARGENTINA