SIERPINSKI SETS AND STRONG FIRST CATEGORY

JAKUB JASINSKI AND TOMASZ WEISS

(Communicated by Andreas R. Blass)

ABSTRACT. We prove that if S is a Sierpinski set and $N \subseteq \mathbb{R}$ is an F_σ set of measure zero, then $(N + t) \cap S = \emptyset$ for some $t \in \mathbb{R}$. A similar result holds for generalized Sierpinski sets under Martin's Axiom.

INTRODUCTION

An uncountable set $S \subseteq \mathbb{R}$ is called a Sierpinski set if $|N \cap S| \leq \omega$ whenever $N \subseteq \mathbb{R}$ is of measure zero. We say that $X \subseteq \mathbb{R}$ has strong first category (SFC), if, for any measure-zero set $N \subseteq \mathbb{R}$, there exists $t \in \mathbb{R}$ such that $(N + t) \cap X = \emptyset$. It is well known that Lusin sets, the category analog of the Sierpinski sets, have strong measure zero (see [M] for details). This leads to the question posed by F. Galvin [M, p. 210]: Do all Sierpinski sets have SFC? It is known that under CH there are Sierpinski sets which have SFC. Simply let $\{N_\alpha\}_{\alpha < \omega_1}$ be all G_δ measure zero sets and inductively pick

$$s_\alpha \in \mathbb{R} \setminus \bigcup_{\beta < \alpha} ((F_\beta \cup \{s_\beta\}) \cup (F_\beta + t_\beta)),$$

where t_α is such that $\{s_\beta : \beta \subseteq \alpha\} \cap (F_\alpha + t_\alpha) = \emptyset$. Then $S = \{s_\alpha : \alpha < \omega_1\}$ is an SFC Sierpinski set. Recently Bartoszynski and Ihoda proved that under CH every Sierpinski set is a union of two SFC sets (although it is not known if the union of two SFC sets always has SFC) and that $\text{Con}(\text{ZF}) \Rightarrow \text{Con}(\text{ZFC} + \text{there are Sierpinski sets} + \text{all Sierpinski sets have SFC})$ [BI].

MAIN RESULT

Let μ be Lebesgue measure on $\mathbb{R}$ and let $L_0 = \{N \subseteq \mathbb{R} : \mu(N) = 0\}$. For $X, Y \subseteq \mathbb{R}$ $X \pm Y = \{x \pm y : x \in X \text{ and } y \in Y\}$.

The following is a partial answer to Galvin's question.

Theorem 1. *If $N \in L_0$ is an F_σ set and S a Sierpinski set, then there exists $t \in \mathbb{R}$ such that $(N + t) \cap S = \emptyset$.*

Received by the editors October 20, 1989 and, in revised form, February 12, 1990.
1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 04A15; Secondary 28A15.

©1991 American Mathematical Society
0002-9939/91 \$1.00 + \$.25 per page

The proof is based on the following simple observation:

Lemma 1. *Let $N \in L_0$ and $Q \subseteq \mathbb{R}$ be such that*

- (1) $N + Q \in L_0$ and
- (2) N can be shifted out of every countable set by a member of Q ; i.e., for every countable set $X \subseteq \mathbb{R}$, $Q \not\subseteq X - N$.

Then N can be shifted out of every Sierpinski set.

Proof. Let S be a Sierpinski set and let $X = S \cap (N + Q)$, so that X is countable. Then any member of Q that shifts N out of X shifts N out of S ; i.e., if $t \in Q$ and $(N + t) \cap X = \emptyset$, then $(N + t) \cap S = \emptyset$.

[EKM, Lemma 9] shows that, if $N \in L_0$, then there exists a perfect set Q such that $N + Q \in L_0$. It follows that every Sierpinski set can be shifted out of any measure-zero set N such that countably many translates of N cannot cover any perfect set. Recall that a set $X \subseteq \mathbb{R}$ has universal measure zero iff, for all atomless measures ν on the Borel sets, there is a Borel set of ν -measure zero covering X . Also recall that X has property s_0 iff for any perfect set P there exists a perfect set $Q \subseteq P$ disjoint from X . It is well known that neither universal-measure-zero sets nor sets with property s_0 can cover a perfect set by countably many translates (for this and other properties of these sets see [M]). We have the following:

Corollary. *If $M \subseteq \mathbb{R}$ is universally null or $M \in L_0$ and has property s_0 and S is a Sierpinski set, then $(M + t) \cap S = \emptyset$ for some $t \in \mathbb{R}$.*

For our main result, we need $\{p_k\}_{k \in \omega}$ -expansions. Let $\{p_k\}_{k \in \omega}$ be a sequence of natural numbers each greater than 1. Every real number $x \in [0, 1)$ has a unique $\{p_k\}_{k \in \omega}$ -expansion of the form $x = \sum_{k \in \omega} \frac{x_k}{p_0 \cdot p_1 \cdot \dots \cdot p_k}$, where $x_k \in p_k$ and $x_k \neq p_k - 1$ for infinitely many k . (We identify p_k with $\{0, 1, 2, \dots, p_k - 1\}$.) In what follows, x_k will be denoted by $(x)_k$.

Lemma 2. *Let $N \in L_0$ be an F_σ subset of $[0, 1)$. Then there exists a sequence $\{p_k\}_{k \in \omega}$ and a sequence $\{\Delta_k\}_{k \in \omega}$ such that, for all $k \in \omega$,*

$$(1) \quad p_k \geq 6, \Delta_k \subseteq p_k \quad \text{and} \quad |\Delta_k| \leq \frac{p_k}{3},$$

and there exists a countable set $Y \subseteq \mathbb{R}$ such that $N \subseteq D + Y$, where

$$D = \left\{ \sum_{k \in \omega} \frac{d_k}{p_0 \cdot p_1 \cdot \dots \cdot p_k} : d_k \in \Delta_k \right\}.$$

Proof. Let $N_0, N_1, \dots \subseteq [0, 1)$ be closed sets such that $N = \bigcup_{k \in \omega} N_k$. Define $I_n^i = [\frac{i}{2^n}, \frac{i+1}{2^n})$, where $n \in \omega$ and $i \in 2^n$. Also, let $Q_k(n) = |\{i: I_n^i \cap N_k \neq \emptyset\}|$. It is not hard to show that, since N_k is closed (in $\mathbb{R}$), $\lim_{n \rightarrow \infty} \frac{Q_k(n)}{2^n} = 0$. Using this fact, we define $\{p_k\}_{k \in \omega}$ by induction.

Let $m \in \omega$ be such that $2^m \geq 6$ and $\frac{Q_0(m)}{2^m} \leq \frac{1}{3}$. Put $p_0 = 2^m$ and

$\Delta_0 = \{i: I_m^i \cap N_0 \neq \emptyset\}$. At stage $k+1$, consider $I_k = [0, \frac{1}{p_0 \cdot p_1 \cdot \dots \cdot p_k})$ and

$$B_k = \left(N_k \cup N_k - \frac{1}{p_0 \cdot p_1 \cdot \dots \cdot p_k} \cup \dots \cup N_k - \frac{p_0 \cdot p_1 \cdot \dots \cdot p_k - 1}{p_0 \cdot p_1 \cdot \dots \cdot p_k} \right) \cap I_k.$$

Now pick m so that $2^m \geq 6$. Then, if we partition I_k into 2^m equal subintervals, at most $\frac{2m}{3}$ of them have nonempty intersections with B_k . Let $p_{k+1} = 2^m$ and $\Delta_{k+1} \subseteq p_{k+1}$ be a set of indices of those subintervals whose intersections with B_k are nonempty. Clearly conditions (1) are satisfied.

Now suppose D is as in Lemma 2 and $a \in N_{k_0}$ for some $k_0 \in \omega$. For $k \geq k_0$, $(a)_k \in \Delta_k$. Therefore, if we let Y be the set of all real numbers from $(-1, 1)$ with finite $\{p_k\}_{k \in \omega}$ -expansions, then $a \in D + Y$.

We now have to modify [EKM, Lemma 9].

Lemma 3. *Let $\{p_k\}_{k \in \omega}$ be a bounded sequence of natural numbers greater than 1. For each set $K \subseteq \omega$, let*

$$Q_K = \left\{ \sum_{k \in \omega} \frac{q_k}{p_0 \cdot p_1 \cdot \dots \cdot p_k} : q_k \in p_k \text{ if } k \in K \text{ and } q_k = 0 \text{ otherwise} \right\}.$$

Given $N \in L_0$, there exists an infinite set $K \subseteq \omega$ such that $N + Q_K \in L_0$.

Proof. The proof is almost the same as the original proof of Erdős, Kunen, and Mauldin. Note that if $f: \omega \rightarrow \omega$ is an increasing function, $I \subseteq \mathbb{R}$ is an interval, and $n \in \omega$, then $\mu(I + Q_{\text{ran}(f)}) \leq p_{f(0)} \cdot p_{f(1)} \cdot \dots \cdot p_{f(n)} (\mu(I) + \frac{1}{p_0 \cdot p_1 \cdot \dots \cdot p_{f(n)}})$. By making f increase rapidly enough we make $p_{f(0)} \cdot p_{f(1)} \cdot \dots \cdot p_{f(n)} / p_0 \cdot p_1 \cdot \dots \cdot p_{f(n)}$ as small as desired so that all steps in the proof given in [EKM] may be performed here without major changes. We leave the details to the reader. However, in the case relevant to the proof of Theorem 1, $p_k = 2^{r_k}$ for some $r_k \in \omega$. For such p_k , Lemma 3 follows easily. By [EKM, Lemma 9], we have an infinite set $H \subseteq \omega$ such that if

$$B_H = \left\{ \sum_{k \in \omega} \frac{q_k}{2^k} : q_k \in \{0, 1\} \text{ if } k \in H \text{ and } q_k = 0 \text{ otherwise} \right\},$$

then $N + B_H \in L_0$ and, if $b \in \omega$ is a bound for $\{r_k\}_{k \in \omega}$ and $H_k = \{k+i: i < b\}$, then there exists an infinite set $X \subseteq \omega$ such that, whenever $k, 1 \in X$, then $H_k \cap H_1 = \emptyset$ and $H_k \subseteq H$. Now simply let $K = \{k+b: k \in X\}$, and observe that $Q_k \subseteq B_H$.

Proof of Theorem 1. Let $N \in L_0$ be an F_σ set. Then $N' = (\bigcup_{n \in \omega} (N \pm n)) \cap [0, 1)$ is also an F_σ set, so that by Lemma 2 there exist sequences $\{p_k\}_{k \in \omega}$, $\{\Delta_k\}_{k \in \omega}$ and a set D such that $N' \subseteq D + Y$ for some countable $Y \subseteq \mathbb{R}$. Since in the proof of Lemma 2 the sets N_k can be made as small as desired, we may assume that the sequence $\{p_k\}_{k \in \omega}$ is bounded. Therefore, by Lemma 3, we have an infinite set $K = \{k_0, k_1, k_2, \dots\} \subseteq \omega$, so that $Q_K + N \in L_0$. By Lemma 1, it suffices to show that N does not cover Q_K by countable translations. Since N' as well as N may be covered by countably many translations of D , let us now

show that for any countable $X \subseteq \mathbb{R}$, $X + D \not\subseteq Q_K$. Let $X = \{x^{(n)} : n \in \omega\}$. By assuming that the addition in $X + D$ is modulo 1, we may consider $X \subseteq [0, 1)$. For each $i \in \omega$, pick $q_{k_i} \in (p_{k_i} - 1) \setminus [(\Delta_{k_i} \cup \Delta_{k_i} \oplus 1) \oplus (x^{(i)})_{k_i}]$, where $\oplus$ signifies algebraic addition modulo p_{k_i} . Set $q_k = 0$ for $k \notin K$ so that the number $q = \sum_{k \in \omega} \frac{q_k}{p_0 \cdot p_1 \cdots p_k}$ is in Q_K and is different from all numbers in $X + D$.

A similar result holds for generalized Sierpinski sets under Martin's Axiom (MA). Recall that $S \subseteq \mathbb{R}$ is a generalized Sierpinski set if $|S| = c$ and $|S \cap N| < c$ whenever $N \in L_0$. Clearly, Lemma 1 remains true for generalized Sierpinski sets if "countable X " is replaced by " $|X| < c$."

Theorem 2. Assume MA. If $N \in L_0$ is an F_σ set, then for every generalized Sierpinski set S there exists a $t \in \mathbb{R}$ such that $(N + t) \cap S = \emptyset$.

Proof. Let $N \in L_0$ be an F_σ set. It suffices to show that N satisfies the generalized Lemma 1 as shown above. Let N' , $\{p_k\}_{k \in \omega}$, $\{\Delta_k\}_{k \in \omega}$, and K be as in the proof of Theorem 1. We would like to show that $D + X \not\subseteq Q_K$ for any $X \subseteq \mathbb{R}$, where $|X| < c$. Let $\mathbf{P} = \{(f, E) : E \subseteq X \text{ is a finite subset and } f \text{ is a finite function, } \text{dom}(f) \subseteq \omega, f(i) \in p_1 \text{ and for every } e \in E \text{ there exists an } i \in K \cap \text{dom}(f) \text{ such that } f(i) \in p_1 \setminus (D + e)_1 \text{ and } f(i) = 0 \text{ for } i \in \text{dom}(f) \setminus K\}$. Define a partial order on $\mathbf{P}$ simply as $(f_2, E_2) \leq (f_1, E_1)$ if $f_2 \supseteq f_1$ and $E_2 \supseteq E_1$. Clearly $(\mathbf{P}, \leq)$ is a ccc poset and $D_F = \{(f, E) : E \supseteq F\}$ and $D_n = \{(f, E) : n \in \text{dom}(f)\}$ are dense subsets of $\mathbf{P}$ for all finite $F \subseteq X$ and $n \in \omega$. By Martin's Axiom, there exists a filter $G \subseteq \mathbf{P}$ which meets all D_F and D_n . Let $q = \bigcup \{f : (f, E) \in G\}$. It is easy to see that

$$\sum_{k \in \omega} \frac{q(k)}{p_0 \cdot p_1 \cdots p_k} \in Q_K \setminus (D + X).$$

Added in proof. I. Reclaw and A. W. Miller have obtained independently a short, elegant proof of the following theorem.

Theorem. If S is a Sierpinski set and F is a F_σ set of measure zero, then $\{x : (x + F) \cap S = \emptyset\}$ is comeager.

T. Bartoszynski and I. Reclaw pointed out that for some dense G_δ sets N there are sets Q such that conditions (1) and (2) of Lemma 1 hold.

REFERENCES

- [BI] T. Bartoszynski and J. I. Ihoda, *On Sierpinski sets*, Proc. Amer. Math. Soc. **108** (1990), 507–512.
- [EKM] P. Erdős, K. Kunen, and R. D. Mauldin, *Some additive properties of sets of real numbers*, Fund. Math. **113** (1981), 187–194.
- [M] A. W. Miller, *Special subsets of the real line*, in *Handbook of set-theoretic topology* (K. Kunen and J. E. Vaughan, eds.), Elsevier Science Publishers B. V., 1984.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SCRANTON, SCRANTON, PENNSYLVANIA 18510

DEPARTMENT OF MATHEMATICS, PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK, PENNSYLVANIA 16802