LEVEL CROSSINGS OF A RANDOM TRIGONOMETRIC POLYNOMIAL

KAMBIZ FARAHMAND

(Communicated by William D. Sudderth)

ABSTRACT. This paper provides an asymptotic estimate for the expected number of K -level crossings of the random trigonometric polynomial $g_1 \cos x + g_2 \cos 2x + \dots + g_n \cos nx$, where g_j ($j = 1, 2, \dots, n$) are independent normally distributed random variables with mean μ and variance one. It is shown that the result for $K = \mu = 0$ remains valid for any finite constant μ and any K such that $(K^2/n) \rightarrow 0$ as $n \rightarrow \infty$.

1. INTRODUCTION

Let

$$(1.1) \quad T(x) \equiv T_n(x, \omega) = \sum_{j=1}^n g_j(\omega) \cos jx$$

where $g_1(\omega), g_2(\omega), \dots, g_n(\omega)$ is a sequence of independent random variables defined on a probability space $(\Omega, \mathcal{A}, P)$, each normally distributed with finite mathematical expectation μ and variance one. The set of equations $y = T(x)$ represents a family of curves in the xy -plane. Some years ago Dunnage [2] showed that the number of times all save a certain exceptional set of these curves crosses the x -axis in the interval $0 \leq x \leq 2\pi$ is $(2n)/\sqrt{3} + O\{n^{11/13}(\log n)^{3/13}\}$. The measure of this exceptional set does not exceed $(\log n)^{-1}$. Later Sambandham and Renganathan [7] and Farahmand [3] considered the case when $\mu \neq 0$ and found that the expected number of crossings of this family with the x -axis is asymptotic to $2n\sqrt{3}$. This asymptotic number of crossings remains invariant in the work of Farahmand [4] when, for $\mu = 0$, he considered the crossings of this family with the line $y = K$ (i.e., with level K), where K is any real value constant such that $(K^2/n) \rightarrow 0$ as $n \rightarrow \infty$.

Here we consider the effect of μ being nonzero on the K -level crossings of this family of curves, and show even in this case the above asymptotic number

---

Received by the editors July 6, 1989 and, in revised form, September 19, 1989.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 60G99; Secondary 42BXX.

Key words and phrases. Number of real roots, Kac-Rice formula, random trigonometric polynomial.

of crossings persists. Denote by $N(\alpha, \beta) \equiv N_K(\alpha, \beta)$ the number of times that the family of curves $y = T(x)$ crosses the line $y = K$ on the interval $\alpha \leq x \leq \beta$, and let $EN(\alpha, \beta)$ be its expectation. We prove the following.

Theorem. *If the coefficients of $T(x)$ in (1.1) are independent normally distributed random variables with mean μ and variance one, then for all sufficiently large n and any constant K the expected number of real roots of the equation $T(x) = K$ satisfies*

$$EN(0, 2\pi) = (2n)/\sqrt{3} + O(n^{3/4}) \quad \text{if } K = O(n^{3/8})$$

and

$$EN(0, 2\pi) = (2n)/\sqrt{3} + o(n) \quad \text{if } K = o(\sqrt{n}).$$

2. A FORMULA FOR THE NUMBER OF CROSSINGS

Let

$$\Phi(t) = (2\pi)^{-1/2} \int_{-\infty}^t \exp(-y^2/2) dy$$

and

$$\phi(t) = \Phi'(t) = (2\pi)^{-1/2} \exp(-t^2/2),$$

then by using the expected number of level crossings given by Cramer and Leadbetter [1, p. 285] for our equation $T(x) - K = 0$ we can obtain

$$(2.1) \quad EN(\alpha, \beta) = \int_{\alpha}^{\beta} (B/A)(1 - \lambda^2)^{1/2} \phi(\xi/A) [2\phi(\eta) + \eta\{2\Phi(\eta) - 1\}] dx$$

where

$$\begin{aligned} A^2 &= \text{Var}\{T(x) - K\}, & B^2 &= \text{Var}\{T'(x)\}, \\ \lambda &= (AB)^{-1} \text{cov}\{[T(x) - K], T'(x)\}, & \xi &= E\{T(x) - K\}, \\ \eta &= B^{-1}(1 - \lambda^2)^{-1/2}(\gamma - B\lambda\xi/A) & \text{and } \gamma &= E\{T'(x)\}. \end{aligned}$$

Since the coefficients of $T(x)$ are independent normal random variables with mean μ and variance one we can easily find that

$$\begin{aligned} \xi &= \mu \sum_{j=1}^n \cos jx - K = m_1 - K, \quad \text{say,} \\ \gamma &= -\mu \sum_{j=1}^n j \sin jx = -m_2, \quad \text{say} \\ A^2 &= \sum_{j=1}^n \cos^2 jx, & B^2 &= \sum_{j=1}^n j^2 \sin^2 jx, \\ C &= \sum_{j=1}^n j \sin jx \cos jx, & \lambda &= -C/AB, \\ \eta &= \{C(m_1 - K) - A^2 m_2\}/A\Delta & \text{and } \Delta^2 &= A^2 B^2 - C^2. \end{aligned}$$

Hence from (2.1) and since $\Phi(t) = 1/2 + (\pi)^{-1/2} \operatorname{erf}(t/\sqrt{2})$, we have the extension of the Kac-Rice formula [5].

$$\begin{aligned}
 & EN(\alpha, \beta) \\
 &= \int_{\alpha}^{\beta} (\Delta/\pi A^2) \exp[\{-B^2(m_1 - K)^2 - A^2 m_2^2 + 2m_2 C(m_1 - K)\}/(2\Delta^2)] dx \\
 &\quad + \int_{\alpha}^{\beta} (\sqrt{2}/\pi) A^{-3} |C(m_1 - K) - A^2 m_2| \exp\{-(m_1 - K)^2/2A^2\} \\
 &\quad \times \operatorname{erf}\{|C(m_1 - K) - A^2 m_2|/\sqrt{2}A\Delta\} dx \\
 (2.2) \quad &= \int_{\alpha}^{\beta} I_1(x) dx + \int_{\alpha}^{\beta} I_2(x) dx.
 \end{aligned}$$

3. OVERVIEW OF THE PROOF OF THE THEOREM AND SOME APPROXIMATIONS

In order to evaluate the expected number of real roots we divide the real roots into two groups: (i) those lying in the intervals $(0, \varepsilon)$, $(\pi - \varepsilon, \pi + \varepsilon)$ and $(2\pi - \varepsilon, 2\pi)$ and (ii) those lying in the intervals $(\varepsilon, \pi - \varepsilon)$ and $(\pi + \varepsilon, 2\pi - \varepsilon)$. For the roots (i) which, it so happens, are negligible, we need some modification to apply Dunnage's [2] approach, which is based on an application of Jensen's theorem [8, p. 125] or [6, p. 332]. For roots (ii) we use the Kac-Rice formula (2.2). The ε should be chosen positive and small enough to facilitate handling type (i) roots, but large enough to allow for the approximation needed to discuss type (ii) roots. We choose $\varepsilon = n^{-1/4}$.

The function

$$S(x) = \sin(2n+1)x / \sin x$$

and, defined by continuity at $x = j\pi$, will occur frequently and so it is convenient to collect together some related inequalities. Since for $\varepsilon \leq x \leq \pi - \varepsilon$ or $\pi + \varepsilon \leq x \leq 2\pi - \varepsilon$ we have $|S(x)| < 1/\sin \varepsilon$, we can obtain

$$(3.1) \quad S(x) = O(1/\varepsilon).$$

Further

$$(3.2) \quad S'(x) = (2n+1) \cos(2n+1)x / \sin x - \cot x S(x) = O(n/\varepsilon)$$

and

$$\begin{aligned}
 (3.3) \quad S''(x) = & -(2n+1)^2 S(x) - (2n+1) \cos x \cos(2n+1)x \sin^{-2} x \\
 & - \cot x S'(x) + \operatorname{cosec}^2 x S(x) = O(n^2/\varepsilon).
 \end{aligned}$$

Hence from (3.1) and since by expanding $\sin x(1 + 2\sum_{j=1}^n \cos 2jx)$, we can show that

$$(3.4) \quad \sum_{j=1}^n \cos 2jx = (1/2)\{S(x) - 1\}$$

for this range of x we have

$$(3.5) \quad A^2 = (1/2) \sum_{j=1}^n (1 + \cos 2jx) = n/2 + (1/4)\{S(x) - 1\} = n/2 + O(1/\varepsilon).$$

Also from (3.3) and since from (3.4)

$$S''(x) = -8 \sum_{j=1}^n j^2 \cos 2jx = 4 \sum_{j=1}^n j^2 (2 \sin^2 jx - 1)$$

we have

$$(3.6) \quad B^2 = n(n+1)(2n+1)/12 + (1/8)S''(x) = n^3/6 + O(n^2/\varepsilon).$$

From (3.2) we obtain

$$(3.7) \quad C = -(1/2) \frac{d}{dx}(A^2) = -(1/8)S'(x) = O(n/\varepsilon).$$

Also from (3.1), (3.2), and (3.4) we can easily obtain the two following estimates:

$$(3.8) \quad m_1 = (\mu/2)\{S(x/2) - 1\} = O(1/\varepsilon)$$

and

$$(3.9) \quad m_2 = (-\mu/4)S'(x/2) = O(n/\varepsilon).$$

Then (3.5), (3.6), and (3.7) give

$$(3.10) \quad \Delta^2 = n^4/12 + O(n^3/\varepsilon).$$

4. PROOF OF THE THEOREM

From (2.2) and (3.5)–(3.10) we can obtain

$$(4.1) \quad \int_{\varepsilon}^{\pi-\varepsilon} I_1(x) dx = (n/\sqrt{3})\{1 + O(\varepsilon)\} \exp\{-K^2/n + O(1/n\varepsilon^2)\}$$

and

$$(4.2) \quad \int_{\varepsilon}^{\pi-\varepsilon} I_2(x) dx = O\{(n^{1/2}/\varepsilon) \exp(-1/n\varepsilon^2 - K^2/n)\}.$$

Hence from (2.2), (4.1), and (4.2) for $K = O(n^{3/8})$, we have

$$(4.3) \quad EN(\varepsilon, \pi - \varepsilon) = EN(\pi + \varepsilon, 2\pi - \varepsilon) = n/\sqrt{3} + O(n^{3/4})$$

and for $K = o(\sqrt{n})$

$$(4.4) \quad EN(\varepsilon, \pi - \varepsilon) = EN(\pi + \varepsilon, 2\pi - \varepsilon) = n/\sqrt{3} + o(n).$$

Now we show that the expected number of real roots in the intervals $(0, \varepsilon)$, $(\pi - \varepsilon, \pi + \varepsilon)$, and $(2\pi - \varepsilon, 2\pi)$ is negligible. The period of $T(x)$ is 2π and so the number of zeros in $(0, \varepsilon)$ and $(2\pi - \varepsilon, 2\pi)$ is the same as the number in $(-\varepsilon, \varepsilon)$. We shall therefore confine ourselves to this last interval; the interval

$(\pi - \varepsilon, \pi + \varepsilon)$ can be treated in exactly the same way to give the same result. We consider the random integral function of the complex variable z ,

$$T(z, \omega) - K = \sum_{j=1}^n g_j(\omega) \cos jz - K.$$

We are seeking an upper bound to the number of real roots in the segment of the real axis joining the points $\pm \varepsilon$, and this certainly does not exceed the number of real roots in the circle $|z| < \varepsilon$. Let $N(r) \equiv N(r, \omega, K)$ denote the number of real roots of $T(z, \omega) - K = 0$ in $|z| < r$. By Jensen's theorem [8, p. 125] or [6, p. 332]

$$\begin{aligned} \int_{\varepsilon}^{2\varepsilon} r^{-1} N(r) dr &\leq \int_0^{2\varepsilon} r^{-1} N(r) dr \\ &= (2\pi)^{-1} \int_0^{2\pi} \log \left| \{T(2\varepsilon e^{ix}, \omega) - K\} / \{T(0) - K\} \right| dx, \end{aligned}$$

assuming that $T(0) \neq K$, from which we have

$$(4.5) \quad N(\varepsilon) \log 2 \leq (2\pi)^{-1} \int_0^{2\pi} \log \left| \{T(2\varepsilon e^{ix}, \omega) - K\} / \{T(0) - K\} \right| dx.$$

Now since the distribution function of $T(0, \omega) = \sum_{j=1}^n g_j(\omega)$ is

$$G(x) = (2\pi n)^{-1/2} \int_{-\infty}^x \exp\{-(t - \mu)^2 / 2n\} dt$$

we can see that for any positive ν

$$\begin{aligned} \text{Prob}(-e^{-\nu} \leq T(0) - K < e^{-\nu}) &= (2\pi n)^{-1/2} \int_{K-e^{-\nu}}^{K+e^{-\nu}} \exp\{-(t - \mu)^2 / 2n\} dt \\ (4.6) \quad &< (2/n\pi)^{1/2} e^{-\nu}. \end{aligned}$$

Also we have

$$(4.7) \quad |T(2\varepsilon e^{ix})| = \left| \sum_{j=1}^n g_j \cos(2j\varepsilon e^{ix}) \right| \leq ne^{2n\varepsilon} \max |g_j|$$

where the maximum is taken over $1 \leq j \leq n$. The distribution function of $|g_j|$ is

$$F(x) = \begin{cases} \sqrt{2/\pi} \int_0^x \exp\{-(t - \mu)^2 / 2\} dt, & x \geq 0, \\ 0, & x < 0, \end{cases}$$

and so for any positive ν and all sufficiently large n

$$\begin{aligned} \text{Prob}\{\max |g_j| > ne^{\nu}\} &\leq n \text{Prob}\{|g_1| > ne^{\nu}\} \\ &= n\sqrt{2\pi} \int_{ne^{\nu}}^{\infty} \exp\{-(t - \mu)^2 / 2\} dt \\ (4.8) \quad &\sim \sqrt{2/\pi} \exp\{-\nu - (ne^{\nu} - \mu)^2 / 2\}. \end{aligned}$$

Therefore from (4.7) and (4.8) except for sample functions in an ω -set of measure not exceeding $(2/\pi)^{1/2} \exp\{-\nu - (ne^\nu - \mu)^2/2\}$,

$$(4.9) \quad |T(2\epsilon e^{ix})| < n^2 \exp(2n\epsilon + \nu).$$

Hence from (4.6), (4.9) and since

$$|n^2 \exp(2n\epsilon + \nu) - K| < 2n^2 \exp(2n\epsilon + \nu)$$

if $K = o(\sqrt{n})$ or if $K = O(n^{3/8})$, we obtain

$$(4.10) \quad \begin{aligned} |\{T(2\epsilon e^{ix}, \omega) - K\} / \{T(0, \omega) - K\}| &\leq e^\nu |2n^2 \exp(2n\epsilon + \nu) - K| \\ &\leq 2n^2 \exp(2n\epsilon + 2\nu) \end{aligned}$$

except for sample functions in an ω -set of measure not exceeding

$$(2/n\pi)^{1/2} e^{-\nu} + (2/\pi)^{1/2} \exp\{-\nu - (ne^\nu - \mu)^2/2\}.$$

Therefore from (4.5) and (4.10) we can show that outside the exceptional set

$$(4.11) \quad N(\epsilon) \leq (\log 2 + 2 \log n + 2n\epsilon + 2\nu) / \log 2.$$

Because $\epsilon = n^{-1/4}$, it follows from (4.11) and for all sufficiently large n

$$(4.12) \quad \text{Prob}\{N(\epsilon) > 3n\epsilon + 2\nu\} \leq (2/n\pi)^{1/2} e^{-\nu} + (2/\pi)^{1/2} \exp\{-\nu - (ne^\nu - \mu)^2/2\}.$$

Let $n' = [3n^{3/4}]$ be the greatest integer less than or equal to $3n^{3/4}$; then from (4.12) and for n large enough we obtain

$$\begin{aligned} EN(\epsilon) &= \sum_{j>0} \text{Prob}\{N(\epsilon) \geq j\} \\ &= \sum_{1 \leq j \leq n'} \text{Prob}\{N(\epsilon) > j\} + \sum_{j \geq 1} \text{Prob}\{N(\epsilon) > n' + j\} \\ &\leq n' + (2/n\pi)^{1/2} \sum_{j \geq 1} e^{-j/2} + (2/\pi)^{1/2} \sum_{j \geq 1} \exp\{-j/2 - (ne^j - \mu)^2/2\} \\ (4.13) \quad &= O(n^{3/4}). \end{aligned}$$

Finally (4.3), (4.4), and (4.13) complete proof of the theorem.

ACKNOWLEDGMENT

The author wishes to thank the referee for his valuable comments.

REFERENCES

1. H. Cramer and M. R. Leadbetter, *Stationary and related stochastic processes*, Wiley, New York, 1967.
2. J. E. A. Dunnage, *The number of real zeros of a random trigonometric polynomial*, Proc. London Math. Soc. **16** (1966), 53-84.

3. K. Farahmand, *On the number of real zeros of a random trigonometric polynomial: coefficient with non-zero mean*, Stochastic Anal. and Appl. **5** (1987), 379–386.
4. ———, *On the average number of level crossings of a random trigonometric polynomial* Annal. of Prob. **18** (1990).
5. S. O. Rice, *Mathematical theory of random noise*, Bell System Tech. J. **25** (1945), 46–156.
6. W. Rudin, *Real and complex analysis*, 2nd ed. McGraw-Hill, 1974.
7. M. Sambandham and N. Renganathan, *On the number of real zeros of a random trigonometric polynomial: coefficient with non-zero means*, J. Indian Math. Soc. **45** (1981), 193–203.
8. E. C. Titchmarsh, *The theory of functions*, 2nd ed. Oxford University Press, 1939.

DEPARTMENT OF MATHEMATICAL STATISTICS, UNIVERSITY OF CAPE TOWN, RONDEBOSCH 7700,
SOUTH AFRICA