COMPARISON THEOREMS FOR THE ν -ZEROES OF LEGENDRE FUNCTIONS $P_\nu^m(z_0)$ WHEN $-1 < z_0 < 1$

FRANK E. BAGINSKI

(Communicated by Kenneth R. Meyer)

ABSTRACT. We consider the problem of ordering the elements of $\{\nu_j^m(z_0)\}$, the set of ν -zeroes of Legendre functions $P_\nu^m(z_0)$ for $m = 0, 1, \dots$ and $z_0 \in (-1, 1)$. In general, we seek to determine conditions on (m, j) and (n, i) under which we can assert that $\nu_j^m < \nu_i^n$. A number of such results were established in [2] for $z_0 \in [0, 1)$, and in the work that we present here we extend a number of these to the case $z_0 \in (-1, 1)$. In addition, we prove $\nu_{j+1}^m < \nu_j^{m+2}$ for $z_0 \in (-1, 0)$ and $\nu_2^3 < \nu_1^6$ for $z_0 \in (-1, 1)$. Using the results established here and in [2], we are able to determine the ordering of the first eleven ν -zeroes of $P_\nu^m(z_0)$ for $0 < z_0 < 1$ and show that the twelfth ν -zero is not necessarily distinct.

1. INTRODUCTION

For a fixed $z_0 \in (-1, 1)$ and $m = 0, 1, \dots$, we will let $\{\nu_j^m(z_0)\}$ denote the set of positive ν -zeroes of Legendre functions $P_\nu^m(z_0)$. The principal goal is determine conditions on (m, j) and (n, i) under which we can assert that $\nu_j^m < \nu_i^n$. One such result which follows from the Sturm-Liouville theory is that

$$(1.1) \quad \nu_j^m(z_0) < \nu_j^{m+1}(z_0) < \nu_{j+1}^m(z_0), \quad -1 < z_0 < 1$$

(See [10].) The problem of ordering the ν_j^m 's when $0 \leq z_0 < 1$ was first considered in [2], where the following results were established:

$$(1.2) \quad \nu_j^{m+2}(z_0) < \nu_{j+1}^m(z_0), \quad 0 < z_0 < 1,$$

$$(1.3) \quad \nu_j^{m+2}(0) = \nu_{j+1}^m(0),$$

$$(1.4) \quad \nu_2^0(z_0) < \nu_1^3(z_0), \quad \nu_2^1(z_0) < \nu_1^4(z_0), \quad \nu_3^0(z_0) < \nu_1^5(z_0), \quad 0 < z_0 < 1.$$

Received by the editors August 8, 1989; some of these results were presented at the 857th Meeting of the AMS at University Park, PA, April 7-8, 1990.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 33A45.

Key words and phrases. Zeroes of Legendre functions, Sturm-Liouville theory.

Combining (1.1)–(1.4), it follows that the first ten ν -zeroes are

$$(1.5) \quad \nu_1^0 < \nu_1^1 < \nu_1^2 < \nu_2^0 < \nu_1^3 < \nu_2^1 < \nu_1^4 < \nu_2^2 < \nu_3^0 < \nu_1^5, \quad 0 < z_0 < 1.$$

The ordering in (1.5) is unique.

The above results lead to several additional questions: Which are the next ν_j^m 's in the chain of inequalities in (1.5)? How do ν_{j+1}^m and ν_j^{m+2} compare for $z_0 \in (-1, 1)$? In the following, we will consider these and related questions. The main results of this paper are contained in §3.

On the basis of numerical calculations, it was conjectured in [2] that the ν -zero which followed ν_1^5 in (1.5) was ν_2^3 . An analytical proof of this follows from the inequality,

$$(1.6) \quad \nu_2^3(z_0) < \nu_1^6(z_0), \quad -1 < z_0 < 1,$$

which is established in Theorem 1. In Lemma 2, we show that

$$(1.7) \quad \nu_{j+1}^m(z_0) < \nu_j^{m+2}(z_0), \quad -1 < z_0 < 0.$$

Theorem 3 combines (1.1)–(1.3) with Lemma 2 and gives the relative ordering of $\nu_i^m(z_0)$, $\nu_j^{m+1}(z_0)$, and $\nu_k^{m+2}(z_0)$ for $-1 < z_0 < 1$.

In addition, we will show that the inequalities in (1.4) hold for all $z_0 \in (-1, 1)$. (See Theorem 2.) Although (1.1)–(1.6) imply that the first eleven ν -zeroes of $P_\nu^m(z_0)$ are distinct for all $0 < z_0 < 1$, in §4 we show that the twelfth ν -zero is not necessarily distinct. Moreover, Theorem 3 shows that the ordering in (1.5) is not preserved for $-1 < z_0 \leq 0$.

2. PRELIMINARIES

In this section, we present some properties of the Legendre functions and their zeroes which will be needed in §3. For convenience, here and in the following sections m will denote a nonnegative integer and n, i, j, k will denote positive integers, unless otherwise stated.

The solution $y = P_\nu^m(z)$ that satisfies

$$(2.1a) \quad \frac{d}{dz} \left((1-z^2) \frac{d}{dz} y \right) + \left(\nu(\nu+1) - \frac{m^2}{1-z^2} \right) y = 0, \quad -1 < z < 1,$$

$$(2.1b) \quad y(1) \text{ is bounded,}$$

is called a Legendre function of the first kind of degree ν and order m . (See [4] for a general discussion of the Legendre functions and their properties.) If $-1 < z \leq 1$ and $\nu > 0$, then $P_\nu^m(z)$ can be expressed [3, p. 148] as

$$(2.2) \quad P_\nu^m(z) = \frac{(-1)^m \Gamma(\nu+m+1)}{2^m m! \Gamma(\nu-m+1)} (1-z^2)^{m/2} \sum_{n=0}^{\infty} \frac{(1+m+\nu)_n (m-\nu)_n}{(m+1)_n n! 2^n} (1-z)^n,$$

where $\Gamma(z)$ is the gamma function and $(a)_n$ denotes the Pochhammer symbol,

$$(a)_0 = 1,$$

$$(a)_n = a(a+1) \dots (a+n-1).$$

Later, we will need the following identity:

$$(2.3) \quad P_{\nu}^{m+2}(z) + 2(m+1)z(1-z^2)^{-1/2}P_{\nu}^{m+1}(z) + (\nu-m)(\nu+m+1)P_{\nu}^m(z) = 0$$

(See [3, p. 161].) For a fixed $z_0 \in (-1, 1)$, the pairs $(\nu(\nu+1), y(z))$ satisfying (2.1) and $y(z_0) = 0$ will be denoted by

$$(2.4) \quad (\nu_j^m(\nu_j^m+1), P_{\nu_j^m}^m(z)),$$

where $P_{\nu_j^m}^m(z_0) = 0$, $m = 0, 1, \dots$, and $j = 1, 2, \dots$.

The next result summarizes a few of the important properties of the ν_j^m 's. Its proof follows the arguments in [2, Lemma 1] with some minor modifications.

Lemma 1. *Let m be a nonnegative integer. There exists a unique sequence $\{\nu_j^m(\tau) | j = 1, 2, \dots\}$ such that for every j , the function $\nu = \nu_j^m(\tau)$ satisfies*

$$(*) \quad P_{\nu}^m(\tau) = 0, \quad \text{for all } \tau \in (-1, 1).$$

Moreover, each $\nu_j^m(\tau)$ is analytic and strictly increasing as a function of τ for $\tau \in (-1, 1)$.

It will be convenient at times to consider $P_{\nu}^m(\cos \phi)$ for $0 < \phi \leq \phi_0 < \pi$ where $z = \cos \phi$ and $z_0 = \cos \phi_0$. As a function of ϕ_0 , we see from Lemma 1 that $\nu_j^m(\phi_0)$ is decreasing for $\phi_0 \in (0, \pi)$. It will be clear from the context whether ν_j^m is to be considered as a function of z_0 or as a function of $\phi_0 = \arccos(z_0)$.

A straightforward calculation shows that if $P_{\nu}^m(z)$ is a solution of (2.1), then $u = \sqrt{\sin \phi} P_{\nu}^m(\cos \phi)$ satisfies

$$(2.5) \quad u'' + \left(\left(\nu + \frac{1}{2} \right)^2 + \frac{1-4m^2}{4 \sin^2 \phi} \right) u = 0.$$

From [7, p. 17], we see that $v = \sqrt{\phi} J_m((\nu + \frac{1}{2})\phi)$ is a solution of

$$v'' + \left(\left(\nu + \frac{1}{2} \right)^2 + \frac{1-4m^2}{4\phi^2} \right) v = 0,$$

where J_m is the Bessel function of order m . From the Sturm-Liouville theory (See [10, Chapter 7]), $P_{\nu_j^m}^m(z)$ has $j-1$ z -zeroes on $(z_0, 1)$. Moreover, for a general ν (not necessarily one of the ν_j^m 's), we see from (2.5) that there are $[\nu - m]$ z -zeroes of $P_{\nu}^m(z)$ on $(-1, 1)$, where $[x] = n$ for $n \leq x < n+1$. $P_n^m(z)$ has exactly $n-m$ z -zeroes on $(-1, 1)$ and P_{n+1}^m has exactly $n-m+1$ z -zeroes on $(-1, 1)$. (See [6, p. 246].) To see this, suppose $\nu = \nu^*$ and $n < \nu^* < n+1$. By applying the Sturm Comparison Theorem [5] to the solutions of (2.5) for $\nu = n$, ν^* and $n+1$, respectively, we see that $P_{\nu^*}^m(z)$ must have at least $n-m$ z -zeroes on $(-1, 1)$ and at most $n-m+1$ z -zeroes on $(-1, 1)$. We conclude that $P_{\nu^*}^m(z)$ has exactly $[\nu^* - m]$ z -zeroes on $(-1, 1)$.

For fixed ν , we will denote the z -zeroes of $P_\nu^m(z)$ that are between -1 and 1 by $z_{\nu,i}^m$, where $z_{\nu,i}^m > z_{\nu,i+1}^m$. By definition of the $\nu_j^m(z_0)$'s, it follows that

$$(2.6) \quad \nu_j^m(z_{\nu,j}^m) = \nu.$$

We define $\phi_{\nu,j}^m$ to be the solution of

$$z_{\nu,j}^m = \cos(\phi_{\nu,j}^m)$$

such that $0 < \phi_{\nu,j}^m < \pi$. It follows that $\phi_{\nu,i}^m < \phi_{\nu,i+1}^m$.

From (2.2), we see that

$$(2.7) \quad P_m^m(z_0) = \frac{(-1)^m \Gamma(2m+1)}{2^m m!} (1 - z_0^2)^{m/2}.$$

From Rodrigue's formula [6, p. 174] and Rolle's Theorem, it follows that $P_{m+2j-1}^m(z)$ has $j-1$ z -zeroes on $(0, 1)$ and $\nu_j^m = m + 2j - 1$ when $z_0 = 0$. Similarly, $P_{m+j-1}^m(z)$ has $j-1$ z -zeroes on $(-1, 1)$ and

$$\lim_{z_0 \rightarrow -1+} \nu_j^m(z_0) = m + j - 1.$$

Since $\nu_j^m(z_0)$ is increasing on $(-1, 1)$, we see that

$$(2.8) \quad m + j - 1 < \nu_j^m(z_0) < m + 2j - 1, \quad -1 < z_0 < 0.$$

From [8], we have

$$(2.9) \quad \frac{1}{3} < \frac{1}{\sin^2 \phi} - \frac{1}{\phi^2} < \alpha(\bar{\phi}), \quad 0 < \phi < \bar{\phi} \leq \pi/2,$$

where $\alpha(\phi) = \sin^{-2} \phi - \phi^{-2}$. Note that $\lim_{\phi \rightarrow 0+} \alpha(\phi) = 1/3$. From (2.9), we see that

$$(2.10) \quad \frac{\phi^2}{1 + \alpha(\bar{\phi})\phi^2} < \sin^2 \phi < \frac{\phi^2}{1 + \frac{1}{3}\phi^2}, \quad 0 < \phi < \bar{\phi} \leq \pi/2.$$

Multiplying (2.9) by $1 - 4m^2$ with $m \geq 1$, we obtain

$$(2.11) \quad \frac{1 - 4m^2}{4\phi^2} - \frac{h^2}{4} > \frac{1 - 4m^2}{4\sin^2 \phi} > \frac{1 - 4m^2}{4\phi^2} - \frac{k^2(\bar{\phi})}{4}, \quad 0 < \phi < \bar{\phi} \leq \pi/2,$$

where $h^2 = (4m^2 - 1)/3$ and $k^2(\bar{\phi}) = (4m^2 - 1)\alpha(\bar{\phi})$.

Next, we consider the following pair of differential equations and their re-

spective solutions which are related to Bessel's equation [7, p. 17],

$$U'' + \left(\left(\nu + \frac{1}{2} \right)^2 - \frac{h^2}{4} + \frac{1 - 4m^2}{4\phi^2} \right) U = 0, \quad 0 < \phi < \bar{\phi},$$

$$U = \sqrt{\phi} J_m \left(\sqrt{\left(\nu + \frac{1}{2} \right)^2 - \frac{h^2}{4}} \phi \right),$$

$$V'' + \left(\left(\nu + \frac{1}{2} \right)^2 - \frac{k^2(\bar{\phi})}{4} + \frac{1 - 4m^2}{4\phi^2} \right) V = 0, \quad 0 < \phi < \bar{\phi},$$

$$V = \sqrt{\phi} J_m \left(\sqrt{\left(\nu + \frac{1}{2} \right)^2 - \frac{k^2}{4}} \phi \right).$$

Let u be a solution of (2.5). From (2.11) and the Sturm Comparison Theorem, it follows that the k th zero of U occurs before the k th zero of u and the k th zero of u occurs before the k th zero of V . In particular, we have for $m \geq 1$,

$$(2.12a) \quad \frac{j_k^m}{\sqrt{\left(\nu + \frac{1}{2} \right)^2 - \frac{h^2}{4}}} < \phi_{\nu, k}^m < \frac{j_k^m}{\sqrt{\left(\nu + \frac{1}{2} \right)^2 - \frac{k^2}{4}}},$$

where j_k^m is the k th positive zero of $J_m(z)$. If $m = 0$, we find that

$$(2.12b) \quad \frac{j_k^0}{\sqrt{\left(\nu + \frac{1}{2} \right)^2 + \frac{a}{4}}} < \phi_{\nu, k}^0 < \frac{j_k^0}{\sqrt{\left(\nu + \frac{1}{2} \right)^2 + \frac{1}{12}}}.$$

3. ORDERING THE ν -ZEROES OF LEGENDRE FUNCTIONS

This section contains the principle results of this paper. We begin with a comparison of ν_2^3 and ν_1^6 :

Theorem 1. $\nu_2^3(z_0) < \nu_1^6(z_0)$ for all $-1 < z_0 < 1$.

Proof. Here, it will be convenient to let $z_0 = \cos \phi_0$ and to consider ν_1^6 and ν_2^3 as functions of ϕ_0 . First, we will show that if $\nu = \nu_1^6 = \nu_2^3$, then $\nu > 7$. Then, we will show that $\nu = \nu_1^6 = \nu_2^3$ is impossible if $\nu > 7$.

Part 1. Since $\nu_2^3(\phi_0)$ and $\nu_1^6(\phi_0)$ are decreasing in ϕ_0 , by (2.8), we have

$$\lim_{\phi_0 \rightarrow \pi^-} \nu_2^3(\phi_0) = 4, \quad \lim_{\phi_0 \rightarrow \pi^-} \nu_1^6(\phi_0) = 6.$$

Since $\nu_1^6(\phi_0) > 6$ for $\phi_0 \in (0, \pi)$ and $\nu_2^3(\phi_0) < 6$ for $\phi_0 \in (\pi/2, \pi)$, we see that if $\nu = \nu_2^3(\phi_0) = \nu_1^6(\phi_0)$ for some ϕ_0 , then $\nu \geq 6$ and $0 < \phi_0 < \pi/2$.

Next, suppose that $\nu = \nu_2^3 = \nu_1^6$ for some $z_0 = \cos \phi_0$ and $0 < \phi_0 < \pi/2$. From (2.3) we see that

$$(A) \quad \begin{pmatrix} 10z_0(1 - z_0^2)^{-1/2} & (\nu - 4)(\nu + 5) \\ 1 & 8z_0(1 - z_0^2)^{-1/2} \end{pmatrix} \begin{pmatrix} P_\nu^5(z_0) \\ P_\nu^4(z_0) \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}.$$

Since by (1.1), $P_\nu^5(z_0)$ and $P_\nu^4(z_0)$ cannot vanish simultaneously, we see that the determinant of the 2×2 matrix in (A) must be zero, and we are led to the condition that

$$(B) \quad z_0^2 = \frac{\nu(\nu+1) - 20}{\nu(\nu+1) + 60}.$$

From (B), we see that for $6 \leq \nu < 7$, $\sqrt{11/51} \leq z_0 < \sqrt{9/29}$. On the other hand, if we set $\bar{\phi} = \arccos \sqrt{11/51}$, $m = 6$, $k = 1$ and $j_1^6 \doteq 9.9361$ in (2.12a), we find that for such a ϕ_0 and ν ,

$$z_0 = \cos \phi_0 = \cos(\phi_{\nu,1}^6) < \cos \left(\frac{j_1^6}{\sqrt{(7.5)^2 - 143/12}} \right) \doteq 0.0784,$$

which is a contradiction. Thus, we have shown that if $\nu = \nu_1^6 = \nu_2^3$, then $\nu \geq 7$. Moreover, we have also shown that if $\nu_1^6 = \nu_2^3$ for some ϕ_0 , then $\phi_0 \leq \arccos \sqrt{9/29}$.

Part 2. By the relationship in (B), we are motivated to define a function $z_0(\nu)$ for $\nu \geq 7$ as follows:

$$(3.1) \quad z_0(\nu) = \left(\frac{\nu(\nu+1) - 20}{\nu(\nu+1) + 60} \right)^{1/2}.$$

For such a $z_0(\nu)$, we also define $\phi_0(\nu) = \arccos(z_0(\nu))$ and observe that

$$(3.2) \quad \sin^2(\phi_0(\nu)) = \frac{80}{\nu(\nu+1) + 60}.$$

From (2.12a), (2.10), and (3.2), with $m = 6$, $k = 1$ and $\bar{\phi} = \arccos \sqrt{9/29}$, we see that for all $\nu \geq 7$,

$$\begin{aligned} (3.3) \quad \sin^2(\phi_{\nu,1}^6) &> \sin^2 \left(\frac{j_1^6}{\sqrt{(\nu + \frac{1}{2})^2 - \frac{h^2}{4}}} \right) \\ &\geq \frac{(j_1^6)^2}{\nu(\nu+1) + \frac{1}{4} - \frac{h^2}{4} + \alpha(\bar{\phi})(j_1^6)^2} \\ &> \frac{80}{\nu(\nu+1) + 60} \\ &= \sin^2(\phi_0(\nu)), \end{aligned}$$

where $j_1^6 \doteq 9.9361$. To complete the proof, we observe that if $\nu^* = \nu_1^6 = \nu_2^3$ for some $\nu^* \geq 7$, then necessarily we must have $\phi_{\nu^*,1}^6 = \phi_{\nu^*,2}^3 = \phi_0(\nu^*)$. However, from (3.3) we see that $\phi_0(\nu) < \phi_{\nu,1}^6$ for all $\nu \geq 7$. It follows that $\nu_1^6 \neq \nu_2^3$ for $0 < \phi_0 < \pi$. Since $\nu_2^3 < \nu_1^6$ for $\pi/2 \leq \phi_0 < \pi$, we conclude that $\nu_2^3 < \nu_1^6$ for all $0 < \phi_0 < \pi$ (or equivalently, for all $z_0 \in (-1, 1)$).

As a consequence of (1.1)–(1.5) and Theorem 1, we see that the first eleven ν -zeroes are

$$(3.4) \quad \nu_1^0 < \nu_1^1 < \nu_1^2 < \nu_2^0 < \nu_1^3 < \nu_2^1 < \nu_1^4 < \nu_2^2 < \nu_3^0 < \nu_1^5 < \nu_2^3, \quad 0 < z_0 < 1,$$

and that this ordering is unique. The inequalities $\nu_2^0 < \nu_1^3$, $\nu_2^1 < \nu_1^4$, and $\nu_3^0 < \nu_1^5$ were established in [2] for $0 \leq z_0 < 1$. By applying (2.8) and arguing as we did at the beginning of the proof of Theorem 1, these inequalities can be shown to hold for $-1 \leq z_0 < 0$ as well. In particular, we have the following:

Theorem 2. $\nu_2^0 < \nu_1^3$, $\nu_2^1 < \nu_1^4$, $\nu_3^0 < \nu_1^5$ for all $z_0 \in (-1, 1)$.

The inequality $\nu_j^{m+2} < \nu_{j+1}^m$ for $0 < z_0 < 1$ was established in [2]. Next, we consider the case $-1 < z_0 < 0$.

Lemma 2. If $-1 < z_0 < 0$, then $\nu_{j+1}^m < \nu_j^{m+2}$.

Proof. The ν_j^m 's are simple zeroes of $P_\nu^m(z_0)$. (See [2].) From (1.1), we have that $\nu_{j+1}^m, \nu_j^{m+2} \in (\nu_j^{m+1}, \nu_{j+1}^{m+1})$. Suppose that $\nu_j^{m+2}(z_0) \leq \nu_{j+1}^m(z_0)$ for some $z_0 \in (-1, 0)$. From (2.7)–(2.8), we see that

$$\text{sign}(P_\nu^m(z_0)) = (-1)^{m+j}, \quad \nu_j^m < \nu < \nu_{j+1}^m, \quad z_0 \in (-1, 0).$$

The signs of $P_\nu^{m+1}(z_0)$ and $P_\nu^{m+2}(z_0)$ can also be determined in this way. We are led to the results summarized in Table 1.

TABLE 1. Suppose $\nu_j^{m+2}(z_0) \leq \nu_{j+1}^m(z_0)$ for $z_0 \in (-1, 0)$

$z_0 \in (-1, 0)$	$\nu \in (\nu_j^{m+1}, \nu_j^{m+2})$	$\nu \in (\nu_j^{m+2}, \nu_{j+1}^m)$	$\nu \in (\nu_{j+1}^m, \nu_{j+1}^{m+1})$
$\text{sign}(P_\nu^m(z_0))$	$(-1)^{m+j}$	$(-1)^{m+j}$	$(-1)^{m+j+1}$
$\text{sign}(P_\nu^{m+1}(z_0))$	$(-1)^{m+j+1}$	$(-1)^{m+j+1}$	$(-1)^{m+j+1}$
$\text{sign}(P_\nu^{m+2}(z_0))$	$(-1)^{m+j+1}$	$(-1)^{m+j+2}$	$(-1)^{m+j+2}$

The second column in Table 1 contradicts (2.3), and we conclude that $\nu_j^{m+2} > \nu_{j+1}^m$ for all $z_0 \in (-1, 0)$.

From (1.1)–(1.3) and Lemma 2, we are led to the following:

Theorem 3. If $\nu_j^m(z_0)$ is a solution of (*), then

- (i) $\nu_j^{m+1} < \nu_{j+1}^m < \nu_j^{m+2} < \nu_{j+1}^{m+1}$, $-1 < z_0 < 0$,
- (ii) $\nu_j^{m+2} = \nu_{j+1}^m = m + 2j - 1$, $z_0 = 0$,
- (iii) $\nu_j^{m+1} < \nu_j^{m+2} < \nu_{j+1}^m < \nu_{j+1}^{m+1}$, $0 < z_0 < 1$.

4. CONCLUDING REMARKS

Since the zeroes of the Bessel functions are distinct [9, p. 484] the elements of $\mathcal{J} = \{j_k^m\}$ can be arranged as an increasing sequence. In particular, we can

define integer-valued functions $m(i)$, $k(i)$ so that $j_{k(i)}^{m(i)}$ denotes the i th element in the sequence $\mathcal{J}$. Clearly, there is no such ordering of all the elements of $\mathcal{N}_{\phi_0} = \{\nu_j^m(\phi_0)\}$ that is independent of ϕ_0 . On the other hand, if we let $\bar{\phi} = \phi_0 = \phi_{\nu,k}^m$ and $\nu = \nu_k^m(\phi_0)$ in (2.12), we see that

$$(4.1) \quad \lim_{\phi_0 \rightarrow 0^+} \phi_0 \left(\nu_k^m(\phi_0) + \frac{1}{2} \right) = j_k^m.$$

The limit in (4.1) is related to the well-known result, $\lim_{n \rightarrow \infty} \phi_{n,k}^0(n + \frac{1}{2}) = j_k^0$ (see [8]) and implies that for ϕ_0 sufficiently small, $\nu_{k(i)}^{m(i)}$ is the i th element in the sequence $\mathcal{N}_{\phi_0} = \{\nu_{j(i)}^{m(i)}(\phi_0)\}$.

In view of (1.6) and the first two inequalities in (1.4), it is natural to conjecture if there is an inequality that relates ν_{j+1}^m and ν_j^{m+3} for $\phi_0 \in (0, \pi/2)$. Such an inequality is not possible. From [1], we see that $j_{k(18)}^{m(18)} = j_1^8 = 12.225$, and $j_{k(19)}^{m(19)} = j_2^5 = 12.338$. Since $\nu_1^8(\pi/2) = 9$, $\nu_2^5(\pi/2) = 8$, and $j_1^8 < j_2^5$, we conclude that $\nu_1^8(\phi_0) = \nu_2^5(\phi_0)$ for some $\phi_0 \in (0, \pi/2)$. Numerical calculations indicate that $\nu_1^8 = \nu_2^5 \doteq 26.706$ when $\phi_0 \doteq 26.134^\circ$.

Although (1.5) and Theorem 1 demonstrate that the first eleven ν -zeroes of $P_\nu^m(\cos \phi_0)$ are distinct for $0 < \phi_0 < \pi/2$, the twelfth ν -zero is not necessarily distinct. Since $\nu_3^1(\pi/2) = 6$, $\nu_1^6(\pi/2) = 7$, and $j_{k(12)}^{m(12)} = j_1^6 = 9.936$, and $j_{k(13)}^{m(13)} = j_3^1 = 10.173$, from (4.1), we see that $\nu_1^6(\phi_0) = \nu_3^1(\phi_0)$ for some $\phi_0 \in (0, \pi/2)$. Numerics indicate $\nu_1^6 = \nu_3^1 \doteq 15.780$ when $\phi_0 \doteq 35.821^\circ$ (see [2]).

REFERENCES

1. M. Abramowitz and I. A. Stegun, *Handbook of mathematical functions. Applied mathematics series 55*, National Bureau of Standards, Washington, DC, 1964.
2. F. Baginski, *Ordering the zeroes of the Legendre functions $P_\nu^m(z_0)$ when considered as a function of ν* , J. Math. Anal. Appl. **147** (1990), 296–308.
3. A. Erdélyi, W. Magnus, and F. Tricomi, *Higher transcendental functions*, vol. 1, McGraw-Hill, New York, 1953.
4. E. Hobson, *The theory of spherical and ellipsoidal harmonics*, Cambridge Univ. Press, London, 1931.
5. A. Laforgia and M. E. Muldoon, *Some consequences of the Sturm Comparison Theorem*, Amer. Math. Monthly **93** (1986), 89–94.
6. G. Sansone, *Orthogonal functions*, Interscience Publishers, New York, 1959.
7. G. Szegő, *Orthogonal polynomials*, Amer. Math. Soc. Colloq. Publ., vol. 23, New York, 1939.
8. —, *Inequalities for the zeroes of Legendre polynomials and related functions*, Trans. Amer. Math. Soc. **39** (1936), 1–17.
9. G. N. Watson, *Theory of Bessel functions*, Cambridge Univ. Press, Cambridge, 1922.
10. H. F. Weinberger, *A first course in partial differential equations with complex variables and transform methods*, Wiley, New York, 1965.