LOWER BOUNDS FOR THE SOLUTIONS IN THE SECOND CASE OF FERMAT'S LAST THEOREM

LE MAOHUA

(Communicated by William Adams)

ABSTRACT. Let p be an odd prime. In this paper, we prove that if $p \equiv 3 \pmod{4}$ and x, y, z are integers satisfying $x^p + y^p = z^p$, $p \mid xyz$, $0 < x < y < z$, then $y > 2^{-1/p} p^{6p-2}$ and $z - x > \frac{1}{2} p^{6p-3}$.

Let p be an odd prime. In [1] and [2], Inkeri showed that if x, y, z are integers satisfying

$$(1) \quad x^p + y^p = z^p, \quad \gcd(x, y) = 1, \quad p \mid xyz, \quad 0 < x < y < z,$$

then $y > \frac{1}{2} p^{3p-1}$ and $z - x > (2p^{20/7})^p$. In this paper, we prove the following result:

Theorem. If $p \equiv 3 \pmod{4}$ and x, y, z are integers satisfying (1), then $y > 2^{-1/p} p^{6p-2}$ and $z - x > \frac{1}{2} p^{6p-3}$.

Proof. It is a well-known fact that (1) is impossible for $p = 3$, so we may assume that $p > 3$. We first deal with the case that $p \mid z$. Let ζ be a p th primitive root of 1. Let S be the set of nonzero squares mod p , and let $\theta = \prod_{i \in S} (x + y\zeta^i)$. Since S represents $\text{Gal}(\mathbb{Q}(\zeta)/\mathbb{Q}(\sqrt{-p}))$, it follows that θ is an algebraic integer in $\mathbb{Q}(\sqrt{-p})$, so $\theta = A + B\sqrt{-p}$, where $A, B \in \frac{1}{2}\mathbb{Z}$. Note that $\theta_0 = \prod_{i \in S} (1 - \zeta^2)$ is an algebraic integer in $\mathbb{Q}(\sqrt{-p})$ with norm p ; hence, it must be $\pm\sqrt{-p}$. From

$$\theta \equiv \begin{cases} \theta_0 = \pm\sqrt{-p} \pmod{2}, & 2 \nmid x, \ 2 \nmid y, \\ \prod_{i \in S} 1 = 1 \pmod{2}, & 2 \nmid x, \ 2 \mid y, \\ \prod_{i \in S} \zeta^i = 1 \pmod{2}, & 2 \mid x, \ 2 \nmid y, \end{cases}$$

we find that $A + B\sqrt{-p}$ is congruent modulo 2 to an element of $\mathbb{Z}(\sqrt{-p})$, so $A, B \in \mathbb{Z}$. Further, since

$$\theta \equiv \prod_{i \in S} (x - x\zeta^i) = 0 \pm x^{(p-1)/2} \sqrt{-p} \pmod{x + y},$$

Received by the editors November 11, 1989 and, in revised form, March 27, 1990.
1980 *Mathematics Subject Classification* (1985 Revision). Primary 11D41.

©1991 American Mathematical Society
0002-9939/91 \$1.00 + \$.25 per page

we get

$$(2) \quad A \equiv 0 \pmod{x+y}.$$

Let $p^\alpha \parallel z$. Classical relations due to Abel [4, p. 54] imply that

$$(3) \quad x+y = p^{\alpha p-1} a^p, \quad \frac{x^p + y^p}{x+y} = pb^p, \quad z = p^\alpha ab,$$

where a, b are positive integers satisfying $p \nmid ab$ and $2 \nmid b$. From the above, we have $pb^p = A^2 + pB^2$. Hence $p \mid A$, $p \nmid B$, and

$$(4) \quad b^p = pA'^2 + B^2,$$

where $A' = A/p$. By (2) and (3), we get

$$(5) \quad A' \equiv 0 \pmod{p^{\alpha p-2}}.$$

The factors $x+y\zeta^i$ are pairwise relatively prime for $1 \leq i \leq p-1$, except for factors of $1-\zeta$. Therefore $\prod_{i \in S}(x+y\zeta^i) = A+B\sqrt{-p}$ and $\prod_{i \notin S}(x+y\zeta^i) = A-B\sqrt{-p}$ are relatively prime, except for primes above p . Recall that $p \nmid B$. We have $\gcd(A, B) = 1$. It also follows that $A'\sqrt{-p} + B$ and $A'\sqrt{-p} - B$ are relatively prime in $\mathbb{Z}[\rho]$, where $\rho = (-1 + \sqrt{-p})/2$. Hence they are p th powers of ideals. Since the class number of $\mathbb{Z}[\rho]$ is less than p , it is prime to p . Therefore $A'\sqrt{-p} + B$ is the p th power of a number in $\mathbb{Z}[\rho]$. It implies that

$$(6) \quad A'\sqrt{-p} + B = (X_1 + Y_1\sqrt{-p})^p,$$

where $X_1, Y_1 \in \frac{1}{2}\mathbb{Z}$. If $X_1 + Y_1\sqrt{-p} \notin \mathbb{Z}(\sqrt{-p})$, then $X_1 = u/2$, $Y_1 = v/2$, with $u, v \in \mathbb{Z}$ and $u \equiv v \equiv 1 \pmod{2}$. By Waring's formula [3, Formula 1.76], we get from (4) and (6) that $u^2 + pv^2 = 4b$ and

$$\begin{aligned} 2B &= \left(\frac{u+v\sqrt{-p}}{2}\right)^p + \left(\frac{u-v\sqrt{-p}}{2}\right)^p \\ &= \sum_{j=0}^{(p-1)/2} (-1)^j \frac{(p-j-1)!p}{(p-2j)!j!} u^{p-2j} v^j \equiv \sum_{j=0}^{(p-1)/2} (-1)^j \frac{(p-j-1)!p}{(p-2j)!j!} \\ &= \left(\frac{1+\sqrt{5}}{2}\right)^p + \left(\frac{1-\sqrt{5}}{2}\right)^p = L_p \pmod{2}, \end{aligned}$$

where L_p is the p th Lucas number. Since $L_p \equiv 0 \pmod{2}$ only if $p = 3$, it follows that $X_1 + Y_1\sqrt{-p} \in \mathbb{Z}[\sqrt{-p}]$, and

$$(7) \quad X_1^2 + pY_1^2 = b, \quad X_1, Y_1 \in \mathbb{Z}.$$

We get from (6) that

$$(8) \quad A' = \binom{p}{1} X_1^{p-1} Y_1 - \binom{p}{3} p X_1^{p-3} Y_1^3 + \cdots + (-1)^{(p-1)/2} \binom{p}{p} p^{(p-1)/2} Y_1^p.$$

Since $p \nmid b$, we have $p \nmid X_1$ by (7), and hence

$$(9) \quad Y_1 \equiv 0 \pmod{p^{\alpha p-3}}$$

by (5) and (8). If $Y_1 = 0$, then $A = 0$, $B = \pm 1$, and $b = 1$ by (4). Since $(x^p + y^p)/(x + y) \geq 2^{p-2} + 1$, it is impossible that $p > 3$. Therefore $Y_1 \neq 0$ and $|Y_1| > p^{\alpha p-3}$ by (9), and $z = p^\alpha ab > p^{2\alpha p+\alpha-5}$ by (7). Note that $\alpha \geq 3$, by [5]. We obtain $z > p^{6p-2}$. Using the same method, we can prove that $y > p^{6p-2}$ or $x > p^{6p-2}$ correspond to $p|y$ or $p|x$. Thus $y > 2^{-1/p} p^{6p-2}$ since $2^{1/p} y > z$. Simultaneously, we have

$$\begin{aligned} z - x &= \frac{y^p}{z^{p-1} + xz^{p-2} + \dots + x^{p-1}} > \frac{y^p}{pz^{p-1}} > \frac{y^p}{p(2^{1/p}y)^{p-1}} \\ &= \frac{y}{2^{p-1/p}p} > \frac{1}{2} p^{6p-3}. \end{aligned}$$

The theorem is proved.

ACKNOWLEDGMENT

The author would like to thank the referee for his valuable suggestions.

REFERENCES

1. K. Inkeri, *Abschätzungen für eventuelle Lösungen der Gleichung im Fermatschen Problem*, Ann. Univ. Turku, 1953, 3-9.
2. —, *Remarks on Fermat's equation: The very knowledge of coding*, Univ. Turku, Turku, 1987, pp. 82-87.
3. R. Lidl and H. Niederreiter, *Finite fields*, Addison-Wesley, Reading, MA, 1983.
4. P. Ribenboim, *13 lectures on Fermat's last theorem*, Springer-Verlag, New York, 1979.
5. H. Vandiver, *A property of cyclotomic integers and its relation to Fermat's last theorem*, Ann. of Math. **21** (1919), 73-80.

RESEARCH DEPARTMENT, CHANGSHA RAILWAY INSTITUTE, CHANGSHA, HUNAN, PEOPLE'S REPUBLIC OF CHINA