STABILITY IN INTERPOLATION OF FAMILIES OF BANACH SPACES

WEI CAO AND YORAM SAGHER

(Communicated by J. Marshall Ash)

ABSTRACT. Let D be a simply connected domain in the complex plane whose boundary Γ is a rectifiable simple closed curve. Let $\{A(\gamma)/\gamma \in \Gamma\}$ and $\{B(\gamma)/\gamma \in \Gamma\}$ be interpolation families of Banach spaces. Let T be a linear operator mapping $A(\gamma)$ continuously into $B(\gamma)$. For $z \in D$ let T_z be the restriction of T to the interpolation space A_z . Then $\{z \in D/\text{cod}(T_z) = d < \infty \text{ and } \dim \text{Ker}(T_z) = 0\}$ and $\{z \in D/\dim \text{Ker}(T_z) = d < \infty \text{ and } T_z \text{ is onto } B_z\}$ are open sets.

1. INTRODUCTION

We begin with a brief review of the theory of interpolation for analytic families of Banach spaces. Throughout the paper, D will denote a simply connected domain in the complex plane whose boundary Γ is a rectifiable simple closed curve.

Definition 1.1. An interpolation family of Banach spaces on Γ is a collection $\{A(\gamma)/\gamma \in \Gamma\}$, of Banach spaces which satisfies:

- (1) Every $A(\gamma)$ is continuously embedded in a common Banach space $\mathcal{U}$ which is called a containing space for $\{A(\gamma)/\gamma \in \Gamma\}$.
- (2) For every $a \in \cap_{\gamma \in \Gamma} A(\gamma)$, $\|a\|_{A(\gamma)}$ is a measurable function on Γ with respect to dP_z , where $z \in D$ and dP_z is a harmonic measure on Γ with respect to z .
- (3) Let $\mathcal{A} = \{a \in \cap_{\gamma \in \Gamma} A(\gamma) | \int_{\Gamma} \log^+ \|a\|_{A(\gamma)} dP_z(\gamma) < \infty\}$, where $\log^+ x = \max(\log x, 0)$. $\mathcal{A}$ is a linear space which is called the log-intersection space for $\{A(\gamma)/\gamma \in \Gamma\}$. We assume that $\mathcal{A}$ is dense in each $A(\gamma)$ and that there exists a measurable function ψ on Γ which satisfies $\int_{\Gamma} \log^+ \psi(\gamma) dP_z < \infty$ such that $\|a\|_u \leq \psi(\gamma) \|a\|_{A(\gamma)}$ for all $a \in \mathcal{A}$ and $\gamma \in \Gamma$.

We shall denote the spaces $A(\gamma)$ by A_γ and their norms by $\|\cdot\|_\gamma$.

Received by the editors October 4, 1989 and, in revised form, February 2, 1990.
 1980 *Mathematics Subject Classification* (1985 Revision). Primary 47A53.

©1991 American Mathematical Society
 0002-9939/91 \$1.00 + \$.25 per page

We denote by $N^+(D)$ the class of all functions $f(z)$ analytic on D such that $f(\phi(w))$ belongs to the class $N^+ = N^+(\Delta)$, where $\Delta = \{w \mid |w| < 1\}$ and ϕ is a conformal map from D to Δ , see [7]. Thus $N^+(D)$ is closed under pointwise addition and multiplication and each $f(z) \in N^+(D)$ possesses a.e. non-tangential limits on Γ . If these non-tangential limits are essentially bounded on Γ , then $f \in H^\infty(D)$, the space of bounded analytic functions on D . A function $f \in N^+(D)$ is termed an outer function in $N^+(D)$, if $f(\phi(w))$ is an outer function in $N^+(\Delta)$.

Let $\mathcal{G}(A(\cdot), \Gamma) = \mathcal{G}(\mathcal{A}) = \mathcal{G}$ be the space of all functions of the form $g(z) = \sum_{j=1}^n \varphi_j(z) a_j$, where $a_j \in \mathcal{A}$ and $\varphi_j \in N^+(D)$, and such that

$$\|g(\cdot)\|_{\mathcal{G}} = \text{ess sup}_{\gamma \in \Gamma} \|g(\gamma)\|_{\gamma} < \infty.$$

Note that, in general, $(\mathcal{G}, \|\cdot\|_{\mathcal{G}})$ is not complete.

Definition 1.2. For $z \in D$ and $a \in \mathcal{A}$, define

$$\|a\|_{A\{z\}} = \inf\{\|g\|_{\mathcal{G}} \mid g \in \mathcal{G}, g(z) = a\}.$$

The interpolation space $\{A_{\gamma}\}\{z\} = A\{z\}$ is defined to be the completion in $\mathcal{U}$ of $(\mathcal{A}, \|\cdot\|_{A\{z\}})$.

In most applications the family of dual spaces $\{A_{\gamma}^*\}$ is itself an interpolation family and $(A\{z\})^* = \{A_{\gamma}^*\}\{z\}$. In this paper we shall assume that the above duality result holds.

We consider in this paper linear operators T mapping $\mathcal{A}$ into $\cap_{\gamma \in \Gamma} B(\gamma)$, with $\|Ta\|_{B(\gamma)} \leq M(\gamma)\|a\|_{A(\gamma)}$ for all $a \in \mathcal{A}$ and $\gamma \in \Gamma$, where $\{A(\gamma)\}_{\gamma \in \Gamma}$ and $\{B(\gamma)\}_{\gamma \in \Gamma}$ are interpolation families on Γ and

$$\int_{\Gamma} |\log M(\gamma)| dP_z(\gamma) < \infty.$$

We will denote the restriction of T to $A\{z\}$ by T_z . If $\ker T_z = \{0\}$, denote by T_z^{-1} the inverse of T_z , i.e., T_z^{-1} is defined on $T_z A\{z\}$. If $T_z A\{z\}$ has finite codimension, by the open mapping theorem, $T_z A\{z\}$ is closed and T_z^{-1} is a bounded linear operator.

The question of the stability of Fredholm property when one changes the parameters which determine the interpolation space has been considered by several authors [1, 4, 9, 12] in the context of interpolation of two spaces and by Vignati and Vignati [11] in the case of families of Banach spaces. We consider families of Banach spaces and prove that, if $\ker T_s = \{0\}$ and $\text{codim } T_s = d < \infty$, then there exists a ball B centered at s such that $z \in B$ implies that $\ker T_z = \{0\}$ and $\text{codim } T_z = d$. This generalizes the result in [11], where the theorem is proved in the case $d = 0$.

Definition 1.3. Suppose $U(A)$ is a subset of $\mathcal{G}(\mathcal{A})$. Let

$$U_s = \{a \in A\{s\} \mid \exists f \in U(A) \text{ such that } f(s) = a\}.$$

Define

$$\|a\|_{U_s(A)} = \inf\{\|f\|_{\mathcal{G}} \mid f(s) = a, f \in U(A)\}.$$

Clearly $\|a\|_{A\{s\}} \leq \|a\|_{U_s(A)}$ for all $a \in U_s$.

Let $U_{\{s\}}$ be a completion in $A\{s\}$ of $\{U_s, \|\cdot\|_{U_s}\}$. Clearly $U_{\{s\}} \subset A\{s\}$. If also $\|a\|_{U_s(A)} < k\|a\|_{A\{s\}}$, for some k and all $a \in U_s$ with $a \neq 0$, we call $U(A)$ an s -Calderón subset (with constant k). If $U(A)$ is an s -Calderón subset and a linear subspace of $\mathcal{G}(\mathcal{A})$, we will say that it is an s -Calderón subspace.

Clearly $U(A)$ is an s -Calderón subspace for some k if and only if $U_{\{s\}}$ is a closed subspace of $A\{s\}$.

An important tool in handling finite dimensional subspaces is Auerbach's lemma:

Lemma 1.4. *Let M be a Banach space with dimension $d < \infty$. Then there exist $\{e_1, \dots, e_d\} \subset M$ and $\{f_1, \dots, f_d\} \subset M^*$ such that*

- (1) $\|e_i\|_M = 1$ and $\|f_i\|_{M^*} = 1$, $i = 1, \dots, d$.
- (2) $f_i(e_j) = \delta_{ij}$.

For a proof see [8].

Given a finite set of vectors $\{e_1, \dots, e_d\}$ in a Banach space A , we define

$$b(e_1, \dots, e_d) = \min \left\{ \left\| \sum_{i=1}^d c_i e_i \right\|_A \mid \max |c_i| = 1 \right\}.$$

Clearly $\{e_1, \dots, e_d\}$ are independent if and only if $b(e_1, \dots, e_d) > 0$. Given $\sum_{i=1}^d c_i e_i$, let $m = \max\{|c_i|\}$. Then $\|\sum_{i=1}^d c_i e_i\|_A = m \|\sum_{i=1}^d c_i e_i / m\|_A \geq mb(e_1, \dots, e_d)$, so that $|c_i| \leq \|\sum_{i=1}^d c_i e_i\|_A / b(e_1, \dots, e_d)$.

Lemma 1.5. *Let M be a Banach space with dimension $d < \infty$. Then $\{e_1, \dots, e_d\}$ can be found so that $\|e_i\|_M = 1$, and $b(e_1, \dots, e_d) = 1$.*

For a proof, see [4].

Lemma 1.6. *Let E be a closed subspace of $A\{s\}$ with finite codimension d . Then there exists $\{v_i, i = 1, \dots, d\} \subset \mathcal{G}(\mathcal{A})$, such that $A\{s\} = E \oplus \text{span}\{v_i(s)\}$.*

Proof. Since E has finite codimension d , $A\{s\} = E \oplus M$ with $\dim M = d$. Assume that $\{e_i, 1 \leq i \leq d\}$ forms a basis for M with $\|e_i\|_{A\{s\}} = 1$ and $b(e_1, \dots, e_d) = 1$. Let

$$\rho = \rho(e_1, \dots, e_d; E) = \inf \left\{ \left\| \sum_{i=1}^d c_i e_i - x \right\|_{A\{s\}} \mid \max |c_i| = 1, x \in E \right\}.$$

Clearly $\rho \leq 1$. If $\rho > 0$ then $E \cap M = \{0\}$ and $\{e_i\}$ are independent. Since E is a closed subspace and M has finite dimension, then $E \cap M = \{0\}$ implies $\rho > 0$.

Since $\{v(s), v \in \mathcal{G}(\mathcal{A})\}$ is dense in $A\{s\}$, we can find $\{v_i, i = 1, \dots, d\} \subset \mathcal{G}(\mathcal{A})$ with $\|v_i(s) - e_i\|_{A\{s\}} < \rho/2d$. Now suppose $\max\{|c_i|\} = 1$ and let $x \in E$; we have

$$\left\|x - \sum_{i=1}^d c_i v_i(s)\right\|_{A\{s\}} \geq \left\|x - \sum_{i=1}^d c_i e_i\right\|_{A\{s\}} - \left\|\sum_{i=1}^d c_i (v_i(s) - e_i)\right\|_{A\{s\}} > \rho/2.$$

Taking infimum we get $\rho(v_1(s), \dots, v_d(s); E) \geq \rho/2 > 0$. Hence $\{v_i(s)\}$ are linearly independent and $A\{s\} = E \oplus \text{span}\{v_i(s)\}$. $\square$

The function ρ introduced in the last proof is, of course, a generalization of the function $b(e_1, \dots, e_d)$. Another application for the function ρ is the following lemma.

Lemma 1.7. *Let E be a closed subspace of a Banach space B , and let M be a finite dimensional subspace of B . Then $E + M$ is closed.*

Proof. We may assume $E \cap M = \{0\}$. Let $\{e_1, \dots, e_d\}$ be a basis for M . Then $\rho = \rho(e_1, \dots, e_d; E) > 0$. Let $x_n \in E + M$, and $x_n \rightarrow x_0 \in B$. $x_n = y_n + \sum_{i=1}^d c_i^n e_i$. Let $M_n = \max\{|c_i^n| \mid i = 1, \dots, d\}$.

Given $\varepsilon > 0$, let N be so that, for all $n \geq N$, $\|x_n\| \leq (1 + \varepsilon)\|x_0\|$. Then $\rho \leq \|\sum_{i=1}^d c_i^n e_i + y_n\|/M_n \leq (1 + \varepsilon)\|x_0\|/M_n$, and $M_n \leq (1 + \varepsilon)\|x_0\|/\rho$. Thus, $\limsup M_n \leq \|x_0\|/\rho$ and $\{\sum_{i=1}^d c_i^n e_i\}$ is compact. Denote $\sum_{i=1}^d c_i^n e_i = z_n$. Let $z_{n_k} \rightarrow z \in M$; then $y_{n_k} = x_{n_k} - z_{n_k} \rightarrow x_0 - z$ and, since E is closed, $x_0 - z \in E$. $\square$

We will also need the following well-known result due to Szegő [10].

Lemma 1.8. *Let $f(\gamma)$ be a positive dP_z measurable function on Γ such that*

$$\int_{\Gamma} |\log f(\gamma)| dP_z(\gamma) < \infty$$

for some (and thus every) $z \in D$. Then there exists a non-vanishing outer function $G(z)$ in $N^+(D)$ whose a.e. non-tangential limits $G(\gamma) = \lim_{z \rightarrow \gamma} G(z)$ satisfy $|G(\gamma)|f(\gamma) = 1$ for a.e. $\gamma \in \Gamma$.

2. STABILITY

An invariant form of Schwarz's lemma states that if $f(z)$ is analytic and $|f(z)| < 1$ on Δ , then for any $z, \zeta \in \Delta$,

$$\left| \frac{f(z) - f(\zeta)}{1 - \overline{f(z)}f(\zeta)} \right| \leq \left| \frac{z - \zeta}{1 - \overline{z}\zeta} \right|.$$

This implies that if $z_0, \zeta_0, z \in \Delta$ and if we denote

$$m_{z_0}(z) = \frac{z - z_0}{1 - \overline{z_0}z} e^{i\alpha},$$

then

$$\left| \frac{m_{z_0}(z) - m_{z_0}(\zeta)}{1 - \overline{m_{z_0}(z)}m_{z_0}(\zeta)} \right| = \left| \frac{z - \zeta}{1 - \overline{z}\zeta} \right|.$$

This, in turn, implies that if $z, \zeta \in D$ and d is a conformal map of D onto Δ , then

$$q(z, \zeta) = \left| \frac{d(z) - d(\zeta)}{1 - \overline{d(z)}d(\zeta)} \right|$$

does not depend on d .

The invariant form of Schwarz's lemma also implies the following lemma.

Lemma 2.1. *Let ϕ be analytic on D , $|\phi(z)| \leq M$. If $s, z \in D$, then*

$$|\phi(z)| \geq M \frac{|\phi(s)| - Mq(z, s)}{M - |\phi(s)|q(z, s)}.$$

For a proof see [9]. Of course, since $q(z, s) < 1$, the inequality above implies $|\phi(z)| \geq |\phi(s)| - q(z, s)M$.

Lemma 2.2. *If $f \in \mathcal{G}(\mathcal{A})$ and $z, s \in D$, then*

$$(2.1) \quad \|f(z)\|_{A\{z\}} \geq \|f(s)\|_{A\{s\}} - q(z, s)\|f\|_{\mathcal{G}}.$$

Proof. Let us fix $s \in D$. Since $f(s) \in A\{s\}$, it is possible to find $\psi \in A^*\{s\}$ such that $\|\psi\|_{A^*\{s\}} = 1$ and $\psi(f(s)) = \|f(s)\|_{A\{s\}}$.

Given $\varepsilon > 0$, we can find $G \in \mathcal{G}(A^*)$ such that $\|G(s) - \psi\|_{A^*\{s\}} < \varepsilon$ and $\|G\|_{\mathcal{G}(A^*)} < 1 + \varepsilon$. Thus $\phi(z) = G(z)(f(z))$ is analytic in D , $\phi \in H^\infty(D)$, and

$$|\phi(z)| \leq (1 + \varepsilon)\|f(z)\|_{A\{z\}} \leq (1 + \varepsilon)\|f\|_{\mathcal{G}}.$$

By Lemma 2.1, we have

$$\begin{aligned} \|f(z)\|_{A\{z\}} &\geq \frac{|\phi(z)|}{1 + \varepsilon} \geq \frac{|\phi(s)|}{1 + \varepsilon} - q(z, s)\|f\|_{\mathcal{G}} \\ &\geq \frac{1 - \varepsilon}{1 + \varepsilon}\|f(s)\|_{A\{s\}} - q(z, s)\|f\|_{\mathcal{G}}. \end{aligned}$$

Letting $\varepsilon \rightarrow 0$, we get the theorem. $\square$

Theorem 2.3. *Let $\{v_1, \dots, v_d\} \subset \mathcal{G}(\mathcal{A})$, and assume that $\{v_i(s)\}$ are independent. Denote $b_z = b(v_1(z), \dots, v_d(z))$. Then there exists $\delta > 0$ so that, for all $|z - s| < \delta$, $b_z > b_s/2$. Consequently, if we denote by U the span of $\{v_i\}$ in $\mathcal{G}(\mathcal{A})$ then, for these z , $\dim(U_{\{z\}}) = d$.*

Proof. Without loss of generality, we can assume that $\|v_i\|_{\mathcal{G}} = 1$. As we consider $v_i(z)$ in $A\{z\}$, b_z will be controlled by inequality (2.1), which will prove the theorem:

$$\left\| \sum_{i=1}^d c_i v_i(z) \right\|_{A\{z\}} \geq \left\| \sum_{i=1}^d c_i v_i(s) \right\|_{A\{s\}} - q(z, s) \left\| \sum_{i=1}^d c_i v_i \right\|_{\mathcal{G}}.$$

Since $\{v_i(s)\}$ are independent, we have $b_s > 0$. Suppose $\max\{|c_i|\} = 1$; then

$$b_s \leq \left\| \sum_{i=1}^d c_i v_i(s) \right\|_{A\{s\}} \quad \text{and} \quad \left\| \sum_{i=1}^d c_i v_i \right\|_{\mathcal{F}} \leq \sum_{i=1}^d |c_i| \cdot \|v_i\|_{\mathcal{F}} \leq d.$$

Hence,

$$\left\| \sum_{i=1}^d c_i v_i(z) \right\|_{A\{z\}} \geq b_s - dq(z, s).$$

We can choose δ small enough such that $|z - s| < \delta$ implies

$$b_s - dq(z, s) > b_s/2.$$

It follows that $b_z > b_s/2$. Therefore, if $|z - s| < \delta$, $\{v_i(z)\}$ are independent and $\dim(U_{\{z\}}) = d$. $\square$

Theorem 2.4. *Let $\{v_1, \dots, v_d\} \subset \mathcal{F}(\mathcal{A})$, and assume that $\{v_i(s)\}$ are independent. Denote by U the span of $\{v_i\}$ in $\mathcal{F}(\mathcal{A})$. Then $\exists \delta > 0$, such that $|z - s| < \delta$ implies U is a z -Calderón space with a uniform constant k .*

Proof. Without loss of generality, we can assume that $\|v_i(s)\|_{A\{s\}} = 1$. Let $m = \max\{\|v_i\|_{\mathcal{F}}\}$ and $b_z = b(v_1(z), \dots, v_d(z))$. Then, as we have seen before, if $b_z > 0$, we have

$$|c_i| \leq \frac{\left\| \sum_{i=1}^d c_i v_i(z) \right\|_{A\{z\}}}{b_z}$$

so that

$$\left\| \sum_{i=1}^d c_i v_i \right\|_{\mathcal{F}} \leq \frac{md}{b_z} \left\| \sum_{i=1}^d c_i v_i(z) \right\|_{A\{z\}}.$$

Since $b_s > 0$, by Theorem 2.3, $\exists \delta > 0$ such that $|z - s| < \delta$ implies that $b_z > b_s/2$ and U is a z -Calderón subspace with constant $2md/b_s$. $\square$

Theorem 2.5. *Let $U(A), V(A)$ be two z -Calderón subspaces with constant k , for all z in a ball S which is contained in D . Then the function*

$$\chi_z = \inf\{d(f(z), U_{\{z\}}) | f \in V(A) \text{ and } \|f(z)\|_{A\{z\}} = 1\}$$

is continuous in S .

Proof. Given $0 < \varepsilon < 1$ and $s \in S$, there are $v \in V(A)$, $u \in U(A)$ so that

$$\begin{aligned} 1 &= \|v(s)\|_{A\{s\}} \leq \|v\|_{\mathcal{F}} \leq k\|v(s)\|_{A\{s\}} = k \\ \|u(s) - v(s)\|_{A\{s\}} &< \chi_s + \varepsilon/2 \end{aligned}$$

with

$$\|u\|_{\mathcal{F}} \leq k\|u(s)\|_{A\{s\}}.$$

Therefore

$$\|u\|_{\mathcal{F}} \leq k(\|u(s) - v(s)\|_{A\{s\}} + \|v(s)\|_{A\{s\}}) < k(\chi_s + 2).$$

Clearly we have

$$\chi_z \leq \frac{\|u(z) - v(z)\|_{A\{z\}}}{\|v(z)\|_{A\{z\}}}.$$

By Lemma 2.2, we have

$$\|v(z)\|_{A\{z\}} \geq \|v(s)\|_{A\{s\}} - q(z, s)\|v\|_{\mathcal{G}} = 1 - q(z, s)\|v\|_{\mathcal{G}}.$$

Hence we can choose $\eta > 0$ small enough such that $|z - s| < \eta$ implies both $\|v(z)\|_z^{-1} < 1 + \varepsilon$ and $z \in S$. Again, invoking Lemma 2.2, we have

$$\begin{aligned} \|u(z) - v(z)\|_{A\{z\}} &\leq \|u(s) - v(s)\|_{A\{s\}} + q(z, s)\|u - v\|_{\mathcal{G}} \\ &\leq \|u(s) - v(s)\|_{A\{s\}} + q(z, s)(\|u\|_{\mathcal{G}} + \|v\|_{\mathcal{G}}) \\ &< (\chi_s + \varepsilon/2) + q(z, s)(\chi_s + 3)k. \end{aligned}$$

Therefore, if necessary, choosing a smaller $\eta > 0$, we have that, for $|z - s| < \eta$,

$$\chi_z \leq \frac{\|u(z) - v(z)\|_{A\{z\}}}{\|v(z)\|_{A\{z\}}} \leq (1 + \varepsilon)(\chi_s + \varepsilon).$$

Similarly $\chi_s \leq (1 + \varepsilon)(\chi_z + \varepsilon)$ in $|z - s| < \eta_1$, and hence χ_z is continuous in S . $\square$

Note that when $V(A)$ is finite dimensional, as it is in the proof below, χ_z is equivalent to the function ρ defined in the proof of Lemma 1.6.

Theorem 2.6. *Let U be a z -Calderón subspace with constant k , for all z in a ball S contained in D . Assume that, for an $s \in S$, $U_{\{s\}}$ has codimension $d < \infty$. Then there exists $\delta > 0$, so that, for all z with $|z - s| < \delta$, we have $\text{codim } U_{\{z\}} = d$.*

Proof. By Lemma 1.6, there exist $\{v_i | i = 1, \dots, d\} \subset \mathcal{G}(\mathcal{A})$ so that $\{v_i(s)\}$ forms a basis for M_s and $A\{s\} = U_{\{s\}} \oplus M_s$. We can assume $\|v_i\|_{\mathcal{G}} = 1$. Denote by M_z the space spanned by $\{v_i(z)\}$. Since $A\{s\} = U_{\{s\}} \oplus M_s$ and $U_{\{s\}}, M_s$ are closed, there exists a projection P of $A\{s\}$ onto $U_{\{s\}}$ with $\ker P = M_s$, and $\|P\| < \infty$.

Since U is a z -Calderón subspace, $U_{\{z\}}$ is a closed subspace of $A\{z\}$. By Lemma 1.7, since M_z is finite-dimensional, $U_{\{z\}} + M_z$ is closed. Hence, if $A\{z\} \neq U_{\{z\}} + M_z$ then there exists $e \in \mathcal{G}(\mathcal{A})$, with $\|e(z)\|_{A\{z\}} = 1$ and $1 \leq \|e\|_{\mathcal{G}(\mathcal{A})} < 2$, such that $\text{dist}(e(z), U_{\{z\}} + M_z) > 2/3$.

Consider $e(s) \in A\{s\}$. We have $e(s) = y_s + z_s$, where $y_s = Pe(s) \in U_{\{s\}}$, $z_s = (I - P)e(s) = \sum_{i=1}^d c_i v_i(s) \in M_s$. We have

$$\begin{aligned} \|y_s\|_{A\{s\}} &\leq \|P\| \|e(s)\|_{A\{s\}} < 2\|P\| \\ \|z_s\|_{A\{s\}} &\leq (\|P\| + 1)\|e(s)\|_{A\{s\}} < 2(\|P\| + 1). \end{aligned}$$

Since U is s -Calderón, there exists $f \in U(A)$ such that $\|f(s) - y_s\|_{A\{s\}} \leq \min\{1/3, \|y_s\|_{A\{s\}}\}$ and

$$(2.2) \quad \|f\|_{\mathcal{G}} \leq k\|f(s)\|_{A\{s\}} \leq 2k\|y_s\|_{A\{s\}} < 4k\|P\|.$$

On the other hand we have, denoting $b = b(v_1(s), \dots, v_d(s))$,

$$|c_i| \leq \frac{\|\sum_{i=1}^d c_i v_i(s)\|_{A\{s\}}}{b} = \frac{\|z_s\|_{A\{s\}}}{b} < 2 \frac{\|P\| + 1}{b}.$$

It follows that

$$(2.3) \quad \left\| \sum_{i=1}^d c_i v_i \right\|_{\mathcal{G}} \leq \sum_{i=1}^d |c_i| \|v_i\|_{\mathcal{G}} \leq \frac{2d(\|P\| + 1)}{b}.$$

Let $g = f + \sum_{i=1}^d c_i v_i$. Clearly $\|g(s) - e(s)\|_{A\{s\}} < 1/3$ and

$$\begin{aligned} \|g - e\|_{\mathcal{G}} &\leq \|f\|_{\mathcal{G}} + \left\| \sum_{i=1}^d c_i v_i \right\|_{\mathcal{G}} + \|e\|_{\mathcal{G}} \\ &\leq 4k\|P\| + \frac{2d(\|P\| + 1)}{b} + 2. \end{aligned}$$

Choose δ such that $|z - s| < \delta$ implies

$$q(z, s) \left[4k\|P\| + \frac{2d(\|P\| + 1)}{b} + 2 \right] < 1/3.$$

Then, by Lemma 2.2,

$$\begin{aligned} \|e(z) - g(z)\|_{A\{z\}} &\leq \|e(s) - g(s)\|_{A\{s\}} + q(z, s)\|e - g\|_{\mathcal{G}} \\ &\leq 1/3 + q(z, s) \left[4k\|P\| + \frac{2d(\|P\| + 1)}{b} + 2 \right] < 2/3 \end{aligned}$$

which contradicts $\text{dist}(e(z), U_{\{z\}} + M_z) > 2/3$. Therefore $U_{\{z\}} + M_z = A\{z\}$.

Since $A\{s\} = U_{\{s\}} \oplus M_s$, $U_{\{s\}}$ is closed and M_s has finite dimension, we have $\chi_s > 0$. By Theorem 2.5, there exists $\delta > 0$ such that $|z - s| < \delta$ implies $\chi_z > 0$, which implies $U_{\{z\}} \cap M_z = \{0\}$. Hence the codimension of $U_{\{z\}}$ is d . $\square$

Theorem 2.7. *If $\ker T_s = \{0\}$ and $\text{codim}(T_s A\{s\}) = d < \infty$, then there exists $\delta > 0$, so that, for all $|z - s| < \delta$, $\ker T_z = \{0\}$ and $\text{codim } T_z A\{z\} = d$.*

Proof. The control over $\ker T_z$ is achieved by considering the following function:

$$r(T_z) = \inf\{\|Ta\|_{B\{z\}} \mid \|a\|_{A\{z\}} = 1\}.$$

Clearly if $r(T_z) > 0$, $\ker T_z = \{0\}$ and the range of T_z is closed. Conversely, if the range of T_z is closed and $\ker T_z = \{0\}$ then, by the open mapping theorem, $r(T_z) = \|T_z^{-1}\|^{-1} > 0$. Since $\text{codim } T_s = d < \infty$, the range of T_s is closed, and, since $\ker T_s = \{0\}$, we have $r(T_s) > 0$. Theorem 2.3 in [11] says that $r(T_z) > r(T_s)/2 > 0$ for all z which satisfy $|z - s| < \delta$, some $\delta > 0$.

Since $\|T\|_{A(\gamma) \rightarrow B(\gamma)} \leq M(\gamma)$ with $\int_{\Gamma} |\log M(\gamma)| dP_z(\gamma) < \infty$, by Lemma 1.8, there exists an outer function $G(\cdot)$ such that

$$(2.4) \quad |G(\gamma)M(\gamma)| = 1 \quad \text{for a.e. } \gamma \in \Gamma.$$

We claim that $G(\cdot)T\mathcal{G}(\mathcal{A})$ is a z -Calderón subspace of $\mathcal{G}(\mathcal{B})$ with constant c , for all z such that $|z - s| < \delta$, where

$$c = 4\|T_s^{-1}\| \sup_{\eta} \left\{ \exp \left(\int \log M(\gamma) dP_{\eta}(\gamma) \right) \mid |\eta - s| < \delta \right\}.$$

In fact, let $y \in G(\cdot)T\mathcal{G}(\mathcal{A})_z$. Then $y \in T_z\mathcal{A}$, and so there exists $x \in \mathcal{A}$ such that $Tx = y$ and $\|x\|_{A\{z\}} \leq \|T_z^{-1}\| \|y\|_{B\{z\}}$. We choose $g \in \mathcal{G}(\mathcal{A})$ such that $g(z) = x$ and

$$(2.5) \quad \|g\|_{\mathcal{G}(\mathcal{A})} \leq 2\|x\|_{A\{z\}} \leq 2\|T_z^{-1}\| \|y\|_{B\{z\}}.$$

Then $g(\zeta)/G(z) \in \mathcal{G}(\mathcal{A})$ and $H(\zeta) = G(\zeta)T(g(\zeta)/G(z)) \in G(\cdot)T\mathcal{G}(\mathcal{A})$. Clearly $H(z) = y$, and, by (2.4), (2.5),

$$\|H\|_{\mathcal{G}(\mathcal{B})} \leq \frac{2\|T_z^{-1}\| \|y\|_{B\{z\}}}{|G(z)|} \leq c\|y\|_{B\{z\}}.$$

Therefore $G(\cdot)T\mathcal{G}(\mathcal{A})$ is a z -Calderón subspace with constant c for all z such that $|z - s| < \delta$.

It is also clear that

$$(G(\cdot)T\mathcal{G}(\mathcal{A}))_{\{z\}} = T_z\mathcal{A}\{z\}.$$

Thus, by Theorem 2.6,

$$\begin{aligned} \operatorname{codim} T_z\mathcal{A}\{z\} &= \operatorname{codim}(G(\cdot)T\mathcal{G}(\mathcal{A}))_{\{z\}} \\ &= \operatorname{codim}(G(\cdot)T\mathcal{G}(\mathcal{A}))_{\{s\}} \\ &= \operatorname{codim} T_s\mathcal{A}\{s\}; \end{aligned}$$

$\ker T_z = \{0\}$, and the theorem is proved. $\square$

Corollary 2.8. *If $\dim \ker T_s = d < \infty$ and T_s is surjective, then there exists $\delta > 0$, so that for all $|z - s| < \delta$, $\dim \ker T_z = d < \infty$ and T_z is surjective.*

Proof. Consider $T^* : B^*(\gamma) \rightarrow A^*(\gamma)$. Clearly $\operatorname{codim} T_s^* = \dim \ker T_s = d < \infty$, $\dim \ker T_s^* = \operatorname{codim} T_s = 0$ and $\|T^*\|_{B^*(\gamma) \rightarrow A^*(\gamma)} = \|T\|_{A(\gamma) \rightarrow B(\gamma)}$. By assumption, $\{A_\gamma^*\}$, $\{B_\gamma^*\}$ are interpolation families and $(A\{z\})^* = \{A_\gamma^*\{z\}$, $(B\{z\})^* = \{B_\gamma^*\{z\}$.

By Theorem 2.7, there exists $\delta > 0$ so that $|z - s| < \delta$ implies $\ker T_z^* = \{0\}$ and $\operatorname{codim} T_z^* = d$. Therefore, for all these z we have $\dim \ker T_z = d$ and T_z is surjective. $\square$

REFERENCES

1. E. Albrecht, *Spectral interpolation*, Oper. Theory: Adv. Appl. **14** (1984), 13–37.
2. J. Bergh and J. Löfström, *Interpolation spaces*, Springer-Verlag, New York and Berlin, 1976.
3. A. P. Calderón, *Intermediate spaces and interpolation, the complex method*, Studia Math. **24** (1964), 113–190.
4. W. Cao and Y. Sagher, *Stability of Fredholm properties in interpolation scales*, Ark. für Mat. (to appear).

5. R. Coifman, M. Cwikel, R. Rochberg, Y. Sagher, and G. Weiss, *The complex method for interpolation of operators acting on families of Banach spaces*, Lecture Notes in Math. vol. 779, Springer-Verlag, Berlin, Heidelberg, and New York, 1980, 123–153.
6. —, *A theory of complex interpolation for families of Banach spaces*, Adv. in Math. **33** (1982), 203–229.
7. P. L. Duren, *Theory of H^p spaces*, Academic Press, New York and London, 1970.
8. J. Lindenstrauss and L. Tzafriri, *Classical Banach spaces*, Springer-Verlag, Berlin, Heidelberg and New York, 1977.
9. I. Ya. Sneiberg, *Spectral properties of linear operators in interpolation families of Banach spaces*, Mat. Issled. **9** (1974), 214–229.
10. G. Szegő, *Über die Randwerte einer analytischen Funktion*, Math. Ann. **84** (1921), 482–492.
11. A. T. Vignati and M. Vignati, *Spectral theory and complex interpolation*, J. Funct. Anal. **80** (1988), 383–397.
12. M. Zafran, *Spectral theory and interpolation of operators*, J. Funct. Anal. **36** (1980), 185–204.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT CHICAGO, CHICAGO, ILLINOIS 60680