THE HYPERSPACES OF SUBCONTINUA OF THE PSEUDO-ARC AND OF SOLENOIDS OF PSEUDO-ARCS ARE CANTOR MANIFOLDS

PAWEŁ KRUPSKI

(Communicated by James E. West)

ABSTRACT. New proofs of the above facts are based on specific homogeneity properties of the pseudo-arc and of solenoids of pseudo-arcs.

The reader is referred to [5] for hyperspace theory. It is known that if X is the pseudo-arc or a solenoid of pseudo-arcs (see [7] for the definition), then the hyperspace $C(X)$ of all nonvoid subcontinua of X is 2-dimensional. It is proved in [6] that if X is the pseudo-arc, then $C(X)$ is also a Cantor manifold, i.e., no 0-dimensional subset separates $C(X)$. In [2] a general theorem is presented that $C(X)$ has this property for an arbitrary metric, nondegenerate continuum X . Our proof of the theorem in the title is an application of the following result [3].

Lemma 1. *If X is an n -dimensional, locally compact, connected, homogeneous, metric space, then no $(n - 2)$ -dimensional subset separates X ($n \geq 1$).*

Lemma 2. *If a dense, connected subset of a metric separable space X is separated by no n -dimensional subset, then the space X has the same property. $\square$*

(1) Let X be the pseudo-arc. To show that $C(X)$ is a Cantor manifold it suffices to observe, by Lemmas 1 and 2, that the subspace $Y \subset C(X)$ of all nondegenerate, proper subcontinua of X is connected, locally compact, homogeneous (see [1]), as well as 2-dimensional and dense.

(2) Let X be a solenoid of pseudo-arcs with the continuous decomposition D into pseudo-arcs such that X/D is a solenoid S . The set D as a subspace of $C(X)$ is homeomorphic to S . As in (1) the open subspace Y of $C(X)$ is connected and dense. The set $Y \setminus D$ is dense in Y and is the union of two disjoint, open, connected, 2-dimensional subsets $M = \{y \in Y : d \neq y \subset d \in D\}$

Received by the editors February 25, 1989 and, in revised form, November 19, 1989.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 54B20; Secondary 54F45.

Key words and phrases. Pseudo-arc, solenoid of pseudo-arcs, hyperspace of subcontinua, homogeneous space, Cantor manifold.

This note was written while the author was visiting the Department of Mathematics, University of Helsinki.

and $N = \{y \in Y : d \neq y \supset d \in D\}$. It follows from [4] and from properties of solenoids of pseudo-arcs [7] that for every pair $y_1, y_2 \in M (y_1, y_2 \in N)$ there exists a homeomorphism $h: X \rightarrow X$ such that $h(y_1) = h(y_2)$. The induced homeomorphism $\hat{h}: C(X) \rightarrow C(X)$ satisfies $\hat{h}(M) = M$, $\hat{h}(N) = N$ and $\hat{h}(y_1) = y_2$, so both M and N are homogeneous and, by Lemma 1, no 0-dimensional subset separates neither M nor N . Suppose a 0-dimensional subset C separates Y . Without loss of generality we may assume that C is a closed subset of Y . It means $Y \setminus C = A \cup B$, where A, B are nonvoid, disjoint and open subsets of $C(X)$. In view of the above properties of M and N we may assume $M \subset A$ and $N \subset B$. Thus $C \subset D$. If there is $d \in D \setminus C$, then some order arc $\alpha \subset C(X)$ passing through d joins M and N , which is impossible, since $\alpha \cap D = \{d\}$ and C separates Y between M and N . Therefore $C = D$, hence C is not 0-dimensional, a contradiction.

Remark. A similar proof works for X being a solenoid. However in this case $C(X)$ is the cone over X , which is evidently a Cantor manifold.

REFERENCES

1. H. Cook, *A locally compact, homogeneous metric space which is not bihomogeneous*, Topology Proc. **11** (1986), 25–27.
2. J. Krasinkiewicz, *No 0-dimensional set disconnects the hyperspace of a continuum*, Bull. Acad. Polon. Sci. **19** (1971), 755–758.
3. P. Krupski, *Homogeneity and Cantor manifolds*, Proc. Amer. Math. Soc. (to appear).
4. G. R. Lechner, *Extending homeomorphisms on the pseudo-arc*, Trans. Amer. Math. Soc. **98** (1961), 369–394.
5. S. B. Nadler, Jr., *Hyperspaces of sets*, Marcel Dekker, New York, 1978.
6. T. Nishiura and C. J. Rhee, *The hyperspace of a pseudo-arc is a Cantor manifold*, Proc. Amer. Math. Soc. **31** (1972), 550–556.
7. J. T. Rogers, Jr., *Solenoids of pseudo-arcs*, Houston J. Math. **3** (1977), 531–537.

MATHEMATICAL INSTITUTE, UNIVERSITY OF WROCLAW, PL. GRUNWALDZKI 2/4, 50-384 WROCLAW, POLAND