APPROXIMATION OF FIXED POINTS OF ASYMPTOTICALLY NONEXPANSIVE MAPPINGS

JÜRGEN SCHU

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. Let T be an asymptotically nonexpansive self-mapping of a non-empty closed, bounded, and starshaped (with respect to zero) subset of a smooth reflexive Banach space possessing a duality mapping that is weakly sequentially continuous at zero. Then, if $\text{id} - T$ is demiclosed and T satisfies a strengthened regularity condition, the iteration process $z_{n+1} := \mu_{n+1} T^n(z_n)$ converges strongly to some fixed point of T , provided (μ_n) has certain properties.

0. INTRODUCTION

Motivated by the papers of B. Halpern [4] and P. Vijayaraju [7], we deal with the convergence of almost fixed points $x_n = \mu_n T^n(x_n)$ of an asymptotically nonexpansive mapping T . This class of functions was introduced by K. Goebel and W. A. Kirk [3] in 1972. In §1, convergence of the sequence (x_n) to some fixed point of T is shown, assuming that T is uniformly asymptotically regular and $\text{id} - T$ is demiclosed. These assumptions have been made by P. Vijayaraju [7] to ensure the existence of a fixed point of T .

By strengthening the regularity condition on T in §2, we establish the convergence of the explicit iteration scheme $z_{n+1} := \mu_{n+1} T^n(z_n)$ to some fixed point of T . This iteration method is similar to one that was introduced by B. Halpern [4].

1. CONVERGENCE OF THE SEQUENCE OF APPROXIMATE FIXED POINTS

We shall begin by recalling some definitions needed in the sequel.

Definition 1.1. Let $(E, \|\cdot\|)$ be a normed space, $\emptyset \neq A \subset E$; $\mu: \mathbb{R}^+ \rightarrow \mathbb{R}^+$; $J: E \rightarrow E^*$; and $x_0 \in E$.

- (1) $(E, \|\cdot\|)$ is called *smooth*: $\iff \|\cdot\|$ is Gateaux-differentiable on $\partial \overline{B}(0, 1)$.
- (2) μ is said to be a *gauge function*: $\iff \mu$ is continuous and strictly increasing with $\mu(0) = 0$ and $\lim_{x \rightarrow \infty} \mu(x) = \infty$.

Received by the editors November 12, 1989 and, in revised form, February 20, 1990.
 1980 *Mathematics Subject Classification* (1985 Revision). Primary 47H10; Secondary 40A05, 47H09.

Key words and phrases. Fixed point, iteration process, asymptotically nonexpansive mapping.

- (3) The set-valued *duality mapping* associated with μ is given by $J_E^\mu(x) := \{u \in E^* | u(x) = \|u\| \|x\| \text{ and } \|u\| = \mu(\|x\|)\}$ and J is called a duality mapping with respect to μ : $\iff J(x) \in J_E^\mu(x)$ for all $x \in E$. For abbreviation, we set $J_E := J_E^{\text{id}}$.
- (4) J is *weakly sequentially continuous* at x_0 : $\iff$ for all $(x_n) \in E^{\mathbb{N}}$ with $(x_n) \rightharpoonup x_0$, it follows that $(J(x_n)) \overset{*}{\rightharpoonup} J(x_0)$.

Remark. It is well known that J_E^μ is single-valued if and only if $(E, \|\cdot\|)$ is smooth (see, e.g., [1]). In this case, we regard J_E^μ as a mapping from E to E^* . In all our proofs we assume, without loss of generality, that J_E^μ is normalized, i.e., $\mu = \text{id}$. Furthermore, in the sequel all normed spaces are assumed to be real Banach spaces.

Definition 1.2. Let $(E, \|\cdot\|)$ be a normed space; $\emptyset \neq A \subset E$; $T: A \rightarrow A$; $(k_n) \in [1, \infty)^{\mathbb{N}}$.

- (1) T is said to be *asymptotically nonexpansive* with sequence (k_n) : $\iff \lim(k_n) = 1$ and $\|T^n(x) - T^n(y)\| \leq k_n \|x - y\|$ for all $n \in \mathbb{N}$ and all $x, y \in A$.
- (2) T is called *demiclosed*: $\iff$ for all $(x_n) \in A^{\mathbb{N}}$ and all $x, y \in E$ with $(x_n) \rightharpoonup x$ and $\lim(T(x_n)) = y$, it follows that $x \in A$ and $T(x) = y$.
- (3) T is called *uniformly asymptotically regular*: $\iff$ for each $\varepsilon > 0$ there is $n_0 \in \mathbb{N}$, such that $\|T^n(x) - T^{n+1}(x)\| \leq \varepsilon$ for all $n \geq n_0$ and all $x \in A$.

Lemma 1.3. Let $(E, \|\cdot\|)$ be a smooth normed space; $\emptyset \neq A \subset E$; $T: A \rightarrow A$ asymptotically nonexpansive with sequence $(k_n) \in [1, \infty)^{\mathbb{N}}$; $x, y \in A$; $\lambda \in (0, 1)$; $n \in \mathbb{N}$; $x = \lambda T^n(x)$; and $y = T^n(y)$. Then

$$J_E(y - x)(x) \geq -\frac{k_n - 1}{1/\lambda - 1} \|x - y\|^2.$$

Proof. Set $\alpha := \frac{1}{\lambda} - 1 > 0$, $z := \alpha x$, and $w := y - x$. Then

$$\begin{aligned} \|z - w\| &= \|(1 + \alpha)x - y\| = \|\tfrac{1}{\lambda}x - y\| \\ &= \|T^n(x) - T^n(y)\| \leq k_n \|x - y\| = k_n \|w\|. \end{aligned}$$

Hence

$$\begin{aligned} J_E(y - x)(x) &= \tfrac{1}{\alpha} J_E(w)(z) = \tfrac{1}{\alpha} J_E(w)(z - w) + \tfrac{1}{\alpha} J_E(w)(w) \\ &\geq -\tfrac{1}{\alpha} \|w\| \|z - w\| + \tfrac{1}{\alpha} \|w\|^2 \geq \tfrac{1}{\alpha} (1 - k_n) \|w\|^2 \\ &= ((1 - k_n)/((\tfrac{1}{\lambda}) - 1)) \|x - y\|^2. \quad \square \end{aligned}$$

The following lemma can be verified by an easy calculation.

Lemma 1.4. Let $\lambda \in (\frac{1}{2}, 1)$. Then

- (1) $\lambda^2/(2\lambda - 1) > 1$, and
 (2) $(\lambda k - \lambda)/(k - \lambda) \leq 1 - \lambda$ for all $k \in [1, \lambda^2/(2\lambda - 1)]$.

Remark. Part (1) of Lemma 1.4 shows that the assumption $k_n \leq \lambda_n^2/(2\lambda_n - 1)$ and $k_n \in [1, \infty)$ of our next lemma makes sense.

Lemma 1.5. *Let $(E, \|\cdot\|)$ be a smooth normed space possessing a duality mapping $J: E \rightarrow E^*$ that is weakly sequentially continuous at 0; $\emptyset \neq A \subset E$ bounded; $T: A \rightarrow A$ asymptotically nonexpansive with sequence $(k_n) \in [1, \infty)^\mathbb{N}$; $(x_n) \in A^\mathbb{N}$; $x \in A$; (x_{φ_n}) a subsequence of (x_n) with $(x_{\varphi_n}) \rightarrow x$; $(\lambda_n) \in (\frac{1}{2}, 1)^\mathbb{N}$ with $\lim(\lambda_n) = 1$; $x = T(x)$; $x_n = (\lambda_n/k_n)T^n(x_n)$ for all $n \in \mathbb{N}$; and $k_n \leq \lambda_n^2/(2\lambda_n - 1)$ for all $n \in \mathbb{N}$. Then*

- (1) $\lim(x_{\varphi_n}) = x$, and
- (2) $J(y - x)(x) \geq 0$ for all $y \in \text{Fix}(T)$.

($\text{Fix}(T) := \{z \in A \mid T(z) = z\}$)

Proof. (1) Since $(\lambda_n/k_n) \in (0, 1)^\mathbb{N}$, it follows from Lemma 1.3 that

$$J(x - x_n)(x_n) \geq -\frac{k_n - 1}{k_n/\lambda_n - 1} \|x_n - x\|^2 \quad \text{for all } n \in \mathbb{N}.$$

We may choose $M > 0$ such that $\|x_n - z\|^2 \leq M$ for all $n \in \mathbb{N}$ and all $z \in A$ because of the boundedness of A . Obviously, $(k_n - 1)/(k_n/\lambda_n - 1) \geq 0$ and from Lemma 1.4 we know, that

$$\frac{k_n - 1}{k_n/\lambda_n - 1} = \frac{\lambda_n k_n - \lambda_n}{k_n - \lambda_n} \leq 1 - \lambda_n.$$

Hence $J(x - x_n)(x_n) \geq -(1 - \lambda_n)M$, and therefore, for all $n \in \mathbb{N}$,

$$\|x - x_n\|^2 = J(x - x_n)(x) - J(x - x_n)(x_n) \leq J(x - x_n)(x) + (1 - \lambda_n)M,$$

where

- (a) $\lim(1 - \lambda_n) = 0$,
- (b) $(x_{\varphi_n} - x) \rightarrow 0$, hence $\lim J(x - x_{\varphi_n})(x) = 0$.

Thus we conclude that $\lim \|x - x_{\varphi_n}\| = 0$.

(2) Fix $y \in \text{Fix}(T)$. As already shown in (1), for $y = T(y)$ instead of x , it follows that

$$(*) \quad J(y - x_n)(x_n) \geq -(1 - \lambda_n)M \quad \text{for all } n \in \mathbb{N}.$$

Since $(E, \|\cdot\|)$ is smooth, J is strong-weak^(*) continuous (see e.g., [2]), and so from $\lim(y - x_{\varphi_n}) = y - x$, we conclude that $(J(y - x_{\varphi_n})) \xrightarrow{*} J(y - x)$. Hence $\lim J(y - x_{\varphi_n})(x_{\varphi_n}) = J(y - x)(x)$. This, together with $(*)$ and $\lim(1 - \lambda_n) = 0$, yields $J(y - x)(x) \geq 0$. $\square$

Remark. Lemmas 1.3 and 1.5 are similar to some related results for nonexpansive mappings; namely, [6, Lemmas 4, 5]. The following theorem was demonstrated by P. Vijayaraju in the course of the proof of [7, Theorems 2.1, 2.2].

Theorem 1.6. Let $(E, \|\cdot\|)$ be a Banach space; $\emptyset \neq A \subset E$ closed, bounded, and starshaped with respect to 0; $T: A \rightarrow A$ asymptotically nonexpansive with sequence $(k_n) \in [1, \infty)^{\mathbb{N}}$ and uniformly asymptotically regular; $(\lambda_n) \in [0, 1)^{\mathbb{N}}$; and $\lim(\lambda_n) = 1$. Then

- (1) for each $n \in \mathbb{N}$ there is exactly one $x_n \in A$ such that $x_n = (\lambda_n/k_n)T^n(x_n)$,
and
- (2) $\lim \|x_n - T(x_n)\| = 0$.

Now we are able to prove the main result of this section.

Theorem 1.7. Let $(E, \|\cdot\|)$ be a smooth reflexive Banach space possessing a duality mapping $J: E \rightarrow E^*$ that is weakly sequentially continuous at 0; $\emptyset \neq A \subset E$ closed, bounded, and starshaped with respect to 0; $T: A \rightarrow A$ asymptotically nonexpansive with sequence $(k_n) \in [1, \infty)^{\mathbb{N}}$ and uniformly asymptotically regular; $\text{id} - T$ demiclosed; $(\lambda_n) \in (\frac{1}{2}, 1)^{\mathbb{N}}$; $\lim(\lambda_n) = 1$; $k_n \leq \lambda_n^2/(2\lambda_n - 1)$ for all $n \in \mathbb{N}$. Then

- (1) for each $n \in \mathbb{N}$ there is exactly one $x_n \in A$ such that $x_n = (\lambda_n/k_n)T^n(x_n)$,
and
- (2) (x_n) converges strongly to some fixed point of T .

Proof. Part (1) follows from Theorem 1.6, so that it remains to prove part (2). Since $(E, \|\cdot\|)$ is reflexive and A is bounded, there exists $z \in E$ and a subsequence (x_{μ_n}) of (x_n) such that $(x_{\mu_n}) \rightharpoonup z$ (Pettis' theorem). From Theorem 1.6, $\lim \|x_n - T(x_n)\| = 0$, so from the demiclosedness of $\text{id} - T$, $z \in A$ and $T(z) = z$.

Applying Lemma 1.5, we get

$$(*) \quad J(y - z)(z) \geq 0 \quad \text{for all } y \in \text{Fix}(T).$$

Fix $\varphi: \mathbb{N} \rightarrow \mathbb{N}$ and injective.

Repeating the above argument, we find a subsequence $(x_{\varphi_{\mu_n}})$ of (x_{φ_n}) converging weakly to some $x \in A$ such that $x = T(x)$ and

$$(**) \quad J(y - x)(x) \geq 0 \quad \text{for all } y \in \text{Fix}(T).$$

Substituting x into $(*)$ and z into $(**)$, it follows that $J(x - z)(z) \geq 0$ and $J(z - x)(x) \geq 0$. Hence $0 \leq J(x - z)(-x) + J(x - z)(z) = -\|x - z\|^2 \leq 0$, so that $x = z$, and therefore $(x_{\varphi_{\mu_n}}) \rightharpoonup z$. This shows that $(x_n) \rightharpoonup z$, and we now apply Lemma 1.5 (with $\varphi := \text{id}$) to obtain $\lim \|x_n - z\| = 0$. $\square$

Remark. For $(k_n) \in (1, \infty)^{\mathbb{N}}$ with $\lim(k_n) = 1$, the conditions $(\lambda_n) \in (\frac{1}{2}, 1)^{\mathbb{N}}$, $\lim(\lambda_n) = 1$, and $k_n \leq \lambda_n^2/(2\lambda_n - 1)$ for all $n \in \mathbb{N}$ are fulfilled, for example, if one chooses $\lambda_n := k_n - \sqrt{k_n^2 - k_n}$ for all $n \in \mathbb{N}$.

2. AN EXPLICIT ITERATION SCHEME

Now we give an explicit iteration scheme for the approximation of a fixed point of T . In analogy to B. Halpern's work [4], we use the fact that the

implicit iteration process $x_n = (\lambda_n/k_n)T^n(x_n)$ already converges to some fixed point of T to show that the same holds for the explicit iteration procedure $z_{n+1} := (\lambda_{n+1}/k_{n+1})T^n(z_n)$.

In addition to several technical assumptions on the values λ_n and k_n , we require that T fulfills a certain strengthened regularity condition (assumption (b) of Theorem 2.2). Although parts of the proof of Theorem 2.2 are parallel to those of the proof of [4, Theorem 3], they are included for the sake of completeness.

Definition 2.1. Let $(\varepsilon_n) \in (0, \infty)^{\mathbb{N}}$ and $(\mu_n) \in (0, 1)^{\mathbb{N}}$. $((\varepsilon_n), (\mu_n))$ is called *admissible* : $\Longleftrightarrow$

- (1) (ε_n) is decreasing;
- (2) (μ_n) is strictly increasing with $\lim(\mu_n) = 1$;
- (3) there exists $(\beta_n) \in \mathbb{N}^{\mathbb{N}}$ such that
 - (i) (β_n) is increasing,
 - (ii) $\lim(\beta_n(1 - \mu_n)) = \infty$,
 - (iii) $\lim((1 - \mu_{n+\beta_n})/(1 - \mu_n)) = 1$, and
 - (iv) $\lim((\varepsilon_n \beta_n \mu_{n+\beta_n})/(1 - \mu_n)) = 0$.

Remark. (a) Parts (2) and (3) (i)–(iii) are due to B. Halpern [4]. (b) $\lim(\beta_n) = \infty$ because of (3)(ii) and (2), so that it follows from (3)(iv) that $\lim(\varepsilon_n) = 0$.

Theorem 2.2. Let $(E, \|\cdot\|)$ be a normed space; $\emptyset \neq A \subset E$ bounded and starshaped with respect to 0; $T: A \rightarrow A$ asymptotically nonexpansive with sequence $(k_n) \in [1, \infty)^{\mathbb{N}}$; $(\lambda_n) \in (0, 1)^{\mathbb{N}}$ with $\lim(\lambda_n) = 1$; $(x_n) \in A^{\mathbb{N}}$; $\mu_n := \lambda_n/k_n$ for all $n \in \mathbb{N}$; $z_0 \in A$; $x_n = (\lambda_n/k_n)T^n(x_n)$ for all $n \in \mathbb{N}$, and $z_{n+1} := (\lambda_{n+1}/k_{n+1})T^n(z_n)$ for all $n \in \mathbb{N}_0$; $(\varepsilon_n) \in (0, \infty)^{\mathbb{N}}$ such that $((\varepsilon_n), (\mu_n))$ is admissible;

- (a) $((1 - \mu_n)/(1 - \lambda_n))$ bounded;
- (b) $\|T^n(x) - T^{n+1}(x)\| \leq \varepsilon_n$ for all $n \in \mathbb{N}$ and all $x \in A$.

Assume further that $q := \lim(x_n)$ exists. Then $\lim(z_n) = q$.

Proof. Observe that $(z_n) \in A^{\mathbb{N}}$ is well defined, because $T(A) \subset A$, $(\mu_n) \in (0, 1)^{\mathbb{N}}$, and A is starshaped with respect to 0. Since A is bounded and T is Lipschitzian, there is a $d > 0$ such that $\|z\| \leq d$ for all $z \in A$ and there is a $c > 0$ such that $\|T(w)\| \leq c$ for all $w \in A$. Additionally, (a) implies the existence of a number $\kappa > 0$, such that $(1 - \mu_n)/(1 - \lambda_n) \leq \kappa$ for all $n \in \mathbb{N}$, and we may choose (β_n) according to Definition 2.1. For $n > p \geq i$, we have

$$\|z_{p+1} - x_i\| \leq \|\mu_{p+1}T^p(z_p) - \mu_iT^i(z_p)\| + \|\mu_iT^i(z_p) - \mu_iT^i(x_i)\|,$$

where

$$\|\mu_iT^i(z_p) - \mu_iT^i(x_i)\| \leq \mu_i k_i \|z_p - x_i\| = \lambda_i \|z_p - x_i\|,$$

and

$$\begin{aligned} & \|\mu_{p+1}T^p(z_p) - \mu_iT^i(z_p)\| \\ & \leq \|\mu_{p+1}T^p(z_p) - \mu_{p+1}T^i(z_p)\| + \|\mu_{p+1}T^i(z_p) - \mu_iT^i(z_p)\| \\ & \leq \mu_{p+1}\|T^p(z_p) - T^i(z_p)\| + |\mu_{p+1} - \mu_i|c. \end{aligned}$$

Since $n > p \geq i$, it follows that $|\mu_{p+1} - \mu_i| \leq \mu_n - \mu_i$. Additionally,

$$\begin{aligned} \|T^p(z_p) - T^i(z_p)\| & \leq \sum_{j=i}^{p-1} \|T^{j+1}(z_p) - T^j(z_p)\| \\ & \leq \sum_{j=i}^{p-1} \varepsilon_j \leq \sum_{j=i}^{p-1} \varepsilon_i = (p-i)\varepsilon_i \leq (n-i)\varepsilon_i. \end{aligned}$$

Combining the above inequalities, we get

$$(1) \quad \|z_{p+1} - x_i\| \leq \lambda_i\|z_p - x_i\| + (\mu_n - \mu_i)c + \mu_n(n-i)\varepsilon_i \quad \text{for all } n > p \geq i,$$

from which it follows by induction that

$$\|z_{m+j} - x_i\| \leq \lambda_i^j\|z_m - x_i\| + (\mu_n - \mu_i)c \sum_{\nu=0}^{j-1} \lambda_i^\nu + \mu_n(n-i)\varepsilon_i \sum_{\nu=0}^{j-1} \lambda_i^\nu,$$

for all $n > m \geq i$ and $j \leq n - m$. Since $\sum_{\nu=0}^{j-1} \lambda_i^\nu \leq 1/(1 - \lambda_i)$, we obtain, defining $j = n - m$,

$$(2) \quad \|z_n - x_i\| \leq \lambda_i^{n-m}\|z_m - x_i\| + (\mu_n - \mu_i)(1 - \lambda_i)^{-1}c + \mu_n(n-i)\varepsilon_i(1 - \lambda_i)^{-1}$$

for all $n > m \geq i$.

Set $m := i$ and $n := i + \beta_i > m$ to get

$$\|z_{i+\beta_i} - x_i\| \leq 2d\lambda_i^{\beta_i} + (\mu_{i+\beta_i} - \mu_i)(1 - \lambda_i)^{-1}c + \mu_{i+\beta_i}\beta_i\varepsilon_i(1 - \lambda_i)^{-1},$$

for all $i \in \mathbb{N}$. Observe that

$$(i) \quad |\mu_{i+\beta_i} - \mu_i|(1 - \lambda_i)^{-1} = |1 - (1 - \mu_{i+\beta_i})(1 - \mu_i)^{-1}||1 - \mu_i|/(1 - \lambda_i)| \\ \leq \kappa|1 - (1 - \mu_{i+\beta_i})(1 - \mu_i)^{-1}|,$$

from which we conclude that $\lim((\mu_{i+\beta_i} - \mu_i)/(1 - \lambda_i)) = 0$, taking into account the admissibility of $((\varepsilon_n), (\mu_n))$. Also,

(ii) $\log(\lambda_i^{\beta_i}) \leq \beta_i(\lambda_i - 1) \leq -\beta_i(1 - \mu_i)1/\kappa$. Since $((\varepsilon_n), (\mu_n))$ is admissible, it follows that $\lim(\log(\lambda_i^{\beta_i})) = -\infty$, and thus $\lim(\lambda_i^{\beta_i}) = 0$. In addition,

(iii) $0 \leq \mu_{i+\beta_i}\beta_i\varepsilon_i(1 - \lambda_i)^{-1} \leq \kappa\mu_{i+\beta_i}\beta_i\varepsilon_i(1 - \mu_i)^{-1}$, from which it follows that $\lim((\mu_{i+\beta_i}\beta_i\varepsilon_i)/(1 - \lambda_i)) = 0$, again using the admissibility of $((\varepsilon_n), (\mu_n))$. Hence

$$(3) \quad \lim \|z_{i+\beta_i} - x_i\| = 0.$$

As already noticed in the proof of [4, Theorem 3], there exists $j: \mathbb{N} \rightarrow \mathbb{N}$ and $n_0 \in \mathbb{N}$ such that $\lim(j_n) = \infty$ and $j_n + \beta_{j_n} < n \leq j_n + 1 + \beta_{j_n+1}$ for all $n \geq n_0$. Now let $n \geq n_0$. With $n > m := j_n + \beta_{j_n} \geq i := j_n + 1$, it follows from (2) that

$$\begin{aligned} \|z_n - x_{j_n+1}\| &\leq \|z_{(j_n+\beta_{j_n})} - x_{j_n+1}\| + (\mu_n - \mu_{j_n+1})(1 - \lambda_{j_n+1})^{-1}c \\ &\quad + \mu_n(n - (j_n + 1))\varepsilon_{j_n+1}(1 - \lambda_{j_n+1})^{-1} \quad \text{for all } n \in \mathbb{N}, \end{aligned}$$

where

(i) $0 \leq (\mu_n - \mu_{j_n+1})(1 - \lambda_{j_n+1})^{-1} \leq (\mu_{(j_n+1+\beta_{j_n+1})} - \mu_{j_n+1})(1 - \lambda_{j_n+1})^{-1}$, which tends to 0 by 2(i).

(ii) $0 \leq \mu_n(n - (j_n + 1))\varepsilon_{j_n+1}(1 - \lambda_{j_n+1})^{-1} \leq \mu_{(j_n+1+\beta_{j_n+1})}\beta_{j_n+1}\varepsilon_{j_n+1}(1 - \lambda_{j_n+1})^{-1}$, which tends to 0 by (2)(iii).

(iii) $\|z_{(j_n+\beta_{j_n})} - x_{j_n+1}\| \leq \|z_{(j_n+\beta_{j_n})} - x_{j_n}\| + \|x_{j_n} - q\| + \|q - x_{j_n+1}\|$, which also tends to 0 because $\lim \|x_n - q\| = 0$ and $\lim \|z_{(j_n+\beta_{j_n})} - x_{j_n}\| = 0$ by (3). Hence $\lim \|z_n - x_{j_n+1}\| = 0$. Finally, since $\|z_n - q\| \leq \|z_n - x_{j_n+1}\| + \|x_{j_n+1} - q\|$, it follows that $\lim \|z_n - q\| = 0$. $\square$

We now prove the main result of this section.

Theorem 2.3. *Let $(E, \|\cdot\|)$ be a reflexive smooth Banach space possessing a duality mapping that is weakly sequentially continuous at 0; $\emptyset \neq A \subset E$ closed, bounded, and starshaped with respect to 0; $T: A \rightarrow A$ asymptotically nonexpansive with sequence $(k_n) \in [1, \infty)^\mathbb{N}$; $\text{id} - T$ demiclosed; $(\lambda_n) \in (\frac{1}{2}, 1)^\mathbb{N}$; $z_0 \in A$; $z_{n+1} := (\lambda_{n+1}/k_{n+1})T^n(z_n)$ for all $n \in \mathbb{N}_0$;*

- (a) $\lim(\lambda_n) = 1$; $\mu_n := \lambda_n/k_n$ for all $n \in \mathbb{N}$; $k_n \leq \lambda_n^2/(2\lambda_n - 1)$ for all $n \in \mathbb{N}$; $((1 - \mu_n)/(1 - \lambda_n))$ bounded; $(\varepsilon_n) \in (0, \infty)^\mathbb{N}$, such that $((\varepsilon_n), (\mu_n))$ is admissible;
- (b) $\|T^n(x) - T^{n+1}(x)\| \leq \varepsilon_n$ for all $n \in \mathbb{N}$ and all $x \in A$.

Then (z_n) converges strongly to some fixed point of T .

Proof. Since $((\varepsilon_n), (\mu_n))$ is admissible, it follows that $\lim(\varepsilon_n) = 0$, so that assumption (b) implies that T is uniformly asymptotically regular. The result follows by a combination of Theorems 1.7, 2.2. $\square$

Remark. (1) For a related result concerning the weak convergence of the successive approximations $x_{n+1} := T(x_n)$ to some fixed point of an asymptotically nonexpansive mapping T in an Opial space we refer the reader to [5].

(2) Theorem 2.3, of course, is applicable to nonexpansive mappings as well (set $k_n := 1$ for all $n \in \mathbb{N}$). In this special case, however, there is a much better result: [6, Theorem 10] allows us to drop the assumptions that $\text{id} - T$ is demiclosed and that $\|T^n(x) - T^{n+1}(x)\| \leq \varepsilon_n$ for all $n \in \mathbb{N}$ and all $x \in A$, provided use is made of the iteration scheme $z_{n+1} := \lambda_{n+1}T(z_n)$, T is nonexpansive, and $(E, \|\cdot\|)$ possesses a duality mapping that is weakly sequentially continuous on the whole of E .

We close this section with a special case of Theorem 2.3, which shows that it is really possible to fulfill the assumptions made on (λ_n) , (μ_n) , (ε_n) , and (k_n) .

Theorem 2.4. *Let $(E, \|\cdot\|)$ be a reflexive smooth Banach space possessing a duality mapping that is weakly sequentially continuous at 0; $\emptyset \neq A \subset E$ closed, bounded, and starshaped with respect to 0; $T: A \rightarrow A$ asymptotically nonexpansive with sequence $(1 + 1/(\sqrt{n+1} - 1))$; $\text{id} - T$ demiclosed; $\mu_n := 1 - (n+1)^{-1/4}$ for all $n \in \mathbb{N}$; $z_0 \in A$; $z_{n+1} := \mu_{n+1} T^n(z_n)$ for all $n \in \mathbb{N}_0$; and*

$$(*) \quad \|T^n(x) - T^{n+1}(x)\| \leq (n+1)^{-7/8} \quad \text{for all } n \in \mathbb{N} \text{ and all } x \in A.$$

Then (z_n) converges strongly to some fixed point of T .

Proof. For $n \in \mathbb{N}$ define $k_n := 1 + 1/(\sqrt{n+1} - 1)$, $\lambda_n := k_n - \sqrt{k_n^2 - k_n}$, $\varepsilon_n := (n+1)^{-7/8}$, and $\beta_n := [(n+1)^{1/2}]_G$ (cf. Definition 2.1), where $[x]_G := \max\{n \in \mathbb{N}_0 | n \leq x\}$. Then easy calculations show that all the assumptions of Theorem 2.3 are fulfilled, especially that $\mu_n = \lambda_n/k_n$ for all $n \in \mathbb{N}$ and that $((\varepsilon_n), (\mu_n))$ is admissible. $\square$

Remark. For $x \in A$ and $n, m \in \mathbb{N}$ with $n > m$, the following holds:

$$\|T^n(x) - T^m(x)\| \leq \sum_{j=m}^{n-1} \|T^{j+1}(x) - T^j(x)\| \leq \sum_{j=m}^{n-1} (j+1)^{-7/8}.$$

Nevertheless, since $\sum_{j=1}^{\infty} (j+1)^{-7/8} = \infty$, $(*)$ does not imply that $(T^n(x))$ is a Cauchy sequence (from which it would follow that $(T^n(x))$ converges strongly to some fixed point of T).

ACKNOWLEDGMENT

The author is grateful to Professor J. Reinermann for his interest in this paper.

REFERENCES

1. B. Beauzamy, *Introduction to Banach spaces and their geometry*, Mathematics Studies, vol. 68, North-Holland, Amsterdam, 1982.
2. J. Diestel, *Geometry of Banach spaces—selected topics*; Lecture Notes in Math., vol. 485, Springer-Verlag, Berlin, Heidelberg, and New York, 1975.
3. K. Goebel and W. A. Kirk, *A fixed point theorem for asymptotically nonexpansive mappings*, Proc. Amer. Math. Soc. **35**(1) (1972), 171–174.
4. B. Halpern, *Fixed points of nonexpanding maps*, Bull. Amer. Math. Soc. **73** (1967), 957–961.
5. S. K. Samanta, *Fixed point theorems in a Banach space satisfying Opial's condition*, J. Indian Math. Soc. **45** (1981), 251–258.

6. J. Schu, *Iterative approximation of fixed points of nonexpansive mappings with starshaped domain*, Comment. Math. Univ. Carolinae (to appear).
7. P. Vijayaraju, *Fixed point theorems for asymptotically nonexpansive mappings*, Bull. Calcutta Math. Soc. **80** (1988), 133–136.

RHEINISCH-WESTFÄLISCHE TECHNISCHE HOCHSCHULE AACHEN, LEHRSTUHL C FÜR MATHEMATIK,
TEMLERGRABEN 55, D-5100 AACHEN, FEDERAL REPUBLIC OF GERMANY