RADII PROBLEMS FOR GENERALIZED SECTIONS OF CONVEX FUNCTIONS

RICHARD FOURNIER AND HERB SILVERMAN

(Communicated by Clifford J. Earle, Jr.)

ABSTRACT. A classical theorem of Szëgo states that for functions $f(z) = z + \sum_{k=2}^{\infty} a_k z^k$ convex in $|z| < 1$, the sequence of partial sums $f_n(z) = z + \sum_{k=2}^n a_k z^k$ must be convex in $|z| < \frac{1}{4}$. For the more general family consisting of functions of the form $z + \sum_{k=2}^{\infty} a_{n_k} z^{n_k}$, where $\{n_k\}$ denotes an increasing (finite or infinite) sequence of integers (≥ 2), we find the radius of convexity (≈ 0.21) and the radius of starlikeness (≈ 0.37). The extremal function in both cases is $z + z^2/(1 - z^2) = z + \sum_{k=1}^{\infty} z^{2k}$ associated with the convex function $z/(1 - z) = z + \sum_{k=2}^{\infty} z^k$.

INTRODUCTION

Denote by S_0 the family consisting of functions $f(z) = z + \sum_{n=2}^{\infty} a_n z^n$ analytic and starlike univalent in $\Delta = \{z | |z| < 1\}$ and by K the subfamily of S_0 containing the convex functions. It is an old and well-known result of Szegő [9] that for any $N \geq 2$,

$$f_N(z) = z + \sum_{n=2}^N \frac{a_n}{4^{n-1}} z^n \in K \quad \text{if} \quad f(z) = z + \sum_{n=2}^{\infty} a_n z^n \in K,$$

in other words, any section of a function in K is convex univalent for $|z| < \frac{1}{4}$. Let $\{n_k\}_{k=2}^{\infty}$ denote an arbitrary increasing sequence (finite or infinite) of positive integers with $k \leq n_k$; in a recent paper [8] one of us conjectured that if $f \in K$ as above, then

$$z + \sum_{k=2}^{\infty} \frac{a_{n_k}}{4^{n_k-1}} z^{n_k} \in K \quad \text{and} \quad z + \sum_{k=2}^{\infty} \frac{a_{n_k}}{2^{n_k-1}} z^{n_k} \in S_0.$$

This would generalize Szegő's theorem to more general sequences.

To see that this conjecture is false, we consider the function $g(z) = z + \sum_{k=1}^{\infty} z^{2k} = z + z^2/(1 - z^2)$ associated with the convex function $f(z) = z + \sum_{k=2}^{\infty} z^k = z/(1 - z)$. Then $1 + zg''(z)/g'(z) = 0$ at $z = -c \sim -.20936$ and

Received by the editors October 10, 1989 and, in revised form, February 13, 1990.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 30C45; Secondary 30C50.

The first author was supported by an ACSAIR (Quebec) grant.

This work was completed while the second author was on sabbatical leave as a Visiting Scholar at the University of California at Davis.

$zg'(z)/g(z) = 0$ at $z = -b \sim -.3715$. It is the object of this note to show that the above g is extremal for the radii of convexity and starlikeness. In fact we shall prove

Theorem 1. *If $\{n_k\}_{k=2}^\infty$ is an increasing sequence of integers with $k \leq n_k$ and $f(z) = z + \sum_{n=2}^\infty a_n z^n \in K$, then $\tilde{f}(z) = z + \sum_{k=2}^\infty a_{n_k} z^{n_k}$ is convex univalent in $|z| < c$ where $c \sim .20936$ is the unique root in $(0,1)$ of the equation*

$$x \frac{1+x^2}{(1-x^2)^3} = \frac{1}{4}.$$

Theorem 2. *If $\{n_k\}_{k=2}^\infty$ is an increasing sequence of integers with $k \leq n_k$ and $f(z) = z + \sum_{n=2}^\infty a_n z^n \in K$ then $\tilde{f}(z) = z + \sum_{k=2}^\infty a_{n_k} z^{n_k}$ is starlike univalent in $|z| < b \sim .3715$ where b is the unique root in $(0,1)$ of the equation*

$$\frac{x}{(1-x^2)^2} = \frac{1}{2}.$$

Since $zf' \in S_0$ if and only if $f \in K$, a consequence of Theorem 1 follows.

Corollary. *Using the notation of Theorem 1, if $f \in S_0$ then $\tilde{f}$ is starlike univalent in $|z| < c$.*

There exists considerable literature concerning sections of univalent functions and we refer the reader to [2, Chapter 8] for a general survey. In particular it has been established (see [2, 4, 5] or [8]) that in the proofs of Theorem 1 and Theorem 2 we may assume $f(z) \equiv z/(1-z)$. This is due to a convolution theorem of Ruscheweyh and Sheil-Small [7].

Let the functions $f(z) = z + \sum_{n=2}^\infty a_n z^n$, $g(z) = z + \sum_{n=2}^\infty b_n z^n$ be analytic in Δ ; their convolution (or Hadamard product) is the function $f * g$ defined by

$$f * g(z) = z + \sum_{n=2}^\infty a_n b_n z^n.$$

For each $T \in \mathbf{R}$ we define

$$h_T(z) \equiv \frac{z/(1-z)^2 + iTz/(1-z)}{1+iT}.$$

Clearly a locally univalent function $u(z) = z + \sum_{n=2}^\infty a_n z^n$ is convex in Δ if and only if $(zu'(z) * h_T(z))/z \neq 0$ for all real T and $z \in \Delta$. Our proofs rely on the following results, due essentially to Ruscheweyh (see [3, 6]).

Theorem A. *Let $u(z) = z + \sum_{n=2}^\infty a_n z^n \in K$ such that*

$$\inf_{\substack{T \in \mathbf{R} \\ z \in \Delta}} \left| \frac{zu'(z) * h_T(z)}{z} \right| \geq \delta > 0.$$

Then $v(z) = z + \sum_{n=2}^\infty b_n z^n \in K$, provided that $\sum_{n=2}^\infty n^2 |a_n - b_n| \leq \delta$.

Note that Theorem A is an extension of a classical result [1]: the case $u(z) \equiv z$ is usually called Alexander's Theorem.

Theorem B. Let $f(z) = z + \sum_{n=2}^{\infty} a_n z^n \in S_0$ such that

$$\inf_{\substack{t \in \mathbb{R} \\ z \in \Delta}} \left| \frac{f * h_T(z)}{z} \right| \geq \delta > 0.$$

Then $g(z) = z + \sum_{n=2}^{\infty} b_n z^n \in S_0$ provided that $\sum_{n=2}^{\infty} n|a_n - b_n| \leq \delta$.

Note [6] that the conclusion of Theorem B holds for any $f \in K \subset S_0$ with $\delta = \frac{1}{4}$.

PROOF OF THEOREM 1

As noticed above it is sufficient to prove Theorem 1 for $f(z) = z/(1-z)$. For sequences $\{n_k\}_{k=2}^{\infty}$ satisfying $n_2 = 2$ and $n_3 > 3$ our approach is to prove that $v(z) = z + \sum_{k=2}^{\infty} c^{n_k-1} z^{n_k} \in K$ by showing that it either satisfies the condition of Alexander's Theorem or the more general Theorem A with $u(z) = z/(1-cz)$. As already mentioned, if $\{n_k\}_{k=2}^{\infty} = \{2k-2\}_{k=2}^{\infty}$ then we obtain the extremal function $z + \sum_{k=2}^{\infty} z^{2(k-1)} = z + z^2/(1-z^2)$.

For the remaining sequences $\{n_k\}_{k=2}^{\infty}$ we now show that $z + \sum_{k=2}^{\infty} z^{n_k}$ is convex univalent in $|z| < \frac{1}{4}$ by using Theorem A with an appropriate choice of the function $u(z)$. If $n_2 > 2$, then

$$(1) \quad z + \sum_{k=2}^{\infty} \frac{1}{4^{n_k-1}} z^{n_k} \in K$$

because

$$\sum_{k=2}^{\infty} \frac{n_k^2}{4^{n_k-1}} \leq \sum_{j=3}^{\infty} \frac{j^2}{4^{j-1}} = \frac{104}{108} < 1.$$

Now consider the function $u(z) = z/(1-z/4)$. A simple calculation gives, for all real T and $z \in \Delta$,

$$\left| \frac{zu'(z) * h_T(z)}{z} \right| = |h'_T(z/4)| \geq \frac{1 - |z|/4}{(1 + |z|/4)^3} > \frac{48}{125} = .384.$$

Since

$$u(z) \equiv z + \frac{1}{4}z^2 + \frac{1}{4^2}z^3 + \frac{1}{4^3}z^4 + \sum_{j=5}^{\infty} \frac{1}{4^{j-1}}z^j,$$

we now obtain from Theorem A

$$(2) \quad v(z) \equiv z + \frac{1}{4}z^2 + \frac{1}{4^2}z^3 + \frac{1}{4^3}z^4 + \sum_{k=5}^{\infty} \frac{1}{4^{n_k-1}}z^{n_k} \in K$$

because, since $n_5 \geq 5$,

$$\sum_{n=2}^{\infty} n^2 \left| \frac{u^{(n)}(0)}{n!} - \frac{v^{(n)}(0)}{n!} \right| \leq \sum_{j=5}^{\infty} \frac{j^2}{4^{j-1}} < .151 < .384.$$

In a similar way if

$$u(z) \equiv \frac{z}{1 - z/4} - \frac{1}{4^3} z^4,$$

we compute

$$\left| \frac{zu'(z) * h_T(z)}{z} \right| = \left| h'_T(z/4) - \frac{1}{4^2} \frac{4 + iT}{1 + iT} z^3 \right| \geq .384 - .250 = .134$$

and, therefore,

$$(3) \quad v(z) \equiv z + \frac{1}{4} z^2 + \frac{1}{4^2} z^3 + \frac{1}{4^4} z^5 + \sum_{k=6}^{\infty} \frac{1}{4^{n_k-1}} z^{n_k} \in K,$$

since

$$(4) \quad \sum_{n=2}^{\infty} n^2 \left| \frac{u^{(n)}(0)}{n!} - \frac{v^{(n)}(0)}{n!} \right| \leq \sum_{j=6}^{\infty} \frac{j^2}{4^{j-1}} < .053 < .134.$$

Finally let $u(z) \equiv z + \frac{1}{4} z^2 + \frac{1}{16} z^3$. Because, for any $z \in \Delta$,

$$\inf_T \left| \frac{zu'(z) * h_T(z)}{z} \right| = \frac{|2 + \frac{3}{2}z + \frac{3}{4}z^2| - |\frac{1}{2}z + \frac{3}{8}z^2|}{2}$$

we have

$$\left| \frac{zu'(z) * h_T(z)}{z} \right| \geq \frac{|2 + \frac{3}{2}z + \frac{3}{4}z^2|^2 - |\frac{1}{2}z + \frac{3}{8}z^2|^2}{2(2 + \frac{3}{2} + \frac{3}{4} + \frac{1}{2} + \frac{3}{8})}.$$

Setting $z = e^{i\theta}$, we see that the numerator of this last expression is $219/64 + (63/8) \cos \theta + 6 \cos^2 \theta$, which attains its minimum when $\cos \theta = -21/32$. Thus,

$$\left| \frac{zu'(z) * h_T(z)}{z} \right| \geq \frac{429/512}{41/4} > .08 > .053.$$

We get from Theorem A and (4),

$$(5) \quad v(z) \equiv z + \frac{1}{4} z^2 + \frac{1}{16} z^3 + \sum_{k=6}^{\infty} \frac{1}{4^{n_k-1}} z^{n_k} \in K.$$

Now it should be clear (see (1), (2), (3), (5)) that the theorem is valid (even with the constant c replaced by $\frac{1}{4}$) for any sequence $\{n_k\}_{k=2}^{\infty}$ except possibly for some sequences with $n_2 = 2$ and $n_3 \geq 4$. The rest of the proof consists of proving $z + cz^2 + \sum_{n_k \geq 4} c^{n_k-1} z^{n_k} \in K$.

We need the following inequalities:

$$(6) \quad \sum_{n=1}^m (2n)^2 c^{2n-1} + \sum_{n=2m+3}^{\infty} n^2 c^{n-1} \leq 1, \quad m \geq 1.$$

$$(7) \quad \sum_{n=2}^m (2n-1)^2 c^{2n-2} + \sum_{n=2m+2}^{\infty} n^2 c^{n-1} \leq \frac{1-c}{(1+c)^3}, \quad m \geq 2.$$

Since $\sum_{n=1}^{\infty} (2n)^2 c^{2n-1} = 1$, the inequality (6) will follow if we can show that the left-hand side is an increasing function of m . This is equivalent to

$$\sum_{n=1}^m (2n)^2 c^{2n-1} + \sum_{n=2m+3}^{\infty} n^2 c^{n-1} \leq \sum_{n=1}^{m+1} (2n)^2 c^{2n-1} + \sum_{n=2m+5}^{\infty} n^2 c^{n-1}$$

or

$$(2m+3)^2 c + (2m+4)^2 c^2 \leq (2m+2)^2, \quad m \geq 1.$$

Now the result follows because

$$1 - \left(\frac{2m+3}{2m+2} \right)^2 c - \left(\frac{2m+4}{2m+2} \right)^2 c^2$$

is minimized when $m = 1$ and

$$1 - \left(\frac{5}{4} \right)^2 c - \left(\frac{3}{2} \right)^2 c^2 > 0.$$

It is easily established that $\sum_{n=2}^{\infty} (2n-1)^2 c^{2n-2} = (1-c)/(1+c)^3$ and (7) will follow if we can show that the left-hand side is an increasing function of m . This amounts to

$$(2m+2)^2 c + (2m+3)^2 c^2 \leq (2m+1)^2, \quad m \geq 2.$$

This last statement is valid because

$$1 - \left(\frac{2m+2}{2m+1} \right)^2 c - \left(\frac{2m+3}{2m+1} \right)^2 c^2$$

is minimized when $m = 2$ and

$$1 - \left(\frac{6}{5} \right)^2 c - \left(\frac{7}{5} \right)^2 c^2 > 0.$$

Clearly we obtain by (6) and Alexander's theorem, for all $m \geq 1$,

$$(8) \quad f(z) \equiv z + \sum_{n=1}^m c^{2n-1} z^{2n} + \sum_{n_k \geq 2m+3} c^{n_k-1} z^{n_k} \in K.$$

Now let us consider

$$\begin{aligned} u(z) &= \frac{z}{1-cz} \\ &= z + \sum_{n=1}^m c^{2n-1} z^{2n} + c^{2m} z^{2m+1} + \sum_{n=2}^m c^{2n-2} z^{2n-1} + \sum_{j=2m+2}^{\infty} c^{j-1} z^j. \end{aligned}$$

We compute for all $z \in \Delta$ and $T \in \mathbf{R}$,

$$(9) \quad \left| \frac{zu'(z) * h_T(z)}{z} \right| = |h'_T(cz)| \geq \frac{1 - c|z|}{(1 + c|z|)^3} > \frac{1 - c}{(1 + c)^3}.$$

It then follows from (7), (9), and Theorem A that

$$(10) \quad v(z) = z + \sum_{n=1}^m c^{2n-1} z^{2n} + c^{2m} z^{2m+1} + \sum_{n_k \geq 2m+2} c^{n_k-1} z^{n_k} \in K$$

for all $m \geq 1$.

We now can complete our proof of Theorem 1. As noticed before we need to prove that

$$f(z) = z + z^2 + \sum_{n_k \geq 4} z^{n_k}$$

is convex univalent in $|z| < c$. Consider an increasing sequence of integers $\{n_k\}$ where $n_1 \geq 4$ and define $2j+1 \geq 5$ as the smallest odd number in this sequence (if $2j+1$ does not exist, our counterexample to the conjecture shows clearly that f is convex univalent in $|z| < c$). Then either

$$f(z) = z + z^2 + \sum_{k=1}^N z^{2t_k} + z^{2j+1} + \sum_{n_k \geq 2j+2} z^{n_k}$$

for some sequence $2 \leq t_1 < t_2 < \dots < t_N \leq j$, $1 \leq N \leq j-1$, or

$$f(z) = z + z^2 + z^{2j+1} + \sum_{n_k \geq 2j+2} z^{n_k}.$$

In the second case f is convex in $|z| < c$, by (8) with $m = 1$. In the first case a finite number of applications of (8) with $m = 2, \dots$ show that we can assume $t_1 = 2$, $t_2 = 3, \dots, t_N = j$, that is we need to prove

$$f(z) = z + \sum_{n=1}^j z^{2n} + z^{2j+1} + \sum_{n_k \geq 2j+2} z^{n_k}$$

to be convex in $|z| < c$. But this follows from (10) with $m = j$. Our proof of Theorem 1 is now complete.

PROOF OF THEOREM 2

It suffices to show that $g(z) = z + \sum_{k=2}^{\infty} b^{n_k-1} z^{n_k} \in S_0$. If $n_2 \geq 3$, then $\sum_{k=2}^{\infty} n_k b^{n_k-1} \leq \sum_{n=3}^{\infty} n b^{n-1} < 1$, a well-known sufficient condition for g to be in S_0 . If $n_2 = 2$, $n_3 = 3$, and $n_4 = 4$, we apply Theorem B with $f(z) = z/(1 - bz) \in K$ and $\delta = \frac{1}{4}$. If $n_2 = 2$, $n_3 = 3$, and $n_4 > 4$, we can apply

Theorem B with $f(z) = z/(1 - bz) - b^3 z^4$ and $\delta = \frac{1}{4}$ because

$$\begin{aligned} \left| \frac{f * h_T(z)}{z} \right| &= \left| \frac{h_T(bz)}{bz} - \frac{b^3(4 + iT)}{1 + iT} z^3 \right| \\ &\geq \left| \frac{h_T(bz)}{bz} \right| - 4b^3 |z|^3 \\ &> \frac{1}{(1+b)^2} - 4b^3 > \frac{1}{4}. \end{aligned}$$

In both cases we obtain $g \in S_0$ because

$$\sum_{n=2}^{\infty} n \left| \frac{f^{(n)}(0)}{n!} - \frac{g^{(n)}(0)}{n!} \right| \leq \sum_{n=5}^{\infty} nb^{n-1} < .17 < 1/4.$$

Thus Theorem 2 is valid except possibly when $n_2 = 2$ and $n_3 \geq 4$.

The remainder of the proof follows along the lines of that of Theorem 1, with appropriate modifications that include substituting b for c , Theorem B for Theorem A, and replacing (6) and (7) with

$$(6') \quad \sum_{n=1}^m 2nb^{2n-1} + \sum_{n=2m+3}^{\infty} nb^{n-1} \leq 1, \quad m \geq 1.$$

$$(7') \quad \sum_{n=2}^m (2n-1)b^{2n-2} + \sum_{n=2m+2}^{\infty} nb^{n-1} \leq \frac{1}{(1+b)^2}, \quad m \geq 2.$$

REFERENCES

1. J. W. Alexander, *Functions which map the interior of the unit circle upon simple regions*, Ann. of Math. **17** (1915/16), 12-22.
2. P. L. Duren, *Univalent functions*, Springer-Verlag, New York, 1983.
3. R. Fournier, *On neighbourhoods of univalent starlike functions*, Ann. Polon. Math. **XLVII** (1986), 189-202.
4. A. W. Goodman and I. J. Schoenberg, *On a theorem of Szegő on univalent convex maps of the unit circle*, J. Analyse Math. **44** (1984/85), 200-204.
5. St. Ruscheweyh, *On the radius of univalence of the partial sums of convex functions*, Bull. London Math. Soc. **4** (1972), 367-369.
6. —, *Neighborhoods of univalent functions*, Proc. Amer. Math. Soc. **31** (1981), 521-527.
7. St. Ruscheweyh and T. Sheil-Small, *Hadamard products of schlicht functions and the Pólya-Schoenberg conjecture*, Comment. Math. Helv. **48** (1973), 119-135.
8. H. Silverman, *Radii problems for sections of convex functions*, Proc. Amer. Math. Soc. **104** (1988), 1191-1196.
9. G. Szegő, *Zur theorie der schlichten Abbildungen*, Math. Ann. **100** (1928), 188-211.

DÉPARTEMENT DE MATHÉMATIQUES ET DE STATISTIQUE, UNIVERSITÉ DE MONTRÉAL, C. P. 6128, SUCCURSALE A, MONTRÉAL, QUEBEC H3C 3J7 CANADA

DEPARTMENT OF MATHEMATICS, COLLEGE OF CHARLESTON, CHARLESTON, SOUTH CAROLINA 29424