ADDITIVE PROPERTIES OF MULTIPLICATIVE SUBGROUPS OF FINITE INDEX IN FIELDS

PEDRO BERRIZBEITIA

(Communicated by Louis J. Ratliff, Jr.)

ABSTRACT. Gallai's theorem, an n -dimensional generalization of Van der Waerden's theorem on arithmetic progression, is used to prove the following theorem:

Let F be a field and $G \subseteq F^*$ a subgroup of finite index n . There is a positive integer N , which depends only on n , so that if $\text{Char } F = 0$ or $\text{char } F \geq N$, then $G + G = F$.

INTRODUCTION

It is well known that if n is a positive integer and p is a prime number, then the equation $x^n + y^n \equiv t \pmod{p}$ has a solution for any t as long as p is sufficiently large. This problem can be solved by using Jacobi Sums to approximate the number of solutions of the equation. This is done in [3, Chapter 8]. The problem can be restated as follows: If G is a subgroup of F_p^* of index at most n , then, for large enough values of p , $G + G \supseteq F_p^*$, with equality if and only if $-1 \notin G$.

For the case $n = 3$ Leep and Shapiro [5] were able to prove the following generalization. Let G be a subgroup of index 3 in the multiplicative group F^* of a field F . Then $G + G = F$, except in the cases $|F| = 4, 7, 13$, or 16. They also conjectured that the conclusion holds for infinite fields if the index is 5.

In §1 of this paper we prove Theorem 1.1 which states: *Let F be a field and $G \supseteq F^*$ a subgroup of finite index n . There is a positive integer N , that depends only on n , so that if $\text{Char } F = 0$ or $\text{char } F \geq N$, then, $G + G = F$.*

Fields for which the conclusion of Theorem 1.1 holds will be called *uniform*. It follows that if F is a uniform field, then, $G + G = F$ if and only if $-1 \in G$. This implies, in particular, that the above conjecture is true for uniform fields. More generally, we show that if m is a positive integer and $m \times G = \{g_1 + g_2 + \cdots + g_m \mid g_i \in G\}$ then $m \times G = F$ if and only if $-1 \in m - 1 \times G$ or, equivalently, if and only if $0 \in m \times G$ (see Corollary 1.2 below). In §2 we let F be an arbitrary field and $G \subseteq F^*$ a subgroup of index n in F^* . Let P denote the additive closure of G in F . We show that if F is uniform, then

Received by the editors February 23, 1990 and, in revised form, May 7, 1990.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 11P99; Secondary 12E99.

©1991 American Mathematical Society
 0002-9939/91 \$1.00 + \$.25 per page

$P = m \times G$ for some $m \leq [P : G] + 1$. (See Corollary 2.2.) Finally, in §3 we let $\mathbf{Q}$ denote the field of the rational numbers. For every odd prime $p \geq 5$, we construct a subgroup of index $n = p - 1$ in $\mathbf{Q}^*$, such that $n + 1 \times G = \mathbf{Q}$ but $m \times G \neq \mathbf{Q}$ for all $m \leq n$.

1. UNIFORM FIELDS

We follow Leep and Shapiro's notation. Let F be a field. Let $G \subseteq F^*$ be a subgroup of finite index n . For each positive integer m we denote by $m \times G$ the set of sums of m elements of G ; that is, $m \times G = \{g_1 + g_2 + \cdots + g_m \mid g_i \in G\}$. The additive closure of G will be denoted by $\sum G$ or simply P . So $P = \bigcup_{i=1}^{\infty} i \times G$. P_m will denote the partial union $P_m = \bigcup_{i=1}^m i \times G$. Finally let $G - G = \{g_1 - g_2 \mid g_i \in G\}$. We now state the main theorem of this paper.

Theorem 1.1. *Let F be a field and $G \subseteq F^*$ a subgroup of finite index n . There is a positive integer N , which depends only on n , so that if $\text{Char } F = 0$ or $\text{char } F \geq N$, then, $G - G = F$.*

For simplicity, fields for which the conclusion of Theorem 1.1 holds will be called *uniform* with respect to subtraction or simply uniform fields.

Proof of Theorem 1.1. We make some preliminary remarks. Note first that $G - G = F$ if and only if $cG - cG = F$ for any $c \in F^*$. Also note that $cG - cG$ consists of a union of cosets of $F^* \bmod G$ and of 0. Hence, to prove that $G - G = F$, it is enough to show that $cG - cG$ contains a complete set of representatives of $F^* \bmod G$ for some $c \in F$.

Let $\mathbf{N}$ be the set of nonnegative integers. For $m \in \mathbf{N}$, m positive, let $\mathbf{N}^m = \{(x_1, x_2, \dots, x_m) \mid x_i \in \mathbf{N}\}$. We will use the following result due to Gallai:

Theorem G. *Let $V \subseteq \mathbf{N}^m$ be finite. For any finite coloring of $\mathbf{N}^m$ there is $a \in \mathbf{N}^m$ and nonzero $d \in \mathbf{N}$, so that all the points of $a + dV$ have the same color.*

Gallai's theorem is a result of Ramsey theory. It is derived from the Hales-Jewett theorem in [2]. To show the existence of $N(n)$ in Theorem 1.1, we will use the following apparently stronger but equivalent form of Gallai's theorem. For $s \in \mathbf{N}$ let $[s] = \{0, 1, 2, \dots, s\}$.

Theorem G'. *Let $V \subseteq \mathbf{N}^m$ be finite. Let $m \in \mathbf{N}$ be positive. There is a natural number $N' = N'(V, m, r)$, such that for any $N \geq N'$ and any r -coloring of $[N]^m$, there is $a \in [N']^m$ and a nonzero $d \in \mathbf{N}$, $d \leq N'$, such that $a + dV$ is monochromatic.*

That Theorem G' is equivalent to Theorem G is a consequence of König's infinity lemma. This modification was first introduced by O. Schreier for the case $m = 1$. See [6].

In our situation we assume $m = n$, the index of G in F^* . $V \in \mathbf{N}^n$ will be the set $V = \{0, e_1, e_2, \dots, e_n\}$, where the e_i are the vectors of the canon-

ical basis for $\mathbf{R}^n$. To each one of the cosets of $F^* \bmod G$ we associate a different color and to $0 \in F$ we associate another color. We will induce a $(n+1)$ -coloring on the set $\mathbf{N}^n$ as follows: Let $g_1, g_2, \dots, g_n$ be a set of representatives of $F^* \bmod G$. Paint each $(x_1, x_2, \dots, x_n) \in \mathbf{N}^n$ with the color associated to the class of $x_1 g_1 + x_2 g_2 + \dots + x_n g_n$. By Gallai's theorem, there is $a = (a_1, a_2, \dots, a_n) \in \mathbf{N}^n$ and a nonzero $d \in \mathbf{N}$ such that $a + dV$ is monochromatic. It follows that if $c = a_1 g_1 + a_2 g_2 + \dots + a_n g_n$, then c and $c + d g_i$ have the same color associated. Hence $d g_i \in cG - cG$ for all $i = 1, 2, \dots, n$. If $d \neq 0$ in F the $d g_i$ run through all classes of $F^* \bmod G$. In this case, by the preliminary remarks we get $G - G = F$. Clearly d is nonzero in F when F has characteristic zero. Also, since $d \leq N' = N'(V, n, n+1)$ given by Theorem G' the same conclusion holds for fields of characteristic larger than N' . Since V depends only on n we let $N(n) = N'$. $\square$

Remark. Following the same techniques used in [1] to derive a proof of Van der Waerden's theorem from the Hales-Jewett theorem, one can derive from Hales-Jewett a particular case of Gallai's theorem that is enough to prove Theorem 1.1. If this is done, one can show that $N(n)$ can be chosen to be no larger than 2^M where $M = N(n+1, 2^n)$ is the number given in the Hales-Jewett theorem. The numbers $N(r, t)$ are defined recursively and, therefore are computable, however, they are already extremely large for small values of t and r .

Corollary 1.2. (a) *Let F be a uniform field. If $m > 1$, then $m \times G = F$ if and only if $-1 \in m-1 \times G$. In particular $G + G = F$ if and only if $-1 \in G$.*

(b) *If G has odd index in F^* for some uniform field F , then $G + G = F$.*

Proof. If $m \times G = F$ then $g_1 + g_2 + \dots + g_m = 0$ for some $g_i \in G$. Then $-1 = g_2 g_1^{-1} + g_3 g_1^{-1} + \dots + g_m g_1^{-1} \in m-1 \times G$. For the converse, if $-1 \in m-1 \times G$ then $-G \subseteq m-1 \times G$. Using Theorem 1.1 we have: $F = G - G = G + (-G) \subseteq G + m-1 \times G = m \times G$. This proves (a). For (b) note that if $[F^* : G] = 2k+1$ then $(-1)^{2k+1} = -1 \in G$. The result follows from (a). (The case $n = 5$ was conjectured in [5] for arbitrary finite fields.) $\square$

Remark. Part (a) of Corollary 1.2 could have been stated in the following equivalent form: *If F is a uniform field, then $m \times G = F$ if and only if $0 \in m \times G$.*

2. INEQUALITIES CONCERNING SETS OF SUMS OF ELEMENTS OF G

Let F be an arbitrary field and $G \subseteq F^*$ be a multiplicative subgroup of finite index. Lemma 1 of [5] shows that if $-1 \in P$ then $P = F$. In any case, we have the following:

Proposition 2.1. *If G has index n in F^* we have:*

- (a) *For $i \geq 1$, $P_i \subseteq P_{i+1}$ with equality if and only if $P_i = P$.*
- (b) *If $-1 \in P$ then $P_{n+1} = F$. If $-1 \notin P$ then $P_{(n/2)+1} = P$.*

Proof. $P_i \subseteq P_{i+1}$ by definition of P_i . If $P_i = P$ then $P_{i+1} = P_i + G = P + G = P = P_i$, so $P_{i+1} \subseteq P_i$. Conversely, if $P_i = P_{i+1}$ then $P_i = P_r$ for all $r > i$.

But $P = \bigcup_{r=1}^{\infty} P_r = P_i$. For (b) just note that every P_m is a union of cosets of F^* and possibly of $\{0\}$. For simplicity $\{0\}$ is considered to be a coset. Suppose that $-1 \in P$ and that $P_n \neq F$. Then, by part (a), for all $i \leq n$, P_{i+1} contains at least one class more than P_i . It follows that P_{n+1} contains all classes including $\{0\}$. So $P_{n+1} = F$. If $-1 \notin P$, $[F^* : P] \geq 2$ then $[P : G] \leq n/2$ and the argument is the same as before. $\square$

Corollary 2.2. *If F is a uniform field we have:*

- (a) $i \times G \subseteq i + 1 \times G$ with equality if and only if $i \times G = P$.
- (b) If $-1 \in P$ then $n + 1 \times G = F$. If $-1 \notin P$ then $n/2 + 1 \times G = P$.

Proof. Using Proposition 3.1 we see that it is enough to show that $i \times G = P_i$ for any i . For this we note that it suffices to show that $G \subseteq G + G$. But, since F is uniform, then $G - G = F$. So $G - G \supseteq G$. This means that there are elements g_1, g_2, g_3 in G such that $g_1 - g_2 = g_3$. So $g_1 = g_2 + g_3$, which implies $G + G \subseteq G$.

Remark. In general it is not true that $n + 1 \times G = P$. Let $F = F_7$, the finite field with 7 elements and let $G = \{1, -1\}$. Then $[F^* : G] = 3$ and $4 \times G \neq F$.

3. BOUNDS IN §2 CANNOT BE IMPROVED

We end this paper with examples of subgroups of finite index n in $\mathbf{Q}^*$ for which $n+1$ is the minimum r such that $\mathbf{Q} = r \times G$. We note that subgroups of finite index in $\mathbf{Q}^*$ are kernels of multiplicative maps of $\mathbf{Q}^*$ into finite groups. These maps are determined by the image of the prime numbers and by the image of -1 .

Let q be an odd prime $q \geq 5$. We will map $\mathbf{Q}^*$ into F_q^* as follows: if $p \neq q$ is a prime then let p be mapped to its class mod q . Let the prime q be mapped into 1 and let -1 be mapped to its class mod q . This way every rational number which is a unit in the field of the q -adic number is mapped into its class mod q . Hence, if G is the kernel of this map, we have, $n \in G$ if and only if $n = q^r u_1$, where r is an integer and u_1 is a unit, $u_1 \equiv 1 \pmod{q}$. It is then easy to verify that $q - 1 \times G = \mathbf{Q}^*$. But $q - 1$ is the index of G in $\mathbf{Q}^*$.

ACKNOWLEDGMENT

I am grateful to Professor D. B. Shapiro for telling me about Van der Waerden's n -dimensional generalization. My original version of Theorem 1.1 used the classical one-dimensional Van der Waerden's theorem and involved only the case $F = \mathbf{Q}$, the field of rational numbers.

REFERENCES

1. R. L. Graham, *Rudiments of Ramsey theory*, Regional Conf. Ser. in Math., vol. 45, Amer. Math. Soc., Providence, RI, 1979.
2. R. L. Graham, B. L. Rothschild, and J. H. Spencer, *Ramsey theory*, Wiley-Interscience, 1980.

3. K. Ireland and M. Rosen, *A classical introduction to modern number theory*, Springer-Verlag, Berlin, 1982.
4. A. Y. Khinchin, *Three pearls of number theory*, Rochester, NY, 1952, pp. 11–17.
5. D. B. Leep and D. B. Shapiro, *Multiplicative subgroups of index three in a field*, Proc. Amer. Math. Soc. **105** (1989), 802–807.
6. B. L. Van der Waerden, *How the proof of Baudet's conjecture was found*, Studies in Pure Mathematics (L. Mirsky, ed.), Academic Press, London, 1971, pp. 251–260.

DEPARTAMENTO DE MATEMÁTICAS PURAS Y APLICADAS, UNIVERSIDAD SIMÓN BOLIVAR,
APARTADO 89000, CARACAS 1080-A, VENEZUELA