THE SPACE $(l_\infty/c_0, \text{WEAK})$ IS NOT A RADON SPACE

JOSE L. DE MARIA AND BALTASAR RODRIGUEZ-SALINAS

(Communicated by Andrew M. Bruckner)

ABSTRACT. Talagrand [10] gives an example of a Banach space with weak topology which is not a Radon space, independently of their weight. This result gives an answer to a question formulated by Schwartz [9]. In this paper, following the papers of Drewnowski and Roberts [1] and Talagrand [10], we prove that the classical space $(l_\infty/c_0, \text{weak})$ is not a Radon space.

Introduction. A Hausdorff topological space E is said to be a Radon space if every finite Borel measure is a Radon measure; i.e.,

$$\mu(A) = \{\mu(K) : K \subset A, K \text{ compact}\}$$

for each Borel subset A of E .

We shall say that a cardinal α is of measure zero (resp. nonmeasurable) if there is not a real-valued, diffuse, nontrivial measure (resp. $\{0, 1\}$ -valued), on the power set of a set A with cardinal α .

The weight (density character) of a topological space E is the smallest cardinal such that there exists in E a dense subset A with this cardinal.

A topological space E has the α -property of Lindelöf, α a transfinite cardinal, if for each family $(G_i)_{i \in I}$ of open subsets of E there exists $J \subset I$ such that $\text{card}(J) \leq \alpha$ and $\bigcup_{i \in I} G_i = \bigcup_{i \in J} G_i$. The smallest cardinal α such that E has the α -property of Lindelöf is called the L -weight of E .

Likewise, E is a Flock space if for every well-ordered family $(G_i)_{i \in I}$ of open sets, with $H_\alpha = G_\alpha \setminus \bigcup_{\beta < \alpha} G_\beta$, every union $\bigcup_{\alpha \in J} H_\alpha$ ($J \subseteq I$) is universally Borel-measurable. This property is an extrapolation of the Montgomery lemma [7], which proves that every metrizable space is Flock. Also, the strongly Lindelöf spaces are Flock spaces.

Let I be a noncountable set and $x \notin I$. Consider the topology T on $I \cup \{x\}$ such that the discrete topology is on I and the neighborhoods of x are the complementary sets of countable subsets of I . Let K be the Stone-Čech compactification of this space. Then, as Talagrand [10] proved, $(C(K), \text{weak})$ is not a Radon space.

Received by the editors February 28, 1990.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 28C15.

This research was supported in part by DGICYT PB88-0141.

Talagrand also proved that (Z, weak) is not a Radon space when Z is the subspace of $l^\infty(I)$ of the countable support functions and $\text{card}(I) = \aleph_1$. Following that proof and the work of Drewnowski and Roberts [1], this paper is devoted to proving that the classical space $(l_\infty/c_0, \text{weak})$ is not a Radon space. Note that this fact could be proved if Z were a subspace of l_∞/c_0 , but this is not yet known. Let us remark that it is easily proved that (E, weak) is not a Radon space if the weight of E is real-measurable.

It is easily deduced that every Borel measure on (l_∞, weak) valued on $\{0, 1\}$ is a δ_x .

In Theorem 18 [2] we proved that a Banach space with the weak topology is a Radon space if and only if it is Flock and its L -weight is of measure zero. Thus, in the last part of this paper, we prove that l_∞/c_0 is not a Flock space. Conversely, in [4], we improved Schachermayer's theorem, proving that a subset Ω of a WCG Banach space with the weak topology whose weight is of measure zero is a Radon space of type $(\mathcal{F})$; that is, every finite Borel measure on (Ω, weak) is τ -additive.

1. Theorem. *On $(l_\infty/c_0, \text{weak})$ there exists a Borel measure $\mu \neq 0$ valued on $\{0, 1\}$ such that the space is not a Radon measure.*

Proof. Let $\beta\omega$ be the Stone-Ćech compactification of $\omega = \mathbb{N}$ with discrete topology. Let $\mathcal{A}$ be the clopen algebra of $K = \beta\omega/\omega$ and $\mathcal{A}_0 = \mathcal{A} \setminus \{\emptyset\}$. $\mathcal{A}$ has the following property (called the Cantor separability property in [11]): For every decreasing sequence (A_n) in $\mathcal{A}_0$ there exists an $A \in \mathcal{A}_0$ such that $A \subset \bigcap_{n=1}^\infty A_n$.

Let $\mathcal{V} = (V_\alpha)_{\alpha \in A}$, a maximal well-ordered family in $\mathcal{A}_0$. Then every countable intersection $\bigcap_{n=1}^\infty V_{\alpha_n}$ contains a $V \in \mathcal{V}$. We denote by F the set of continuous functions from K in $\{0, 1\}$ which vanish in $H = \bigcap \{V : V \in \mathcal{V}\}$.

Like Talagrand, we are going to make two classes $\mathcal{C}$ and $\mathcal{D}$ of weak-Borel sets (Borel with the weak topology) in $F \subset C(K)$ such that

- (i) the smallest σ -algebra containing $\mathcal{C}$ is the class $\mathcal{B}$ of the weak-Borel sets of F ;
- (ii) if $C \in \mathcal{C}$, then either $C \in \mathcal{D}$ or $F \setminus C \in \mathcal{D}$;
- (iii) every countable intersection of elements of $\mathcal{D}$ is not empty; and
- (iv) for every $t \in K \setminus H$, $\{f \in F : f(t) = 1\} \in \mathcal{D}$.

Now, we define a Borel measure μ on (F, weak) by $\mu(B) = 1$ if $B \in \mathcal{B}$ and contains a countable intersection of members of $\mathcal{D}$, and $\mu(B) = 0$ in the other case. The measure μ is not τ -additive, because F is the union of the open sets $G_t = \{f \in F : f(t) = 0\}$ when $t \in K \setminus H$, $\mu(F) = 1$, and each $\mu(G_t) = 0$. It follows that there exists a Borel measure μ_0 on $(C(K), \text{weak}) \approx (l_\infty/c_0, \text{weak})$ (see [6]) valued in $\{0, 1\}$ which is not a Radon measure.

Let k be an integer, J a set, $(\mu_j^p)_{j \in J, p \leq k}$, of Radon measures on K , and $(a^p)_{p \leq k}$, $(b^p)_{p \leq k}$ rational numbers such that $a^p < b^p$. By definition, $\mathcal{C}$ is the

class of sets $C = \bigcup_{j \in J} U_j$, where

$$U_j = \{f \in F : \forall p \leq k, \mu_j^p(f) \in (a^p, b^p)\},$$

varying over all k, J and measures μ_j^p .

The class $\mathcal{D}$ is built up by the sets $C \in \mathcal{E}$ such that, for every $V \in \mathcal{V}$, there exist $j \in J$ and $f \in U_j$ with $f = 1$ in $K \setminus V$, and with the complementaries $F \setminus C$ of the sets $C \in \mathcal{E}$ such that they do not have this condition. Only (iii) has to be proved. To do this, it is enough that, for every sequence $(C_n) \subset \mathcal{E} \cap \mathcal{D}$ and every $V_0 \in \mathcal{V}$, there exists $f \in \bigcap_n C_n$ such that $f = 1$ in $K \setminus V_0$. It is easily proved (with the evident notations) that for every $n \in \mathbb{N}$, there exists $\varepsilon_n > 0$ such that, if $W \in \mathcal{V}$, then there exist $j \in J_n$ and $f \in U'_j$ with $f = 1$ on $K \setminus W$, where

$$U'_j = \{f \in F : \forall p \leq k_n, \mu_{j,n}^p(f) \in (a_n^p + \varepsilon_n, b_n^p - \varepsilon_n)\}.$$

Let $\mathcal{U}_n$ be an ultrafilter on J_n which contains all sets

$$\{j \in J_n : \exists f \in U'_j, f = 1 \text{ on } K \setminus V\}$$

when $V \in \mathcal{V}$. Given $p \leq k_n$, let $v_n^p = \lim_{\mathcal{U}_n} \mu_{j,n}^p$. Then there exists $V \in \mathcal{V}$, $V \subset V_0$, such that $v_n^p(V \setminus W) = 0$ for every $W \in \mathcal{V}$ and $p \leq k_n$, $n \in \mathbb{N}$. Similarly, by means of an easy induction, there exist $f_n \in F$, $j_n \in J_n$, V_{α_n} , $V'_{\alpha_n} \in \mathcal{V}$ and clopen sets H_{n0} , H_{n1} in $\beta\omega$ such that

- (i) $\mu_{j_n,n}^p(V \setminus V_{\alpha_n}) \leq \varepsilon_n/2$, for every n and $p \leq k_n$;
- (ii) $f_n \in U'_{j_n}$ and $f_n = 1$ on $K \setminus V_{\alpha_n}$;
- (iii) $f_n = 0$ on $V'_{\alpha_n} \subset V_{\alpha_n}$ and $\mu_{j_n,n}^p(V'_{\alpha_n} \setminus H) = 0$ for every n and $p \leq k_n$;
- (iv)

$$\begin{aligned} A_{00} &= \emptyset, & A_{01} &= K \setminus V, \\ A_{n0} &= \{t \in V_{\alpha_n} \setminus V'_{\alpha_n} : f_n(t) = 0\}, \\ A_{n1} &= \{t \in V_{\alpha_n} \setminus V'_{\alpha_n} : f_n(t) = 1\}; \end{aligned}$$

- (v) H , H_{n0} , H_{n1} are pairwise disjoint sets such that $H_{ni} \supset A_{ni} \cup H_{n-1,i}$, $(H_{0i} = A_{0i})$ for $i = 0, 1$;
- (vi) The set $V_{\alpha_{n+1}}$ verifies that $V_{\alpha_{n+1}} \subset V'_{\alpha_n}$ and $V_{\alpha_{n+1}} \cap (H_{n0} \cup H_{n1}) = \emptyset$.

Clearly the open sets $G_0 = \bigcup_{n=1}^\infty H_{n0}$ and $G_1 = \bigcup_{n=1}^\infty H_{n1}$ are disjoint, but $\overline{G_0} \cap \overline{G_1} = \emptyset$ also is, because if $t \in \overline{G_0} \cap \overline{G_1}$, then $t \in \beta\omega$ would be a cluster point of two disjoint sets of integer numbers contained in G_0 and G_1 , respectively. As $(\overline{G_0} \cup \overline{G_1}) \cap H = \emptyset$, then there exists a clopen H_1 in K such that $\overline{G_1} \cap K \subset H_1$, $\overline{G_0} \cap K \cap H_1 = \emptyset$, and $H \cap H_1 = \emptyset$.

Function $f = \chi_{H_1} \in F$ satisfies $f = f_n$ on $A_n = A_{n0} \cup A_{n1}$. Moreover, since for every n and every $p \leq k_n$, we have that $\mu_{j_n}^p(f \neq f_n) < \varepsilon_n$. It follows that $f \in U_{j_n}$ for every n . Then $f \in \bigcap_{n=1}^\infty C_n$ and $f = 1$ on $K \setminus V_0$.

2. **Corollary.** $(l_\infty/c_0, \text{weak})$ is not a Radon space.

3. **Theorem.** Let Ω be a Flock space whose L -weight is nonmeasurable. Then every finite perfect Borel measure μ on Ω is τ -additive.

Proof. It is enough to prove that, for every family of open sets $(G_\alpha)_{\alpha \in A}$ in Ω ,

$$\mu \left(\bigcup_{\alpha \in A} G_\alpha \right) = \sup_J \mu \left(\bigcup_{\alpha \in J} G_\alpha \right),$$

where the supremum is taken over all finite subsets J in A .

From Zermelo's theorem, and since the L -weight of Ω is nonmeasurable, then we can suppose A is well ordered and cardinally nonmeasurable. Let $H_\alpha = G_\alpha \setminus \bigcup_{\beta < \alpha} G_\beta$. Since Ω is a Flock space, the union $\bigcup_{\alpha \in A} H_\alpha$ of every subfamily of $(H_\alpha)_{\alpha \in A}$ is a universally Borel-measurable set. By [5, Theorem 2.5], if $S = \{\alpha \in A : \mu(H_\alpha) \neq 0\}$, then S is countable and $\mu(\bigcup_{\alpha \notin S} H_\alpha) = 0$.

Now,

$$\mu \left(\bigcup_{\alpha \in A} G_\alpha \right) = \mu \left(\bigcup_{\alpha \in A} H_\alpha \right) = \mu \left(\bigcup_{\alpha \in S} H_\alpha \right) \leq \mu \left(\bigcup_{\alpha \in S} G_\alpha \right) \leq \sup_J \mu \left(\bigcup_{\alpha \in J} G_\alpha \right),$$

so $\mu(\bigcup_{\alpha \in A} G_\alpha) = \sup_J \mu(\bigcup_{\alpha \in J} G_\alpha)$.

4. **Corollary.** $(l_\infty/c_0, \text{weak})$ is not a Flock space.

Proof. Follows from Theorems 1 and 3, since cardinal of l_∞/c_0 is nonmeasurable.

Remark. Since [3, Theorem 6] says that every Radon space is a Flock space, Corollary 4 improves Corollary 2. Even this theorem proves that a regular space E is Flock and its weight is of measure zero if and only if E is a Radon space of type $(\mathcal{F})$.

5. **Theorem.** If (φ_n) is a sequence of continuous functions which separates points on a topological space E , then every Borel measure $\mu \neq 0$ on E which takes values in $\{0, 1\}$ is concentrated in a point.

Proof. Let ν be the Borel measure on $\mathbb{R}$ defined by

$$\nu(B) = \mu(\varphi_1^{-1}(B))$$

on the class $\mathcal{B}$ of Borel measure on $\mathbb{R}$. Then, as ν takes only the values 0 and 1, there exists $\alpha_1 \in \mathbb{R}$ such that $\mu(\varphi_1^{-1}(\alpha_1)) = 1$. Now, by induction, we can construct a sequence (α_n) in $\mathbb{R}$ such that

$$\mu \left(\bigcap_{k=1}^n \varphi_k^{-1}(\alpha_k) \right) = 1$$

for every n . To do this, it is enough to consider the following measure:

$$\nu(B) = \mu \left(\bigcap_{k=1}^n \varphi_k^{-1}(\alpha_k) \cap \varphi_{n+1}^{-1}(B) \right) \quad (B \in \mathcal{B}).$$

Then

$$\mu \left(\bigcap_{n=1}^{\infty} \varphi_n^{-1}(\alpha_n) \right) = 1.$$

Finally, as (φ_n) separates points of E , then $\bigcap_{n=1}^{\infty} \varphi_n^{-1}(\alpha_n)$ contains only a point x . So μ is concentrated in x .

6. **Corollary.** Every Borel measure $\mu \neq 0$ on (l_∞, weak) taking values in $\{0, 1\}$ is concentrated in a point.

This allows us to formulate the following open question: Is (l_∞, weak) a Flock space?

Remark. Drewnowski and Roberts [1] have proved that for every Schauder decomposition (finite or infinite) $C = l_\infty/c_0 = X_1 + X_2 + \dots$, at least one of X_n contains a space isomorphic to C , complemented in C . It can easily be proved, as in Theorem 5, that if C is a Banach space with a Schauder decomposition and there exists a nontrivial Borel measure μ on C valued in $\{0, 1\}$, then there exist $n \in \mathbb{N}$ and a nontrivial Borel measure μ_n on X_n taking values in $\{0, 1\}$. So, taking into account that $C \approx C \oplus C$ and $C \approx l_\infty \oplus C$ ($C = l_\infty/c_0$) the following question naturally appears:

Open question. If X is a complemented subspace in l_∞/c_0 on which there exists a nontrivial measure taking values in $\{0, 1\}$, then does X have a complemented copy of C ?

Let $L = L_\infty([0, 1]^c)$, where $c = 2^{\aleph_0}$ and $[0, 1]$ is considered with the product Lebesgue measure. If we suppose $c = \aleph_1(CH)$, then L is isometric to a complemented and closed subspace X of C .

Another question arises:

Open question. Does there exist on L a nontrivial Borel measure μ which only takes 0, 1 values?

REFERENCES

1. L. Drewnowski and J. W. Roberts, *On the primariness of the Banach space l_∞/c_0* , preprint.
2. J. de Maria and B. Rodriguez-Salinas, *On measures on Banach spaces with the weak topology* (to appear).
3. —, *On measurable sets of a τ -additive measure* (to appear).
4. —, *Banach spaces which are Radon spaces with the weak topology* (to appear).
5. G. Koumoullis, *On perfect measures*, Trans. Amer. Math. Soc. **264** (1981), 521–537.
6. I. E. Leonard and J. H. M. Whitfield, *A classical Banach space: l_∞/c_0* , Rocky Mountain J. Math. **13** (1981), 531–539.
7. D. Montgomery, *Non-separable metric spaces*, Fund. Math. **25** (1935), 527–533.
8. W. Schachermayer, *Eberlein compacts et espaces de Radon*, C. R. Acad. Sci. Paris **284** (1977), 405–407.
9. L. Schwartz, *Seminaire Maurey-Schwartz*, 1975–1976, expos XXIII.

10. M. Talagrand, *Sur un theoreme de L. Schwartz*, C. R. Acad. Sci. Paris **286** (1978), 265–267.
11. R. C. Walker, *The Stone-Čech compactification*, Springer-Verlag, New York, 1974.

DEPARTAMENTO MATEMATICAS FUNDAMENTALES, FACULTAD DE CIENCIAS, U.N.E.D., 28040 MADRID, SPAIN

DEPARTAMENTO DE ANALISIS MATEMATICO, FACULTAD DE MATEMATICAS, UNIVERSIDAD COMPLUTENSE, 28040 MADRID, SPAIN