HOMOTOPY-COMMUTATIVE H -SPACES

JAMES P. LIN AND FRANK WILLIAMS

(Communicated by Frederick R. Cohen)

ABSTRACT. Let X be an H -space with $H^*(X; \mathbb{Z}_2) \simeq \mathbb{Z}_2[x_1, \dots, x_d] \otimes \Lambda(y_1, \dots, y_d)$, where $\deg x_i = 4$ and $y_i = \text{Sq}^1 x_i$. In this article we prove that X cannot be homotopy-commutative. Combining this result with a theorem of Michael Slack results in the following theorem: *Let X be a homotopy-commutative H -space with mod 2 cohomology finitely generated as an algebra. Then $H^*(X; \mathbb{Z}_2)$ is isomorphic as an algebra over $A(2)$ to the mod 2 cohomology of a torus producted with a finite number of $CP(\infty)$ s and $K(\mathbb{Z}_2, 1)$ s.*

0. INTRODUCTION

In this article we prove the following theorem:

Theorem A. *Let X be an H -space with*

$$H^*(X; \mathbb{Z}_2) = \mathbb{Z}_2[x_1, \dots, x_d] \otimes \Lambda(y_1, \dots, y_d),$$

where $\deg x_i = 4$ and $y_i = \text{Sq}^1 x_i$. Then X cannot be homotopy-commutative.

The significance of Theorem A lies in its relationship to the following theorem, due to Michael Slack:

Theorem (Slack). *Let X be a homotopy-commutative H -space with mod 2 cohomology finitely generated as an algebra. Then*

- (1) *All even-degree generators have infinite height and are in degrees two and four.*
- (2) *All odd-degree generators lie in degrees one and five. The one-dimensional generators have infinite height and the five-dimensional generators are exterior.*
- (3) *$\text{Sq}^1: QH^4(X; \mathbb{Z}_2) \rightarrow QH^5(X; \mathbb{Z}_2)$ is an isomorphism.*

Received by the editors February 27, 1990.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 55P45, 55S40.

Key words and phrases. Homotopy-commutative H -space, cohomology operation, Steenrod algebra.

The first author's work was supported by a grant from the National Science Foundation.

Combining Theorem A with Slack's theorem results in the following:

Theorem B. *Let X be a homotopy-commutative H -space with mod 2 cohomology finitely generated as an algebra. Then $H^*(X; \mathbb{Z}_2)$ is isomorphic as an algebra over $A(2)$ to the mod 2 cohomology of a torus producted with a finite number of $CP(\infty)$ s and $K(\mathbb{Z}_2, 1)$ s.*

This theorem extends work of Hubbuck and Lin [3, 4] which proves any finite homotopy-commutative H -space has the homotopy type of a torus. Here we treat the larger class of H -spaces whose cohomology is finitely generated as an algebra.

Theorem A also has the following immediate corollary:

Corollary C. *The 3-connective cover of $S^3, \widehat{S}^3$, is not mod 2 homotopy-commutative, hence neither is any product of $\widehat{S}^3$ s homotopy-commutative.*

Remarks. (1) Corollary C can also be proved by using results in [8] and verifying that the Samelson product $\langle \alpha, \alpha \rangle \neq 0$, where α is a generator of $\pi_4(\widehat{S}^3)$. This approach was suggested to the authors by Chuck McGibbon. (The referee has observed that Corollary C can also be obtained quickly from results of Zabrodsky.)

(2) Our original proof of Theorem A rested upon an analysis of a partial Postnikov tower for X . The proof we give here is shorter and is based on a factorization of $Sq^8 Sq^4$ (Theorem 1.1 below) that may well be of independent interest.

(3) In the rest of this paper, cohomology will be taken to have coefficients in $\mathbb{Z}_2$ unless otherwise specified.

1. FACTORIZATION OF $Sq^8 Sq^4$

Theorem 1.1. *Let $x \in H^8(X; \mathbb{Z})$ be such that $\rho(x)$ lies in the kernel of Sq^2 , where $\rho: H^*(X; \mathbb{Z}) \rightarrow H^*(X; \mathbb{Z}_2)$ is reduction mod 2. Then there is a formula*

$$Sq^8 Sq^4 x = Sq^4 \tilde{\varphi}_{0,3}(x) + (Sq^9 + Sq^{7,2})\varphi_{1,1}(x) + x(Sq^4 x),$$

where the secondary operations $\tilde{\varphi}_{0,3}$ and $\varphi_{1,1}$ are defined by the following relations:

$$\tilde{\varphi}_{0,3}: Sq^9 = (Sq^7 + Sq^{4,2,1}) Sq^2 \text{ on an integral class}$$

$$\varphi_{1,1}: Sq^2 Sq^2 = 0 \text{ on an integral class.}$$

Proof. Consider the following two-stage Postnikov system.

$$\begin{array}{ccc} K(Z_2, 9) & & \\ \downarrow j & & \\ E & & \\ \downarrow p & & \\ K(Z, 8) & \xrightarrow{g} & K(Z_2, 10) \end{array}$$

in which $g^*(\iota_{10}) = \text{Sq}^2 \iota_8$. Then there are elements $\tilde{v}_{0,3} \in H^{16}(E)$ and $v_{1,1} \in H^{11}(E)$ such that

$$j^*(\tilde{v}_{0,3}) = (\text{Sq}^7 + \text{Sq}^{4,2,1})\iota_9$$

and

$$j^*(v_{1,1}) = \text{Sq}^2 \iota_9.$$

Further, $\bar{\Delta}(\tilde{v}_{0,3}) = p^*(\iota_8) \otimes p^*(\iota_8)$, where $\bar{\Delta}$ denotes the reduced diagonal. We compute

$$j^*((\text{Sq}^9 + \text{Sq}^{7,2})v_{1,1} + \text{Sq}^4 \tilde{v}_{0,3}) = 0$$

and

$$\bar{\Delta}((\text{Sq}^9 + \text{Sq}^{7,2})v_{1,1} + \text{Sq}^4 \tilde{v}_{0,3}) = \bar{\Delta}(p^*(\iota_8) \text{Sq}^4 p^*(\iota_8)).$$

Hence

$$z = (\text{Sq}^9 + \text{Sq}^{7,2})v_{1,1} + \text{Sq}^4 \tilde{v}_{0,3} - p^*(\iota_8) \text{Sq}^4 p^*(\iota_8)$$

is an element of $PH^{20}(E) \cap \ker(j^*)$. So $z \in p^*PH^{20}(K(Z, 8))$, and thus by [2] $z = c \text{Sq}^{8,4} p^*(\iota_8)$, for some $c \in Z_2$.

To prove the theorem, we must show that $c = 1$. To do this, we loop the diagram four times, obtaining

$$\begin{array}{ccc} K(Z_2, 5) & & \\ \downarrow j & & \\ \Omega^4 E & & \\ \downarrow \tilde{p} & & \\ K(Z, 4) & \xrightarrow{\tilde{g}} & K(Z_2, 6) \end{array}$$

Since $\text{Sq}^5 \iota_4 = \text{Sq}^{2,1} \text{Sq}^2 \iota_4$, there exists $\tilde{v}_{0,2} \in H^8(\Omega^4 E)$ such that $\tilde{j}^*(\tilde{v}_{0,2}) = \text{Sq}^{2,1} \iota_5$ and $\bar{\Delta}(\tilde{v}_{0,2}) = \tilde{p}^*(\iota_4) \otimes \tilde{p}^*(\iota_4)$. So $\bar{\Delta}(\text{Sq}^4 \tilde{v}_{0,2}) = \bar{\Delta}(\tilde{p}^*(\iota_4)^3)$. Thus

$$\begin{aligned} \text{Sq}^4 \tilde{v}_{0,2} + \tilde{p}^*(\iota_4)^3 + (\sigma^*)^4(\tilde{v}_{0,3}) &\in PH^{12}(\Omega^4 E) \cap \ker(\tilde{j}^*) \\ &\subset \tilde{p}^*PH^{12}(K(Z, 4)) = 0. \end{aligned}$$

We have

$$\begin{aligned}
 c\tilde{p}^*(\iota_4)^4 &= c\operatorname{Sq}^{8,4}\tilde{p}^*(\iota_4) \\
 &= (\operatorname{Sq}^9 + \operatorname{Sq}^{7,2})(\sigma^*)^4 v_{1,1} + \operatorname{Sq}^4(\sigma^*)^4 \tilde{v}_{0,3} \\
 &= \operatorname{Sq}^{7,2}(\sigma^*)^4 v_{1,1} + \operatorname{Sq}^4(\sigma^*)^4 \tilde{v}_{0,3} \\
 &= \operatorname{Sq}^{7,2}(\sigma^*)^4 v_{1,1} + \operatorname{Sq}^4 \operatorname{Sq}^4 \tilde{v}_{0,2} + \operatorname{Sq}^4 \tilde{p}^*(\iota_4)^3.
 \end{aligned}$$

Let $f: K(Z, 2) \rightarrow K(Z, 4)$ be defined by $f^*(\iota_4) = \iota_2^2$. Let $\hat{f}: K(Z, 2) \rightarrow \Omega^4 E$ be a lifting of f . Then

$$\begin{aligned}
 c\iota_2^8 &= \hat{f}^*(\operatorname{Sq}^{7,2}(\sigma^*)^4 v_{1,1} + \operatorname{Sq}^4 \operatorname{Sq}^4 \tilde{v}_{0,2} + \operatorname{Sq}^4 \tilde{p}^*(\iota_4)^3) \\
 &= 0 + \lambda \operatorname{Sq}^{4,4} \iota_2^4 + \operatorname{Sq}^4 \iota_2^6, \quad \lambda = 0 \text{ or } 1 \\
 &= \iota_2^8.
 \end{aligned}$$

So $c = 1$. $\square$

2. RELATIONS IN P_2X

Henceforth, we will assume that X is an H -space with mod 2 cohomology,

$$H^*(X) = Z_2[x_1, \dots, x_d] \otimes \Lambda(y_1, \dots, y_d)$$

as in the statement of Theorem A. For degree reasons, $H^*(X)$ is a primitively generated Hopf algebra.

According to [1], there is an exact triangle

$$\begin{array}{ccc}
 H^*(P_2X) & \xrightarrow{\iota} & \overline{H}^*(X) \\
 (2.1) \quad \swarrow \lambda & & \nwarrow \overline{\Delta} \\
 & \overline{H}^*(X) \otimes \overline{H}^*(X) &
 \end{array}$$

in which ι has degree -1 , λ has degree 2, and $\overline{\Delta}$ has degree 0.

Therefore there exist unique elements $u_i \in H^5(P_2X)$ and $v_i \in H^6(P_2X)$ such that $\iota(u_i) = x_i$, $\iota(v_i) = y_i$, and $\operatorname{Sq}^1 u_i = v_i$. A simple calculation following [1] shows that

$$(2.2) \quad \frac{Z_2[u_1, \dots, u_d, v_1, \dots, v_d, \operatorname{Sq}^4 u_1, \dots, \operatorname{Sq}^4 u_d]}{(\dots)^3} \subset H^*(P_2X).$$

Note that $\operatorname{Sq}^2 H^6(P_2X) = 0$. Using the Adem relation $\operatorname{Sq}^{2,3} = \operatorname{Sq}^5 + \operatorname{Sq}^{4,1}$ and the fact that $\operatorname{Sq}^3 u_i = 0$, we obtain

$$\operatorname{Sq}^4 v_i = \operatorname{Sq}^{4,1} u_i = \operatorname{Sq}^5 u_i = \operatorname{Sq}^{1,4} u_i.$$

Hence, if $\rho: H^*(P_2X; Z) \rightarrow H^*(P_2X; Z_2)$ is mod 2 reduction, we have

$$(2.3) \quad \operatorname{Sq}^4 H^6(P_2X; Z_2) \subset \operatorname{Im}(\rho).$$

Consider the following commutative diagram of infinite loop maps

$$\begin{array}{ccc}
 K(Z_2; 7, 9) & \longrightarrow & K(Z_2; 7, 9) \\
 \downarrow j & & \downarrow \bar{j} \\
 E & \xrightarrow{h} & \bar{E} \\
 \downarrow p & & \downarrow \bar{p} \\
 K(Z_2, 5) \times K(Z, 10) & \xrightarrow{\delta} & K(Z; 6, 10) \\
 \downarrow w & & \downarrow \bar{w} \\
 K(Z_2; 8, 10) & \longrightarrow & K(Z_2; 8, 10)
 \end{array}
 \quad (2.4)$$

in which

$$\begin{aligned}
 \bar{w}^*(\iota_8) &= \text{Sq}^2 \iota_6; & \bar{w}^*(\iota_{10}) &= \text{Sq}^4 \iota_6 - p(\iota_{10}); \\
 w^*(\iota_8) \text{Sq}^2 \text{Sq}^1 \iota_5; & & w^*(\iota_{10}) &= \text{Sq}^4 \text{Sq}^1 \iota_5 - p(\iota_{10});
 \end{aligned}$$

and

$$\delta^*(\iota_{10}) = \iota_{10}; \quad \delta^*(\iota_6) = \text{Sq}^1 \iota_5.$$

There exist elements $\bar{v}_{1,1} \in PH^9(\bar{E})$, $\bar{v}_{0,2} \in PH^{10}(\bar{E})$ such that $\bar{j}^*(\bar{v}_{1,1}) = \text{Sq}^2 \iota_7$ and $\bar{j}^*(\bar{v}_{0,2}) = \text{Sq}^{2,1} \iota_7 + \text{Sq}^1 \iota_9$.

Lemma 2.1. *If $v_{1,1} = h^*(\bar{v}_{1,1})$ and $v_{0,2} = h^*(\bar{v}_{0,2})$, then $\text{Sq}^2 v_{1,1} = \text{Sq}^1 v_{0,2}$.*

Proof. Note that

$$\text{Sq}^2 \bar{v}_{1,1} + \text{Sq}^1 \bar{v}_{0,2} \in \ker \bar{j}^* = \bar{p}^*(PH^{11}(K(Z; 6, 10))) = 0.$$

Therefore $\text{Sq}^2 \bar{v}_{1,1} = \text{Sq}^1 \bar{v}_{0,2}$. Applying h^* gives the lemma. $\square$

Define $\varphi_{1,1}$ and $\varphi_{0,2}$ to be the secondary operations defined by the elements $v_{1,1}$ and $v_{0,2}$, respectively. We note that if $y \in H^5(P_2 X)$, then $\varphi_{1,1}$ and $\varphi_{0,2}$ are defined on $\text{Sq}^1 y$ with zero indeterminacy.

We have the following relations:

- (1) $\varphi_{1,1} \text{Sq}^1 H^5(P_2 X) \subset \text{Sq}^4 H^5(P_2 X)$, since $H^9(P_2 X) = \text{Sq}^4(P_2 X)$.
- (2) $\text{Sq}^2 \varphi_{1,1} \text{Sq}^1 H^5(P_2 X) = \text{Sq}^1 \varphi_{0,2} \text{Sq}^1 H^5(P_2 X)$, by Lemma 2.1.

Proposition 2.2. $\varphi_{0,2} \text{Sq}^1 H^5(P_2 X) \subset \text{Sq}^5 H^5(P_2 X)$.

Proof. Let y be in $H^5(P_2X)$. Then

$$\begin{aligned} \text{Sq}^1 \varphi_{0,2} \text{Sq}^1 y &= \text{Sq}^2 \varphi_{1,1} \text{Sq}^1 y \quad \text{by Lemma 2.1} \\ &= \text{Sq}^2 \text{Sq}^4 y' \quad \text{by 1.} \\ &= \text{Sq}^5 \text{Sq}^1 y' \\ &= \text{Sq}^1 \text{Sq}^4 \text{Sq}^1 y' \\ &= \text{Sq}^1 \text{Sq}^1 \text{Sq}^4 y' \\ &= 0. \end{aligned}$$

But $H^{10}(P_2X) \cap \ker \text{Sq}^1 \subset \text{Sq}^5 H^5(P_2X)$. $\square$

In the proofs of the next two theorems we will be using properties of the c -invariant and the H -deviation. We refer the reader to the appendix of [5] for a summary of general properties of these operations.

Theorem 2.3. $\varphi_{1,1} \text{Sq}^1 H^5(P_2X) = 0$.

Proof. We now construct a Postnikov system which reflects the following relations for $u \in H^5(P_2X)$:

- (1) $\varphi_{1,1} \text{Sq}^1 u = \text{Sq}^4 u'$,
- (2) $\varphi_{0,2} \text{Sq}^1 u = \text{Sq}^5 u''$.

Consider the diagram

$$\begin{array}{ccc} & K(Z_2; 8, 9) & \\ & \downarrow & \\ & E_1 & \\ & \downarrow & \\ (2.5) & E & \xrightarrow{w} K(Z_2; 9, 10) \\ & \downarrow & \\ & K(Z_2; 5, 5, 5) \times K(Z; 10, 10) & \xrightarrow{w_0} K(Z_2; 8, 10, 10) \end{array}$$

in which

$$w_0^*(\iota_8) = \text{Sq}^{2,1} \iota_5, \quad w_0^*(\iota_{10}) = \text{Sq}^{4,1} \iota_5 - \iota_{10}, \quad w_0^*(\iota'_{10}) = \text{Sq}^{4,1} \iota'_5 - \iota'_{10},$$

and

$$w^*(\iota_9) = \varphi_{1,1} \text{Sq}^1 \iota_5 - \text{Sq}^4 \iota'_5, \quad w^*(\iota_{10}) = \varphi_{0,2} \text{Sq}^1 \iota_5 - \text{Sq}^5 \iota''_5.$$

We shall retain the notation ι_5 and ι'_5 to denote the images of these elements in E and E_1 .

In $H^*(E_1)$ we have

$$\begin{aligned}
 \text{Sq}^6 i'_5 &= (\text{Sq}^2 \text{Sq}^4 + \text{Sq}^5 \text{Sq}^1) i'_5 \\
 &= \text{Sq}^2 \text{Sq}^4 i'_5 + \text{Sq}^1 (\text{Sq}^4 \text{Sq}^1 i'_5) \\
 &= \text{Sq}^2 \varphi_{1,1} \text{Sq}^1 i'_5 + \text{Sq}^1 i'_{10} \\
 &= \text{Sq}^1 \varphi_{0,2} \text{Sq}^1 i_5 \\
 &= \text{Sq}^1 \text{Sq}^5 i''_5.
 \end{aligned}$$

This Postnikov system can be delooped. It follows that there is an element $v \in H^{10}(E_1)$ with $j_1^*(v) = \text{Sq}^2 i_8 + \text{Sq}^1 i'_9$ and $\bar{\Delta}v = i'_5 \otimes i'_5$. Therefore $c(\sigma^*v) = \sigma^*(i'_5) \otimes \sigma^*(i'_5)$. Since P_2X lifts to E_1 , there is an H -lifting f_1 of X to ΩE_1 . Checking degrees, one sees that X lifts to ΩE_1 by a c -map. This implies that

$$c(f_1^* \sigma^* v) = x' \otimes x' \neq 0, \quad \text{where } u' = i(x').$$

Therefore $f_1^* \sigma^* v$ is a nonzero primitive in $H^9(X)$. But $PH^9(X) = 0$. $\square$

3. PROOF OF THEOREM A

We can now prove Theorem A.

Proof. By (2.2), if $u \in H^5(P_2X)$, then $(\text{Sq}^4 u)^2 \neq 0$. We have

$$\begin{aligned}
 (\text{Sq}^4 u)^2 &= \text{Sq}^{9,4} u \\
 &= \text{Sq}^{8,4,1} u \quad \text{since } \text{Sq}^2 u = 0 \\
 &= \text{Sq}^4 \tilde{\varphi}_{0,3} \text{Sq}^1 u + (\text{Sq}^9 + \text{Sq}^{7,2}) \varphi_{1,1} \text{Sq}^1 u \quad \text{by Theorem 1.1} \\
 &= \text{Sq}^4 \tilde{\varphi}_{0,3} \text{Sq}^1 u \quad \text{by Theorem 2.3.}
 \end{aligned}$$

Consider the following Postnikov system:

$$\begin{array}{ccc}
 & & K(Z_2; 8, 13) \\
 & \downarrow j_1 & \\
 & E_1 & \\
 & \downarrow p_1 & \\
 & E & \xrightarrow{w_1} K(Z_2; 9, 14) \\
 & \downarrow r & \\
 K(Z_2, 5) & \xrightarrow{w_0} & K(Z_2; 7, 8)
 \end{array}$$

in which

$$\begin{aligned}
 w_0^*(i_7) &= \text{Sq}^2 i_5; & w_0^*(i_8) &= \text{Sq}^{2,1} i_5; \\
 w_1^*(i_9) &= \varphi_{1,1} \text{Sq}^1 i_5; & w_1^*(i_{14}) &= \tilde{\varphi}_{0,3} \text{Sq}^1 i_5.
 \end{aligned}$$

In $H^*(E_1)$,

$$\begin{aligned} (\text{Sq}^4 \iota_5)^2 &= \text{Sq}^{8,5} \iota_5 \\ &= \text{Sq}^{8,4,1} \iota_5 \\ &= \text{Sq}^4 \tilde{\phi}_{0,3} \text{Sq}^1 \iota_5 + (\text{Sq}^9 + \text{Sq}^{7,2}) \phi_{1,1} \text{Sq}^1 \iota_5 \\ &= 0. \end{aligned}$$

Therefore in $H^{16}(\Omega E_1)$ there exists an element v with

$$\begin{aligned} (\Omega j_1)^*(v) &= \text{Sq}^4 \iota_{12} + (\text{Sq}^9 + \text{Sq}^{7,2}) \iota_7, \\ \bar{\Delta} v &= \iota_4^2 \otimes \iota_4^2. \end{aligned}$$

We have the following commutative diagram

$$\begin{array}{ccc} & & K(Z_2; 7, 12) \\ & & \downarrow \\ & \Omega E_1 & \\ & \nearrow f_1 & \\ & \Omega E & \xrightarrow{\Omega w_1} K(Z_2; 8, 13) \\ & \nwarrow f_0 & \\ & \downarrow r & \\ X \xrightarrow{f} K(Z_2, 4) & \xrightarrow{\Omega w_0} & K(Z_2; 6, 7) \end{array}$$

Here f is an H -map, since the relations defined by w_0 hold in $P_2 X$. Further, $c(f)$ factors through $K(Z_2; 4, 5)$, so in fact f is a c -map. It follows that $(1 - T^*)[D_{f_1}] = \Omega w_1 \circ c(f) \simeq *$. So $[D_{f_1}] \in H^{12}(X \wedge X) \cap \ker(1 - T^*)$.

Any terms of the form $x_i \otimes x_j^2 + x_j^2 \otimes x_i$ may be eliminated by changing f_1 by $x_i x_j^2$. It follows that

$$[D_{f_1}] = \sum_{j \neq k} b_{ijk} (x_i \otimes x_j x_k + x_j x_k \otimes x_i), \quad b_{ijk} \in Z_2.$$

Therefore

$$\bar{\Delta} f_1^*(v) = x^2 \otimes x^2 + \text{Sq}^4 \sum b_{ijk} (x_i \otimes x_j x_k + x_j x_k \otimes x_i) \neq 0.$$

This implies that if $\langle t, x^2 \rangle \neq 0$, then $t^2 \neq 0$ for $t \in H_*(X)$. But all squares are zero in $H_*(X)$, since $H_*(X)$ is primitively generated. We conclude that X is not homotopy-commutative. $\square$

REFERENCES

1. W. Browder and E. Thomas, *On the projective plane of an H -space*, Ill. J. Math. **7** (1963), 492–502.
2. J. Harper, *On the cohomology of stable two-stage Postnikov systems*, Trans. Amer. Math. Soc. **152** (1970), 375–388.

3. J. Hubbuck, *On homotopy commutative H -spaces*, *Topology* **8** (1969), 119–126.
4. J. Lin, *A cohomological proof of the torus theorem*, *Math. Z.* **190** (1985), 469–476.
5. J. Lin and F. Williams, *On 6-connected finite H -spaces with two torsion*, *Topology* **28** (1989), 7–34.
6. M. Slack, *Maps between iterated loop spaces*, *J. Pure Appl. Algebra* (to appear).
7. —, *A classification of homotopy commutative finitely generated H -spaces*, *Mem. Amer. Math. Soc.* (to appear).
8. H. Toda, *Composition methods in homotopy groups of spheres*, *Ann. of Math. Studies* No. 49, Princeton Univ. Press, Princeton, NJ, 1961.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA AT SAN DIEGO, LA JOLLA, CALIFORNIA 92093

DEPARTMENT OF MATHEMATICS, NEW MEXICO STATE UNIVERSITY, LAS CRUCES, NEW MEXICO 88001