DIVISIBILITY CONSTRAINTS ON DEGREES OF FACTOR MAPS

PAUL TROW

(Communicated by Kenneth R. Meyer)

ABSTRACT. We show that the degree of a finite-to-one factor map $f: \sum_A \rightarrow \sum_B$ between shifts of finite type is constrained by the factors of χ_A and χ_B . A special case of these constraints is that if $2 \operatorname{rank}^* B > \operatorname{rank}^* A$, then the degree of f is a unit in $\mathbb{Z}[1/\det^* B]$ (where $\operatorname{rank}^* A$ is the rank of the Jordan form away from 0 of A , and $\det^* B$ is the determinant of the Jordan form away from 0 of B).

0. INTRODUCTION

The degree of a finite-to-one factor map $f: \sum_A \rightarrow \sum_B$ between irreducible shifts of finite type is the largest positive integer d such that every $y \in \sum_B$ has d mutually separated inverse images. This concept was introduced in [H] for endomorphisms of full shifts. In [B], M. Boyle showed that, given $\sum_A$ and $\sum_B$, there exists a finite set E of positive integers such that if $f: \sum_A \rightarrow \sum_B$ has degree d , then $d = eu$, where $e \in E$ and u is a unit in $\mathbb{Z}[1/\lambda]$, where λ is the dominant eigenvalue of A . In [T2], this result was extended to apply to any nonzero eigenvalue γ that has the same algebraic multiplicity in χ_A and χ_B . In this paper we give further constraints on the possible degrees of a finite-to-one factor map, given certain assumptions on the factors of χ_A and χ_B (see Theorem 8 and Corollary 9).

1. BACKGROUND

We give some basic definitions concerning shifts of finite type. For a more thorough introduction to symbolic dynamics, see [AM] or [PT].

Let A be an $n \times n$, nonnegative, integral matrix, and let $G(A)$ be the directed graph on n vertices, with A_{ij} edges from vertex i to vertex j . Let $\mathcal{E}_A = \{\text{edges of } G(A)\}$. We define the *shift of finite type* $\sum_A = \{x \in \mathcal{E}_A^{\mathbb{Z}} : x_{i+1} \text{ follows } x_i \text{ for all } i\}$, where x_{i+1} follows x_i if the terminal vertex of x_i is the initial vertex of x_{i+1} . $\sum_A$ is a compact metric space under the product topology.

A is *irreducible* if for each i, j , there exists $n = n(i, j)$ such that $A_{ij}^n > 0$. $\sum_A$ is irreducible if A is. A *factor map* $f: \sum_A \rightarrow \sum_B$ is a continuous,

Received by the editors April 28, 1990.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 28D05.

©1991 American Mathematical Society
 0002-9939/91 \$1.00 + \$.25 per page

surjective map which commutes with the shift. f is *finite-to-one* if there exists $N \in \mathbb{Z}^+$ such that $\#f^{-1}(y) \leq N$, for all $y \in \sum_B$. By [CP, Corollary 4.5], this is equivalent to $h(\sum_A) = h(\sum_B)$.

If D is a principal ideal domain, and $g, h \in D$, we denote the greatest common divisor of g and h by (g, h) . If $p(x) \in \mathbb{Z}[x]$, we denote the degree of $p(x)$ by $\deg p(x)$.

If A is a square, integral matrix, and $\chi_A = p(x)x^r$, where r is the largest power of x which divides χ_A , let $\chi_A^* = p(x)$, $\text{rank}^* A = \deg p(x)$, and $\det^* A = \text{constant term of } p(x)$.

If F is a field, A is a square matrix over F , and $g(x)$ is a factor of χ_A , we denote the null space of $g(A)$ by $N(g(A))$. If V is a vector space over F , we denote the dual space of V by V^* . If W is a subspace of V , the annihilator of W , denoted $\text{Ann } W$, is $\{f \in V^*: f(w) = 0 \text{ for all } w \in W\}$.

If $\mathcal{L} \subseteq \mathbb{Z}^n$, $\mathcal{L}$ is called a *sublattice*. $\mathcal{L}$ is *closed* if $\mathcal{L} = \langle \mathcal{L} \rangle \cap \mathbb{Z}^n$, where $\langle \mathcal{L} \rangle$ denotes the rational span of $\mathcal{L}$.

2. MAIN RESULTS

Definition 1. Let $p(x)$ be a monic, integral polynomial, and $p(x) = \prod_{i=1}^t p_i^{e_i}(x)$ be the unique factorization of $p(x)$ in $\mathbb{Z}[x]$, where the $p_i(x)$ are distinct, monic, irreducible polynomials. Any factor of $p(x)$ which is a product of some of the $p_i^{e_i}(x)$ is called a *complete factor* of $p(x)$. If $g(x)$ is a complete factor of $p(x)$, and $h(x)$ is a complete factor of $p(x)/g(x)$, then $h(x)$ is called a *complementary complete factor* of $p(x)$.

If $f: \sum_A \rightarrow \sum_B$ is a finite-to-one factor map, then χ_B^* divides χ_A^* (see [K, Corollary A]). If $g_B(x) = \prod p_i^{e_i}(x)$ is a complete factor of $p(x) = \chi_B^*$, then $g_A(x) = \prod p_i^{f_i}(x)$, $f_i \geq e_i$, is a complete factor of χ_A^* , where f_i is the power to which p_i is raised in χ_A^* . Then $g_A(x)$ is called the *corresponding complete factor* of χ_A^* .

Lemma 2. Let B be a $k \times k$ integral matrix, and let $g(x)$ be a complete factor of χ_B . Then there is a closed, B -invariant sublattice $\mathcal{L} \subseteq N(g(B))$ such that $\text{rank } \mathcal{L} = \deg g(x)$.

Proof. Let $\mathcal{L} = N(g(B)) \cap \mathbb{Z}^k$. By the Primary Decomposition Theorem ([HK, Theorem 12, p. 189]), $\text{rank } \mathcal{L} = \deg g(x)$. $\square$

We now consider a sublattice $\mathcal{L}$ reduced over $\mathbb{Z}/p\mathbb{Z}$, for p a prime. Lemmas 4–6 are elementary results from linear algebra.

Definition 3. For p a fixed prime, let $\pi: \mathbb{Z} \rightarrow \mathbb{Z}/p\mathbb{Z}$ denote the natural homomorphism $k \rightarrow k + p\mathbb{Z}$. Denote $\pi(k)$ by $\bar{k}$. If $A = (a_{ij})$ is an $n \times n$ integral matrix, let $\bar{A}$ denote the $n \times n$ matrix over $\mathbb{Z}/p\mathbb{Z}$ given by $\bar{A}_{ij} = \bar{a}_{ij}$. If $v = (v_1, \dots, v_n) \in \mathbb{Z}^n$, let $\bar{v} \in (\mathbb{Z}/p\mathbb{Z})^n$ be defined by $\bar{v} = (\bar{v}_1, \dots, \bar{v}_n)$. If $\mathcal{L} \subseteq \mathbb{Z}^n$, let $\bar{\mathcal{L}} = \{\bar{v}: v \in \mathcal{L}\}$.

Lemma 4. *If A is an $n \times n$ integral matrix and $\mathcal{L} \subseteq \mathbb{Z}^n$ is a closed sublattice, then $\overline{\mathcal{L}}$ is an $\overline{A}$ -invariant subspace of $(\mathbb{Z}/p\mathbb{Z})^n$ and $\text{rank } \mathcal{L} = \dim \overline{\mathcal{L}}$.*

Proof. It is easy to check that $\mathcal{L}$ is an $\overline{A}$ -invariant subspace. To see that $\text{rank } \mathcal{L} = \dim \overline{\mathcal{L}}$, choose an integral basis $\mathcal{B} = \{\beta_1, \dots, \beta_k\}$ for $\mathcal{L}$. We claim that $\overline{\mathcal{B}} = \{\overline{\beta}_1, \dots, \overline{\beta}_k\}$ is a basis for $\overline{\mathcal{L}}$. If $\overline{v} \in \overline{\mathcal{L}}$, where $v \in \mathcal{L}$, then $v \in \sum_{i=1}^k c_i \beta_i$, for some $c_i \in \mathbb{Z}$, since $\mathcal{B}$ is an integral basis for $\mathcal{L}$. Therefore $\overline{v} = \sum_{i=1}^k \overline{c_i} \overline{\beta_i}$, so $\overline{\mathcal{B}}$ spans $\overline{\mathcal{L}}$. If $\sum_{i=1}^k \overline{c_i} \overline{\beta_i} = (\overline{0}, \dots, \overline{0})$, then p divides $(\sum_{i=1}^k c_i \beta_i)_j$, for $1 \leq j \leq n$. It follows that $v = \sum_{i=1}^k (c_i/p) \beta_i \in \mathcal{L}$, since $v \in \langle \mathcal{L} \rangle \cap \mathbb{Z}^n$ and $\mathcal{L}$ is closed. This implies that $c_i/p \in \mathbb{Z}$, since $\mathcal{B}$ is an integral basis for $\mathcal{L}$. Therefore $\overline{c_i} = \overline{0}$, so $\overline{\mathcal{B}}$ is linearly independent. It follows that $\text{rank } \mathcal{L} = \dim \overline{\mathcal{L}}$. $\square$

Lemma 5. *Let $\overline{A}$ be an $n \times n$ matrix over a field F . Let $\overline{g}(x)$ and $\overline{h}(x)$ be factors of $\chi_{\overline{A}}$. If $(\overline{g}(x), \overline{h}(x)) = 1$ in $F[x]$, and $v \in N(\overline{h}(\overline{A}))$, $w \in N(\overline{g}(\overline{A}^T))$, then $vw^T = 0$.*

Proof. Because $(\overline{g}(x), \overline{h}(x)) = 1$, there exist $q(x), r(x) \in F[x]$ such that $\overline{g}(x)q(x) + \overline{h}(x)r(x) = 1$. Therefore $v = v\overline{g}(A)q(A)$ and $w^T = \overline{h}(A)r(A)w^T$. So $vw^T = v\overline{g}(A)q(A)\overline{h}(A)r(A)w^T = v\overline{h}(A)q(A)r(A)\overline{g}(A)w^T = 0$. $\square$

Lemma 6. *Let U and V be subspaces of $(\mathbb{Z}/p\mathbb{Z})^n$. Assume that $\overline{v}\overline{w}^T = \overline{0}$ for all $\overline{v} \in U$, $\overline{w} \in V$. Then $\dim U \leq \dim \text{Ann}(V)$.*

Proof. It is easy to check that the linear map $\phi: (\mathbb{Z}/p\mathbb{Z})^n \rightarrow (\mathbb{Z}/p\mathbb{Z})^{n^*}$ given by $v \mapsto f_v$, where $f_v(w) = vw^T$, is injective. Since $\phi(U) \subseteq \text{Ann}(V)$, the result follows. $\square$

Theorem 7. *Let A be an $n \times n$ integral matrix, p a prime, $g(x)$ a complete factor of χ_A^* , with constant term c , and $h(x)$ a complementary complete factor of χ_A^* . Assume that $(\overline{g}(x), \overline{h}(x)) = \overline{1}$ and that $(p, c) = 1$. Let $\mathcal{L}_1$ be a closed, A -invariant sublattice of $N(g(A))$ and $\mathcal{L}_2$ a closed, A^T -invariant sublattice of $N(g(A^T))$. If p divides vw^T , for all $v \in \mathcal{L}_1$, $w \in \mathcal{L}_2$, then $\text{rank } \mathcal{L}_1 + \text{rank } \mathcal{L}_2 \leq \text{rank}^* A - \deg h(x)$.*

Proof. Let $\deg h(x) = s$, and let r be the largest power of x which divides χ_A . Clearly, $\text{rank}^* A = n - r$. Since $(p, c) = 1$, it is easy to see that $(\overline{g}(x), \overline{x}) = \overline{1}$. Let $h_1(x) = h(x)x^r$. Then $(\overline{g}(x), \overline{h}_1(x)) = \overline{1}$. We have $\dim N(h_1(A)) \leq \dim N(\overline{h}_1(\overline{A}))$, since $N(h_1(A)) \cap \mathbb{Z}^n$ projects onto a subspace of $N(\overline{h}_1(\overline{A}))$ whose dimension equals $\text{rank } N(h_1(A)) \cap \mathbb{Z}^n$, by Lemma 4. Also, $\dim N(h_1(A)) = \deg h_1(x) = s + r$ by the Primary Decomposition Theorem ([HK, Theorem 12]), since $h_1(x)$ is a complete factor of χ_A . So $\dim N(\overline{h}_1(\overline{A})) \geq s + r$. Also, we have $\overline{v}\overline{w}^T = \overline{0}$, for all $\overline{v} \in \overline{\mathcal{L}}_1$, $\overline{w} \in \overline{\mathcal{L}}_2$, since p divides vw^T . Let U be the span over $\mathbb{Z}/p\mathbb{Z}$ of $N(\overline{h}_1(\overline{A}))$ and $\overline{\mathcal{L}}_1$. Since $\overline{\mathcal{L}}_1 \subseteq N(\overline{g}(\overline{A}))$ and $(\overline{g}(x), \overline{h}_1(x)) = \overline{1}$, it is easy to see that $N(\overline{h}_1(\overline{A})) \cap \overline{\mathcal{L}}_1 = \{0\}$, so that $\dim U = \dim N(\overline{h}_1(\overline{A})) + \dim \overline{\mathcal{L}}_1 \geq s + r + \dim \overline{\mathcal{L}}_1$. Also, since

$\overline{\mathcal{L}}_2 \subseteq N(\overline{g}(\overline{A}^T))$, it follows from Lemma 5 that $\overline{v}\overline{w}^T = \overline{0}$, for all $\overline{v} \in U$, $\overline{w} \in \overline{\mathcal{L}}_2$. By Lemma 6, $s + r + \dim \overline{\mathcal{L}}_1 \leq \dim U \leq \dim \text{Ann}(\overline{\mathcal{L}}_2)$. Since $\dim \overline{\mathcal{L}}_2 + \dim \text{Ann}(\overline{\mathcal{L}}_2) = \dim(\mathbb{Z}/p\mathbb{Z})^n = n$ (by [HK, Theorem 17, p. 98]), we have $s + r + \dim \overline{\mathcal{L}}_1 + \dim \overline{\mathcal{L}}_2 \leq n$ or $\dim \overline{\mathcal{L}}_1 + \dim \overline{\mathcal{L}}_2 \leq n - r - s = \text{rank}^* A - \deg h(x)$. Since $\dim \overline{\mathcal{L}}_1 = \text{rank} \mathcal{L}_1$ and $\dim \overline{\mathcal{L}}_2 = \text{rank} \mathcal{L}_2$, by Lemma 4, the result follows. $\square$

Theorem 8. Suppose that $\sum_A$ and $\sum_B$ are irreducible shifts of finite type, and that $f: \sum_A \rightarrow \sum_B$ is a finite-to-one factor map of degree d . Suppose that $g_B(x)$ is a complete factor of χ_B^* and $g_A(x)$ is the corresponding complete factor of χ_A^* , where the constant term of $g_A(x)$ is c , and that $h(x)$ is a complementary complete factor of χ_A^* . Let p be a prime which divides d , and assume that $(\overline{g}_A(x), \overline{h}(x)) = \overline{1}$ and that $2 \deg g_B(x) > \text{rank}^* A - \deg h(x)$. Then p divides c .

Proof. By [KMT, Theorem 2.3] and [T2, Definition 1.4 and Lemma 1.6] (substituting $\mathbb{Q}$ for $\mathbb{C}$), there exist $\sum_{\tilde{A}}$ topologically conjugate to $\sum_A$, and $\sum_{\tilde{B}}$ conjugate to $\sum_B$, a rational $\tilde{A}$ -invariant subspace V , a rational $\tilde{A}^T$ -invariant subspace W , and surjective linear maps $\phi: V \rightarrow \mathbb{Q}^k$ and $\theta: W \rightarrow \mathbb{Q}^k$ (where $\tilde{B}$ is $k \times k$) such that $v\tilde{A}\phi = v\phi\tilde{B}$ and $w\tilde{A}^T\theta = w\theta\tilde{B}^T$, for all $v \in V$, $w \in W$. By [W, Corollary 4.8], $\chi_A^* = \chi_{\tilde{A}}^*$ and $\chi_B^* = \chi_{\tilde{B}}^*$, so $g_B(x)$ is a complete factor of $\chi_{\tilde{B}}^*$, $g_A(x)$ is the corresponding complete factor of $\chi_{\tilde{A}}^*$, and $h(x)$ is a complementary complete factor of $\chi_{\tilde{A}}^*$. It follows from Lemma 2 and duality that there exist a closed $\tilde{A}$ -invariant sublattice $\tilde{\mathcal{L}}_1 \subseteq N(g_A(\tilde{A})) \cap V$ and a closed $\tilde{A}^T$ -invariant sublattice $\tilde{\mathcal{L}}_2 \subseteq N(g_A(\tilde{A}^T)) \cap W$ such that $\text{rank} \tilde{\mathcal{L}}_1 = \text{rank} \tilde{\mathcal{L}}_2 = \deg g_B(x)$. So $\text{rank} \tilde{\mathcal{L}}_1 + \text{rank} \tilde{\mathcal{L}}_2 = 2 \deg g_B(x) > \text{rank}^* A - \deg h(x)$. In [T2, Theorem 2.5], it is shown that if $v \in V$ and $w \in W$ are $\mathbb{Z}$ -vectors, then d divides vw^T , and so p divides vw^T . If $(p, c) = 1$, then by Theorem 7 there would exist $v \in \tilde{\mathcal{L}}_1$ and $w \in \tilde{\mathcal{L}}_2$ such that p does not divide vw^T , contradicting the previous statement. Therefore p divides c . $\square$

Corollary 9. Suppose that $\sum_A$ and $\sum_B$ are irreducible shifts of finite type, and $f: \sum_A \rightarrow \sum_B$ is a finite-to-one factor map of degree d . If $2 \text{rank}^* B > \text{rank}^* A$, then every prime which divides d also divides $\det^* B$. Therefore d is a unit in $\mathbb{Z}[1/\det^* B]$.

Proof. Take $g_B(x) = \chi_B^*$, $g_A(x) = \chi_A^*$, and $h(x) = 1$. For any prime p , the condition $(\overline{g}_B(x), \overline{h}(x)) = \overline{1}$ holds trivially. Since $\deg g_B(x) = \text{rank}^* B$, $\deg g_A(x) = \text{rank}^* B$, and the constant term of χ_B^* is $\det^* B$, the result follows from Theorem 8. $\square$

Example 10. Suppose that $\sum_A$ and $\sum_B$ are irreducible shifts of finite type, and that $\chi_A^* = (x-2)(x-1)^3$ and $\chi_B^* = (x-2)(x-1)^2$. Take $g_B(x) = (x-1)^2$, $g_A(x) = (x-1)^3$, and $h(x) = x-2$. Observe that for any prime p , $(\overline{g}_A(x), \overline{h}(x)) = \overline{1}$. Since $2 \deg g_B(x) = 4 > \text{rank}^* A - \deg h(x) = 4 - 1 = 3$,

and the constant term of $g_A(x)$ is 1, if $f: \sum_A \rightarrow \sum_B$ is a factor map, then the degree of f is 1, by Theorem 8.

We remark that if we assume only that $\chi_A^* = (x-2)(x-1)^2$ and $\chi_B^* = (x-2)(x-1)$, we cannot draw the same conclusion. However, by applying Corollary 9, we can conclude that the degree of f is a power of 2, but we do not have an example of a factor map between shifts with the given characteristic polynomials (modulo x) which has degree 2. Does such a factor map exist?

Example 11. This example shows that Corollary 9 is sharp, in general. Let

$$A = \begin{bmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}.$$

Then there exists a factor map $f: \sum_A \rightarrow \sum_B$, of degree 2, which is right and left resolving (and therefore constant-to-one). Then f is defined as a map on states, sending states 1 and 2 of $\sum_A$ to state 1 of $\sum_B$ and states 3 and 4 of $\sum_A$ to state 2 of $\sum_B$. Note that $\det^* B = -1$, so 2 does not divide $\det^* B$, but that $2 \operatorname{rank}^* B = 4 = \operatorname{rank}^* A$.

Example 12. This example shows that the condition that $(\bar{g}(x), \bar{h}(x)) = \bar{1}$ in Theorem 8 is necessary. Let

$$A = \begin{bmatrix} 1 & 0 & 0 & 2 \\ 0 & 1 & 2 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{bmatrix} \quad \text{and} \quad B = \begin{bmatrix} 1 & 2 \\ 1 & 0 \end{bmatrix}.$$

There exists a factor map $f: \sum_A \rightarrow \sum_B$ of degree 2, similar to the one in the previous example, and f maps states 1 and 2 of $\sum_A$ to state 1 of $\sum_B$ and states 3 and 4 of $\sum_A$ to state 2 of $\sum_B$. Let $p = 2$. Observe that $\chi_B = (x-2)(x+1)$ and $\chi_A = (x-2)(x+1)(x^2-x+2)$. Let $g_B(x) = x+1 = g_A(x)$ and $h(x) = (x-2)(x^2-x+2)$. Note that $2 \deg g_B(x) = 2 > \operatorname{rank}^* A - \deg h(x) = 4 - 3 = 1$, but that $(\bar{g}_A(x), \bar{h}(x)) \neq \bar{1}$, since $\bar{g}_A(x) = x+1$ and $\bar{h}(x) = x^2(x+1)$. Also, the conclusion of Theorem 8 does not hold, since the degree of f is not 1.

REFERENCES

- [AM] R. L. Adler and B. Marcus, *Topological entropy and equivalence of dynamical systems*, Mem. Amer. Math. Soc., no. 219, Amer. Math. Soc., Providence, RI, 1979.
- [B] M. Boyle, *Constraints on the degree of a sofic homomorphism and the induced multiplication of measures on unstable sets*, Israel J. Math. **53** (1986), 52–68.
- [CP] E. Coven and M. Paul, *Sofic systems*, Israel J. Math. **20** (1975), 165–177.
- [H] G. A. Hedlund, *Endomorphisms and automorphisms of the shift dynamical system*, Math. Systems Theory **3** (1969), 320–375.
- [HK] K. Hoffman and R. Kunze, *Linear algebra*, Prentice-Hall, Englewood Cliffs, NJ, 1961.
- [K] B. Kitchens, *An invariant for continuous factors of Markov chains*, Proc. Amer. Math. Soc. **83** (1981), 825–828.

- [KMT] B. Kitchens, B. Marcus, and P. Trow, *Eventual factor maps and compositions of closing maps*, Ergodic Theory Dynamical Systems (to appear).
- [PT] W. Parry and S. Tuncel, *Classification problems in ergodic theory*, London Math. Soc. Lecture Notes Ser., vol. 67, Cambridge University Press, Cambridge and New York, 1982.
- [T1] P. Trow, *Degrees of constant-to-one factor maps*, Proc. Amer. Math. Soc. **103** (1988), 184–188.
- [T2] ———, *Degrees of finite-to-one factor maps*, Israel J. Math. (to appear).
- [W] R. F. Williams, *Classification of one-dimensional attractors*, Proc. Sympos. Pure Math., vol. 14, Amer. Math. Soc., Providence, RI, 1970, 341–361.

DEPARTMENT OF MATHEMATICAL SCIENCES, MEMPHIS STATE UNIVERSITY, MEMPHIS, TENNESSEE 38152