PRODUCT SHIFTS ON $B(H)$

P. J. STACEY

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. A shift on $B(H)$ is a $*$ -endomorphism α for which $\bigcap_r \alpha^r(B(H)) = \mathbb{C}P$ for some projection P . The paper discusses some aspects of the classification of shifts on $B(H)$ up to conjugacy by $*$ -automorphisms, with a focus on the shifts arising from an infinite tensor product decomposition of H .

1. INTRODUCTION

Let H be a separable Hilbert space and let $B(H)$ be the algebra of all bounded linear operators on H . Slightly extending the definition given in [7] to include the nonunital case, a $*$ -endomorphism α of $B(H)$ will be called a shift if $\bigcap_r \alpha^r(B(H)) = \mathbb{C}P$ for some (possibly zero) projection P on $B(H)$. The present paper discusses some aspects of the classification of shifts on $B(H)$ up to conjugacy by $*$ -automorphisms, with a focus on the shifts arising from an infinite tensor product decomposition of H .

2. SPATIAL DESCRIPTIONS

It was observed in [2, Proposition 2.1] that every nonzero normal $*$ -endomorphism α of $B(H)$ is implemented by a sequence of isometries. An alternative proof of this result can be obtained by imitating the usual proof that $*$ -automorphisms of $B(H)$ are inner, as described, for example, in [5, Lemma 7.5.3], and this alternative proof shows that the normality assumption can be omitted (and hence every $*$ -endomorphism of $B(H)$ is normal). For later reference we now state this minor extension of [2, Proposition 2.1].

Proposition 2.1. *Let α be a nonzero $*$ -endomorphism of $B(H)$. Then there is a (finite or infinite) sequence of isometries $V_1, V_2, \dots$ in $B(H)$ having mutually orthogonal ranges such that*

$$\alpha(A) = \sum_n V_n A V_n^*$$

Received by the editors May 7, 1990.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 46L40; Secondary 46L55, 47A65.

Key words and phrases. Endomorphism, isometry, Hilbert space, shift, infinite tensor product, unitary equivalence.

for each $A \in B(H)$. The linear space of operators

$$E_\alpha = \{T \in B(H) : \alpha(A)T = TA \text{ for all } A \in B(H)\}$$

is a Hilbert space relative to the inner product defined by

$$T^*S = \langle S, T \rangle 1$$

and $\{V_1, V_2, \dots\}$ is an orthonormal basis for E_α .

The following simple result relates the conjugacy of α and β to the Hilbert spaces E_α and E_β .

Proposition 2.2. *Let α, β be *-endomorphisms of $B(H)$ and let $U \in B(H)$ be unitary. Then $\alpha = (\text{Ad } U)\beta(\text{Ad } U^*)$ if and only if $E_\alpha = UE_\beta U^*$.*

Proof. If $\alpha = (\text{Ad } U)\beta(\text{Ad } U^*)$ then an easy calculation yields $E_\alpha = UE_\beta U^*$. This same result shows that if $E_\alpha = UE_\beta U^*$ then $E_\alpha = E_\gamma$ where $\gamma = (\text{Ad } U)\beta(\text{Ad } U^*)$: hence there exist orthonormal bases $\{V_1, V_2, \dots\}$ and $\{W_1, W_2, \dots\}$ of E_α with $\alpha(A) = \sum V_n A V_n^*$ and $\gamma(A) = \sum W_n A W_n^*$ for each $A \in B(H)$. A simple calculation using the unitary transition matrix between the two orthonormal bases then yields the conclusion $\alpha = \gamma$. $\square$

A natural question arising from Propositions 2.1 and 2.2 is the extent to which the specification of the isometries V_i up to unitary equivalence determines the corresponding *-endomorphism up to conjugacy. The classification of an isometry V up to unitary equivalence is given by the *Wold decomposition*, described, for example, in [4, Problem 118], which decomposes V as a direct sum of copies of the unilateral shift and a unitary mapping on a subspace; when the unitary summand is absent the isometry is said to be *pure*. It will now be shown that if α is a shift, then at most one of any corresponding family of isometries is not pure; this result extends [1, Theorem 3] which is the case of two unitarily equivalent isometries. The proof is based on the following lemma in which Cuntz's notation $V_\mu V_\nu^*$ is used to denote $V_{\mu(1)} \cdots V_{\mu(r)} V_{\nu(s)}^* \cdots V_{\nu(1)}^*$, where $\mu = (\mu(1), \dots, \mu(r))$ has $r = |\mu|$ components and $\nu = (\nu(1), \dots, \nu(s))$ has $s = |\nu|$ components.

Lemma 2.3. *Let $V_1, V_2, \dots$ be isometries from a Hilbert space H onto a family of mutually orthogonal subspaces; α be the *-endomorphism defined by $\alpha(A) = \sum_n V_n A V_n^*$; and $h \in \bigcap_r V_i^r H$.*

- (i) *If P_h is the orthogonal projection from H onto the closed linear span S of the vectors $V_\mu V_\nu^* h$ with $|\mu| \geq 0$ and $|\nu| \geq 0$, then $\alpha(P_h) = P_h$ and hence $P_h \in \bigcap_r \alpha^r(B(H))$.*
- (ii) *If Q_h is the orthogonal projection from H onto the closed linear span T of the vectors $V_\mu V_\nu^* h$ with $|\mu| = |\nu| \geq 0$, then $Q_h \in \bigcap_r \alpha^r(B(H))$.*

Proof. (i) Let $|\mu| = r$, $|\nu| = s$, and $h = V_i^{s+1} k$, where $k \in H$. Then $\alpha(1)V_\mu V_\nu^* h = \sum V_n V_n^* V_\mu V_\nu^* V_i^{s+1} k = V_\mu V_\nu^* h$ (even if $r = 0$) and hence $\alpha(1) \geq P_h$. Since S is invariant under both V_n and V_n^* , it follows that $P_h V_n = V_n P_h$

for each n and hence $\alpha(P_h) = \sum V_n P_h V_n^* = P_h \sum V_n V_n^* = P_h \alpha(1) = P_h$.

(ii) Since T is invariant under each $V_\gamma V_\delta^*$ with $|\gamma| = |\delta|$, it follows that $Q_h V_\gamma V_\delta^* = V_\gamma V_\delta^* Q_h$. Thus

$$\begin{aligned} \alpha^r(V_1^{*r} Q_h V_1^r) &= \sum_{|\gamma|=r} V_\gamma V_1^{*r} Q_h V_1^r V_\gamma^* \\ &= \sum_{|\gamma|=r} V_\gamma V_1^{*r} V_1^r V_\gamma^* Q_h = \sum_{|\gamma|=r} V_\gamma V_\gamma^* Q_h \\ &= \alpha^r(1) Q_h. \end{aligned}$$

However, using part (i), $Q_h \leq P_h = \alpha^r(P_h) \leq \alpha^r(1)$ so $Q_h = \alpha^r(V_1^{*r} Q_h V_1^r)$ and hence $Q_h \in \bigcap_r \alpha^r(B(H))$. $\square$

Proposition 2.4. *Let $V_1, V_2, \dots$ be a sequence of isometries from a Hilbert space H onto a family of orthogonal subspaces and let α be the *-endomorphism of $B(H)$ defined by $\alpha(A) = \sum_n V_n A V_n^*$. If α is a shift, then at most one of the isometries V_n is not pure and if V_n is not pure then the subspace $\bigcap_r V_n^r H$ associated with the unitary summand of V_n is one-dimensional.*

Proof. Suppose, to obtain a contradiction, that there exist nonzero vectors $h \in \bigcap_r V_i^r H$ and $k \in \bigcap_r V_j^r H$ with $\langle h, k \rangle = 0$. Then, by part (ii) of the lemma, there exist associated nonzero projections Q_h, Q_k in $\bigcap_r \alpha^r(B(H))$. Let $|\mu| = |\nu|$ and $|\delta| = |\gamma|$. The proof will be completed if it is shown that $\langle V_\mu V_\nu^* h, V_\gamma V_\delta^* k \rangle = 0$ for then $Q_h Q_k = 0$ and, therefore, α is not a shift. To this end let $V_\delta V_\gamma^* V_\mu V_\nu^* = V_\rho V_\sigma^*$; $|\rho| = |\sigma| = r$; $h = V_i^{r+1} h'$; and $k = V_j^{r+1} k'$. Then $\langle V_\mu V_\nu^* h, V_\gamma V_\delta^* k \rangle = \langle V_\rho V_\sigma^* h, k \rangle = \langle V_\sigma^* h, V_\rho^* k \rangle = \delta_{\sigma(i, \dots, i) \rho(j, \dots, j)} \langle V_i h', V_j k' \rangle$. But, when $i \neq j$, $\langle V_i h', V_j k' \rangle = \langle h', V_i^* V_j k' \rangle = 0$ and, when $i = j$, $\langle V_i h', V_j k' \rangle = \langle V_i^{*r} h, V_i k' \rangle = \langle h, V_i^{r+1} k' \rangle = \langle h, k \rangle = 0$, as required. $\square$

Using Proposition 2.4, it is possible to give a complete classification up to conjugacy of the simplest shifts α , for which E_α is one-dimensional.

Proposition 2.5. (i) *Let V be an isometry on a separable infinite-dimensional Hilbert space H and let α be the *-endomorphism of $B(H)$ defined by $\alpha(A) = V A V^*$. Then α is a shift if and only if V is either a pure isometry or $\bigcap_r V^r H$ is one-dimensional.*

(ii) *Let α, β be shifts on $B(H)$ defined by $\alpha(A) = V A V^*$ and $\beta(A) = W A W^*$, where V and W are isometries on H . Then α is conjugate to β if and only if V is unitarily equivalent to λW for some $\lambda \in S^1$ i.e. if and only if V and W have the same Wold decomposition.*

Proof. (i) By Proposition 2.4, the conditions on V are necessary for α to be a shift. Conversely, if V satisfies the given conditions, then V can be assumed to be a direct sum of copies of the unilateral shift and (possibly) a unitary map on a

one-dimensional subspace. Let $\{e_i: i \in \mathbb{N}\}$ be the orthonormal basis associated with one of the shift summands. If $T \in \alpha'(B(H))$ then $T = V'SV^{*r}$ for some $S \in B(H)$ and hence $Te_i = 0$ for $1 \leq i \leq r$. It follows that, if $T \in \bigcap_r \alpha'(B(H))$ then the restriction of T to $(\bigcap_r V'H)^\perp$ is 0. The same argument applies to T^* and hence T leaves $\bigcap_r V'H$ globally invariant. Therefore $\bigcap_r \alpha'(B(H)) = \mathbb{C}P$ where P is the projection from H onto the zero or one-dimensional space $\bigcap_r V'H$.

(ii) This follows from Proposition 2.2. $\square$

The case in which E_α has dimension one is very special. For example, in contrast to the result of Proposition 2.5(i), if E_α has dimension greater than one, then each isometry V in E_α is onto a space of infinite codimension and hence must be equivalent to an infinite direct sum of copies of the unilateral shift, possibly with an additional one-dimensional unitary summand.

Later in the paper an example will be produced to show that it may occur that $\alpha(A) = \sum V_n A V_n^* = \sum W_n A W_n^*$ where all the V_n are pure and one of the W_n is not. It will also be shown that the condition that all the V_n are pure is not sufficient to ensure that α is a shift. Both these examples rely on the notion of a product shift, that will now be introduced.

3. PRODUCT SHIFTS

Let K be a Hilbert space and, for each $i \in \mathbb{N}$, let $k_i \in K$ be a unit vector such that $\sum_i |1 - \langle k_i, k_{i+1} \rangle| < \infty$. Recall that, as described in [3] or [6], the incomplete infinite tensor product $\bigotimes^{k_i} K$ is the Hilbert space inductive limit of the sequence $K \rightarrow K \otimes K \rightarrow K \otimes K \otimes K \cdots$, where the i th map takes h to $h \otimes k_{i+1}$. Let ϕ_i be the natural map from the i -fold tensor product $K \otimes K \otimes \cdots \otimes K$ into $\bigotimes^{k_i} K$ and let ψ_i be the corresponding map from $K \otimes K \otimes \cdots \otimes K$ into $\bigotimes^{k_{i+1}} K$. Then, by [3, Proposition 1.1], for each h in the r -fold tensor product $K \otimes K \otimes \cdots \otimes K$, the sequence $\phi_r(h), \phi_{r+1}(h \otimes k_{r+2}), \phi_{r+2}(h \otimes k_{r+2} \otimes k_{r+3}), \dots$ has a limit in $\bigotimes^{k_i} K$ and, by [3, Proposition 1.3], the map taking $\psi_r(h)$ to this limit extends to an isomorphism θ from $\bigotimes^{k_{i+1}} K$ onto $\bigotimes^{k_i} K$. Informally, we can describe θ as taking $\bigotimes x_i$ (where $x_i = k_{i+1}$ for almost all i) to $\bigotimes x_i$ (interpreting the last expression as the limit of $\phi_1(x_1), \phi_2(x_1 \otimes x_2), \dots$).

Proposition 3.1. *Let P be a projection on K ; k_i be a sequence of unit vectors in K with $\sum |1 - \langle k_i, k_{i+1} \rangle| < \infty$; and θ be the isomorphism from $\bigotimes^{k_{i+1}} K$ onto $\bigotimes^{k_i} K$ taking $\bigotimes x_i$ (where $x_i = k_{i+1}$ for almost all i) to $\bigotimes x_i$. Then the map $\alpha_{k,P}$ defined on $B(H) = B(\bigotimes^{k_i} K)$ by $\alpha_{k,P}(T) = \theta(P \otimes T)\theta^{-1}$ is a shift.*

Proof. The map $\alpha_{k,P}$ is clearly a $*$ -endomorphism. In the unital case, $\alpha_{k,1}'(B(H))' = B(K \otimes \cdots \otimes K) \otimes \mathbb{C}1$ (where the tensor product has r factors) and hence $\bigcup_r \alpha_{k,1}'(B(H))'$ is weakly dense in $B(H)$, from which it follows that $\alpha_{k,1}$ is a shift. In the nonunital case let q be the greatest lower bound

of the projections $\alpha'_{k,p}(1)$. If $q = 0$ then, since $\alpha'_{k,p}(1)T = T$ for each $T \in \bigcap_r \alpha'_{k,p}(B(H))$, $T = qT = 0$ for all such T and hence $\alpha_{k,p}$ is a shift. If $q \neq 0$ then $q\alpha'_{k,1}(T)q = q\alpha'_{k,p}(T)q$ for all T because, as can be checked by induction, $\alpha'_{k,p}(1)\alpha'_{k,1}(T)\alpha'_{k,p}(1) = \alpha'_{k,p}(T)$. Thus, since $\alpha_{k,p}(q) = q$, $\alpha'_{k,p}(qB(H)q) = q\alpha'_{k,p}(B(H))q = q\alpha'_{k,1}(B(H))q$ and so, when both algebras are restricted to qH , $[\alpha'_{k,p}(qB(H)q)]' = q\alpha'_{k,1}(B(H))'q$. Hence, as in the unital case, $\bigcap_r \alpha'_{k,p}(qB(H)q) = \mathbb{C}q$; however, $\bigcap_r \alpha'_{k,p}(B(H)) \subseteq qB(H)q$ and, therefore, $\bigcap_r \alpha'_{k,p}(B(H)) = \mathbb{C}q$, as required. $\square$

If there exists a unitary U from a Hilbert space H onto an infinite tensor product $\bigotimes^{k_i} K$ with the properties above, then any $*$ -endomorphism α of $B(H)$ of the form $(\text{Ad } U^*)\alpha_{k,p}(\text{Ad } U)$ will be called a *product shift* on $B(H)$.

Not every shift on $B(H)$ is a product as can be seen from Proposition 2.4. Any product shift $\alpha_{k,p}$ has $\alpha_{k,p}(1)$ of infinite codimension whereas if α is the shift corresponding to a single unilateral shift V then $\alpha(1)$ has codimension 1. Nevertheless the author is not aware of examples of nonproduct unital shifts on $B(H)$: the following proposition shows that any sequence of isometries defining such a shift must be all pure.

Proposition 3.2. *Let α be a unital shift on $B(H)$ defined by $\alpha(A) = \sum_n V_n A V_n^*$ where $V = V_1$ is not pure. Then α is a product shift.*

Proof. Let h be a unit vector in $\bigcap_r V^r H$ and note that, by Proposition 2.4, $\bigcap_r V^r H$ is one-dimensional. It follows that $Vh = \lambda h$ for some $\lambda \in S^1$ and, replacing V by $\bar{\lambda}V$, we can assume that $Vh = h$.

Let c_{00} denote the space of functions $f: \mathbb{N} \rightarrow \mathbb{N}$ such that $f(r) = 1$ for all sufficiently large r and, for each $f \in c_{00}$, let $b_f = \phi_s(V_{f(1)} \otimes \cdots \otimes V_{f(s)}) \in \bigotimes^V E_\alpha$, where ϕ_s is the natural map from the s -fold tensor product $E_\alpha \otimes \cdots \otimes E_\alpha$ into $\bigotimes^V E_\alpha$ and where s is chosen sufficiently large so that $f(r) = 1$ for all $r \geq s$. Then, by [6, Lemma 4.1.4], $\{b_f: f \in c_{00}\}$ is an orthonormal basis for $\bigotimes^V E_\alpha$. Define $\psi: \bigotimes^V E_\alpha \rightarrow H$ extending $\psi(b_f) = V_{f(1)} \cdots V_{f(s)} h$ (which is well defined since $Vh = h$). Then, for $t \geq s$, $\langle \psi(b_f), \psi(b_g) \rangle = \langle V_{f(1)} \cdots V_{f(s)} h, V_{g(1)} \cdots V_{g(t)} h \rangle = \delta_{f(1)g(1)} \cdots \delta_{f(s)g(s)} \langle V_{g(t)}^* \cdots V_{g(s+1)}^* h, h \rangle = \delta_{fg} \langle h, h \rangle = \delta_{fg}$ so that ψ defines a unitary mapping from $\bigotimes^V E_\alpha$ onto a subspace K of H .

For each μ, ν with $|\mu| \geq 0$ and $|\nu| = r \geq 0$, $V_\mu V_\nu^* h = V_\mu V_\nu^* V^r h = \delta_{\nu(1, \dots, 1)} V_\mu h$, so that K is equal to the closed linear span of the vectors $V_\mu V_\nu^* h$ with $|\mu| \geq 0$ and $|\nu| \geq 0$. By Lemma 2.3 (i) it follows that the projection onto K belongs to $\bigcap_r \alpha'(B(H))$ and hence, since α is a unital shift, $K = H$.

Note that

$$\begin{aligned} (\psi^* V_n \psi)(\phi_r(V_{f(1)} \otimes \cdots \otimes V_{f(r)})) &= \psi^* V_n V_{f(1)} \cdots V_{f(r)} h \\ &= \phi_{r+1}(V_n \otimes V_{f(1)} \otimes \cdots \otimes V_{f(r)}), \end{aligned}$$

from which it follows that

$$(\psi^* V_n^* \psi)(\phi_r(V_{f(1)} \otimes \cdots \otimes V_{f(s)})) = \langle V_{f(1)}, V_n \rangle \phi_{r-1}(V_{f(2)} \otimes \cdots \otimes V_{f(r)});$$

hence

$$\begin{aligned} & (\text{Ad } \psi^*)\alpha(\text{Ad } \psi)(T_1 \otimes T_2 \otimes \cdots \otimes T_s \otimes 1) \\ &= \sum_n (\psi^* V_n \psi)(T_1 \otimes \cdots \otimes T_s \otimes 1)(\psi^* V_n \psi)^* = 1 \otimes T_1 \otimes \cdots \otimes T_s \otimes 1 \end{aligned}$$

and thus that α is a product shift. $\square$

It will now be shown how product shifts can be defined to illustrate the comments made after Proposition 2.5.

Example 3.3. Let $K = \mathbb{C}^2$; $k_i = (1, 0)$ for each i ; and the isometries V_1, V_2, W_1, W_2 be defined on $H = \bigotimes^{k_i} K$ by

$$\begin{aligned} V_1(h) &= (1/\sqrt{2}, 1/\sqrt{2}) \otimes h, & V_2(h) &= (1/\sqrt{2}, -1/\sqrt{2}) \otimes h, \\ W_1(h) &= (1, 0) \otimes h, & W_2(h) &= (0, 1) \otimes h. \end{aligned}$$

Then $\alpha_{k,1}(A) = V_1 A V_1^* + V_2 A V_2^* = W_1 A W_1^* + W_2 A W_2^*$ for each $A \in B(H)$. However, both V_1 and V_2 are pure isometries (as is W_2) but $\bigcap_n W_1^n H = \mathbb{C}k$, where $k = \bigotimes(1, 0)$. $\square$

Example 3.4. Let H, V_1, V_2 , and $\alpha_{k,1}$ be as defined in Example 3.3 and let $\mathcal{H}$ be any Hilbert space. Then $\alpha = \alpha_{k,1} \otimes \text{id}$ on $B(H) \otimes B(\mathcal{H})$ is not a shift; however,

$$\alpha(A) = (V_1 \otimes 1)A(V_1^* \otimes 1) + (V_2 \otimes 1)A(V_2^* \otimes 1)$$

for each A , and both $V_1 \otimes 1$ and $V_2 \otimes 1$ are pure isometries. Hence the converse of Proposition 2.4 is false. $\square$

The next main result gives a criterion for two unital product shifts to be conjugate. The proof is based on the following lemma:

Lemma 3.5. Let $W: \bigotimes^{k_i} K \rightarrow \bigotimes^{h_i} K$ be a unitary for which $(\text{Ad } W)\alpha_{k,1}(\text{Ad } W^*) = \alpha_{h,1}$. Then there exists a unitary V on K such that, for each $r \geq 1$ and each $T_1, \dots, T_r \in B(K)$, $(\text{Ad } W)(T_1 \otimes \cdots \otimes T_r \otimes 1) = (V T_1 V^*) \otimes (V T_2 V^*) \otimes \cdots \otimes (V T_r V^*) \otimes 1$.

Proof. For notational convenience, let $H_h = \bigotimes^{h_i} K$ and $H_k = \bigotimes^{k_i} K$. Then $\alpha_{h,1}(B(H_h))' = \text{Ad}(W)[\alpha_{k,1}(B(H_k))']$ and hence, for each $T \in B(K)$,

$$(\text{Ad } W)(T \otimes 1 \otimes 1 \otimes \cdots) = V T V^* \otimes 1 \otimes 1 \otimes \cdots$$

for some unitary V on K . Then, for each $r \geq 0$,

$$\begin{aligned} (\text{Ad } W)\alpha_{k,1}^r(T \otimes 1 \otimes 1 \otimes \cdots) &= \alpha_{h,1}^r(\text{Ad } W)(T \otimes 1 \otimes 1 \otimes \cdots) \\ &= \alpha_{h,1}^r(V T V^* \otimes 1 \otimes 1 \otimes \cdots). \end{aligned}$$

Hence, for each $r \geq 0$, $(\text{Ad } W)(T_1 \otimes \cdots \otimes T_r \otimes 1) = (VT_1V^*) \otimes (VT_2V^*) \otimes \cdots \otimes (VT_rV^*) \otimes 1$. $\square$

Theorem 3.6. *The product shifts $\alpha_{k,1}$ and $\alpha_{h,1}$ are conjugate if and only if there exists a unitary map V on K with $\sum |1 - |\langle Vk_i, h_i \rangle|| < \infty$.*

Proof. If $\sum |1 - |\langle Vk_i, h_i \rangle|| < \infty$ then, by [3, Proposition 1.3], the map taking $\bigotimes x_i$ (where $x_i = k_i$ for almost all i) to $\bigotimes x_i$ extends to an isomorphism from $\bigotimes^{k_i} K$ onto $\bigotimes^{\alpha_i V^* h_i} K$, where α_i is the complex number of unit modulus such that $|\langle k_i, V^* h_i \rangle| = \langle k_i, \alpha_i V^* h_i \rangle$. However there is an isomorphism from $\bigotimes^{\alpha_i V^* h_i} K$ onto $\bigotimes^{h_i} K$ taking $\bigotimes x_i$ to $\bigotimes \bar{\alpha}_i V x_i$ and hence there is an isomorphism W from $\bigotimes^{k_i} K$ onto $\bigotimes^{h_i} K$ taking $\bigotimes x_i$ (where $x_i = k_i$ for almost all i) to $\bigotimes \bar{\alpha}_i V x_i$. Then, for $T_1, \dots, T_r \in B(K)$,

$$(\text{Ad } W)\alpha_{k,1}(\text{Ad } W^*)(T_1 \otimes \cdots \otimes T_r \otimes 1) = \alpha_{h,1}(T_1 \otimes \cdots \otimes T_r \otimes 1),$$

from which it follows, by linearity and continuity that $\alpha_{k,1}$ is conjugate to $\alpha_{h,1}$.

Conversely, if $\alpha_{k,1}$ is conjugate to $\alpha_{h,1}$ then, by Lemma 3.5, there exists an isomorphism W from $\bigotimes^{k_i} K$ onto $\bigotimes^{h_i} K$ and a unitary V on K such that $(\text{Ad } W)(T_1 \otimes \cdots \otimes T_r \otimes 1) = (VT_1V^*) \otimes \cdots \otimes (VT_rV^*) \otimes 1$ for all $T_1, \dots, T_r \in B(K)$. In particular, this equation holds when T_i is the one-dimensional projection onto the subspace spanned by k_i . In this case, the decreasing sequence of projections $T_1 \otimes \cdots \otimes T_r \otimes 1$ is weakly convergent as $r \rightarrow \infty$; by considering the effect of $T_1 \otimes \cdots \otimes T_r \otimes 1$ on an orthonormal basis $\{e_{1i_1} \otimes e_{2i_2} \otimes \cdots : i_1, i_2, \dots \in \mathbb{N}\}$ (where $e_{r1} = k_r$ for each r and $i_k = 1$ for all sufficiently large k) it can be seen that the weak limit is the one-dimensional projection onto the subspace spanned by $k = \bigotimes k_i$. Hence the projections $(VT_1V^*) \otimes \cdots \otimes (VT_rV^*) \otimes 1$ are weakly convergent to a one-dimensional projection in $\bigotimes^{h_i} K$; let x be a unit vector in the corresponding one-dimensional subspace. If ϕ_r denotes the natural map from the r -fold tensor product $K \otimes \cdots \otimes K$ into $\bigotimes^{h_i} K$ then there exists $s \in \mathbb{N}$ and a unit vector x_s in $\phi_s(K \otimes \cdots \otimes K)$ such that $\|x - x_s\| < \frac{1}{2}$. Then, for each r ,

$$\begin{aligned} & \|[(VT_1V^*) \otimes \cdots \otimes (VT_rV^*) \otimes 1]x_s\| \\ & \geq \|x\| - \|x - [(VT_1V^*) \otimes \cdots \otimes (VT_rV^*) \otimes 1]x_s\| > \frac{1}{2}. \end{aligned}$$

It follows that $\prod_{i=s+1}^r |\langle k_i, V^* h_i \rangle| \not\rightarrow 0$ as $r \rightarrow \infty$ and hence, by Lemma 2.4.1 (II) of [6], $\sum |1 - |\langle k_i, V^* h_i \rangle|| < \infty$, as required. $\square$

Theorem 3.6 can be used to produce examples of nonconjugate but outer-conjugate unital shifts. Let $\log^k$ denote the k -fold composite $\log \circ \cdots \circ \log$ and, for each $r \in \mathbb{N}$, define a sequence f_r by

$$f_r(n) = \begin{cases} 1 & \text{if } \log^r(n) \leq 0, \\ 1/(n \log(n) \log^2(n) \cdots \log^r(n)) & \text{if } \log^r(n) > 0. \end{cases}$$

Let K be a separable Hilbert space (of dimension at least 2) with an orthonormal basis $\{e_i\}$ and, for each $n \in \mathbb{N}$, define $k_{r,n} \in K$ by $k_{r,n} = (1 - f_r(n))e_1 + [f_r(n)(2 - f_r(n))]^{1/2}e_2$.

Lemma 3.7. (i) $\|k_{r,n}\| = 1$ for each r and n .

(ii) $\sum_n |1 - \langle k_{r,n}, k_{r,n+1} \rangle| < \infty$ for each r .

Proof. Claim (i) is easily verified. To check claim (ii), note that

$$0 \leq \langle k_{r,n}, k_{r,n+1} \rangle \leq \|k_{r,n}\| \|k_{r,n+1}\| = 1$$

and hence that

$$\begin{aligned} 0 &\leq |1 - \langle k_{r,n}, k_{r,n+1} \rangle| \\ &= f_r(n) + f_r(n+1) - f_r(n)f_r(n+1) \\ &\quad - [f_r(n)f_r(n+1)(2 - f_r(n))(2 - f_r(n+1))]^{1/2} \\ &\leq 2f_r(n) - f_r(n)f_r(n+1) - f_r(n+1)(2 - f_r(n)) \\ &= 2(f_r(n) - f_r(n+1)). \end{aligned}$$

However $\sum_n 2(f_r(n) - f_r(n+1))$ is convergent, from which the result follows. $\square$

Proposition 3.8. For each $r \in \mathbb{N}$, let α_r be the product shift $\alpha_{k_{r,1}}$ on $\bigotimes_{n=1}^{\infty} k_{r,n} K$, where $k_{r,n}$ is defined above. Then, if $r \neq s$, α_r is outer conjugate but not conjugate to α_s .

Proof. For definiteness suppose that $r > s$. By construction α_r and α_s have the same multiplicity $\dim(K)$ and hence are outer conjugate by [7, Theorem 2.4]. If they are conjugate then, by Theorem 3.6, there exists a unitary V on K such that $\sum_n |1 - |\langle V k_{r,n}, k_{s,n} \rangle|| < \infty$. It follows that $|\langle V k_{r,n}, k_{s,n} \rangle| \rightarrow 1$ as $n \rightarrow \infty$ and hence that $|\langle V e_1, e_1 \rangle| = 1$, so that $V e_1 = \alpha e_1$ for some $\alpha \in S^1$. Hence,

$$\begin{aligned} |\langle V k_{r,n}, k_{s,n} \rangle| &= |(1 - f_r(n))(1 - f_s(n))\langle \alpha e_1, e_1 \rangle \\ &\quad + [f_r(n)f_s(n)(2 - f_r(n))(2 - f_s(n))]^{1/2} \langle V e_2, e_2 \rangle| \\ &\leq (1 - f_r(n))(1 - f_s(n)) + 2f_s(n)^{1/2} f_r(n)^{1/2} \\ &= 1 - f_s(n)[1 - f_r(n) + f_r(n)f_s(n)^{-1} - 2f_r(n)^{1/2} f_s(n)^{-1/2}]. \end{aligned}$$

However, for all sufficiently large n , $f_r(n)f_s(n)^{-1} = 1/(\log^{s+1}(n) \cdots \log^r(n)) \rightarrow 0$ as $n \rightarrow \infty$ and therefore, for all sufficiently large n , $|\langle V k_{r,n}, k_{s,n} \rangle| \leq 1 - f_s(n)/2$ and therefore, $\sum |1 - |\langle V k_{r,n}, k_{s,n} \rangle||$ is not convergent, giving a contradiction. $\square$

REFERENCES

1. R. Archbold, *On the 'flip-flop' automorphism of $C^*(S_1, S_2)$* , Quart. J. Math. Oxford (2), **30** (1979), 129–132.
2. W. Arveson, *Continuous analogues of Fock space*, Mem. Amer. Math. Soc., no. 409, Amer. Math. Soc., Providence, RI, 1989.

3. A. Guichardet, *Produits tensoriels infinis et représentations des relations d'anticommutation*, Ann. Sci. Ecole Norm. Sup. (3) **83** (1966), 1–52.
4. P. R. Halmos, *A Hilbert space problem book*, Springer-Verlag, Berlin and New York, 1974.
5. H. Hanche-Olsen and E. Størmer, *Jordan operator algebras*, Pitman, 1984.
6. J. von Neumann, *On infinite direct products*, Compositio Math. **6** (1938), 1–77.
7. R. T. Powers, *An index theory for semigroups of $*$ -endomorphisms of $B(H)$ and type II_1 factors*, Canad. J. Math. **40** (1988), 86–114.

DEPARTMENT OF MATHEMATICS, LA TROBE UNIVERSITY, BUNDOORA, VICTORIA 3083, AUSTRALIA