“LEBESGUE MEASURE” ON $\mathbf{R}^\infty$

RICHARD BAKER

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. We construct a translation invariant Borel measure λ on $\mathbf{R}^\infty = \prod_{i=1}^\infty \mathbf{R}$ such that for any infinite-dimensional rectangle $R = \prod_{i=1}^\infty (a_i, b_i)$, $-\infty < a_i \leq b_i < +\infty$, if $0 \leq \prod_{i=1}^\infty (b_i - a_i) < +\infty$, then $\lambda(R) = \prod_{i=1}^\infty (b_i - a_i)$. Because $\mathbf{R}^\infty$ is an infinite-dimensional locally convex topological vector space, the measure λ can not be σ -finite.

1. INTRODUCTION

If $\mathcal{X}$ is any infinite-dimensional locally convex topological vector space, then it is known that there does not exist a nontrivial translation invariant σ -finite Borel measure on $\mathcal{X}$ [2, p. 143]. Because of this impossibility theorem, it is part of the folklore of infinite-dimensional functional analysis that there is no satisfactory analogue of Lebesgue measure on the space $\mathbf{R}^\infty = \prod_{i=1}^\infty \mathbf{R}$. In this paper, without upsetting this folklore, we show that there exists a nontrivial translation invariant Borel measure λ on $\mathbf{R}^\infty$ which is analogous to Lebesgue measure in the sense that if $R = \prod_{i=1}^\infty (a_i, b_i)$ is any infinite-dimensional rectangle such that the “volume” $\prod_{i=1}^\infty (b_i - a_i)$ of R is a nonnegative real number, then

$$\lambda(R) = \prod_{i=1}^\infty (b_i - a_i).$$

Therefore, it would be appropriate to call λ *infinite-dimensional Lebesgue measure on $\mathbf{R}^\infty$* . Of course, according to the above no-go result, the measure λ can not be σ -finite.

The following theorem is the main result of this paper.

Theorem I. *Let $\mathcal{R}$ be the class of all infinite-dimensional rectangles $R \in \mathbf{R}^\infty$ of the form*

$$R = \prod_{i=1}^\infty (a_i, b_i), \quad -\infty < a_i \leq b_i < +\infty,$$

Received by the editors June 1, 1990.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 28A35.

©1991 American Mathematical Society
 0002-9939/91 \$1.00 + \$.25 per page

such that $0 \leq \prod_{i=1}^{\infty} (b_i - a_i) < +\infty$. Let τ be the set of function on $\mathcal{R}$ defined by

$$\tau(R) = \prod_{i=1}^{\infty} (b_i - a_i), \quad R \in \mathcal{R}.$$

Finally, let λ be the set function defined on all subsets of $\mathbf{R}^{\infty}$ by

$$\lambda(E) = \inf_{\substack{R_j \in \mathcal{R} \\ \bigcup R_j \supseteq E}} \sum_{j=1}^{\infty} \tau(R_j), \quad E \subseteq \mathbf{R}^{\infty}.$$

We adopt the convention that any infimum taken over an empty set of real numbers has the value $+\infty$. Then λ is a translation invariant Borel measure on $\mathbf{R}^{\infty}$ such that for all $R = \prod_{i=1}^{\infty} (a_i, b_i) \in \mathcal{R}$, we have

$$\lambda(R) = \prod_{i=1}^{\infty} (b_i - a_i).$$

The plan of the paper is as follows. In §2 we present the preliminary measure-theoretic machinery that we will need for the construction of λ . In §3 we prove Theorem I.

2. PRELIMINARIES

The purpose of this section is to collect together in one place the measure-theoretic facts that we will need for the construction of the measure λ . All of the material in this section has been transcribed from Chapter 1 of [1].

Throughout this section Ω will denote an arbitrary nonempty set. Definition 2.1 and Theorem 2.2 below give a general method for constructing outer measures on Ω .

Definition 2.1. A function τ on a class $\mathcal{C}$ of subsets of Ω will be call a *pre-measure*, if:

- (a) $\emptyset \in \mathcal{C}$;
- (b) $0 \leq \tau(C) \leq +\infty$ for all C in $\mathcal{C}$;
- (c) $\tau(\emptyset) = 0$.

Theorem 2.2. If τ is a pre-measure defined on a class $\mathcal{C}$ of subsets of Ω , the set function

$$\mu(E) = \inf_{\substack{C_j \in \mathcal{C} \\ \bigcup C_j \supseteq E}} \sum_{j=1}^{\infty} \tau(C_j)$$

is an outer measure on Ω .

Theorem 2.3. Let μ be an outer measure on Ω . Then μ is the outer measure, constructed via Theorem 2.2, from the pre-measure τ , defined on the class of all subsets of Ω to coincide with μ .

Now assume that Ω is a metric space with metric ρ . The next theorem gives another general method of constructing an outer measure on Ω .

Theorem 2.4. If τ is a pre-measure defined in a class $\mathcal{C}$ of sets, in a metric space Ω with metric ρ , the set function

$$\mu(E) = \sup_{\delta > 0} \mu_\delta(E),$$

where

$$\mu_\delta(E) = \inf_{\substack{C_j \in \mathcal{C}, \\ \bigcup C_j \supseteq E}} \sum_{j=1}^{\infty} \tau(C_j) \quad d(C_j) \leq \delta$$

is an outer measure on Ω . Here $d(C)$ is the diameter of C with respect to ρ , $C \subseteq \mathbf{R}^\infty$.

The last two theorems of this section, Theorems 2.7 and 2.8, state that the outer measure μ constructed in Theorem 2.4 has the desirable property that every Borel subset of Ω is μ -measurable.

Definition 2.5. If A and B are disjoint nonempty sets in a metric space Ω with metric ρ , A and B are said to be *positively separated* if the distance

$$\inf_{a \in A, b \in B} \rho(a, b)$$

separating A and B is positive.

Definition 2.6. An outer measure μ defined on a metric space Ω is called a *metric outer measure*, if

$$\mu(A \cup B) = \mu(A) + \mu(B),$$

for every pair of disjoint nonempty sets A, B that are positively separated.

Theorem 2.7. if μ is a metric outer measure on a metric space Ω , then all Borel sets in Ω are μ -measurable. Recall that a subset $E \subseteq \Omega$ is μ -measurable if for all sets $A \subseteq E$, $B \subseteq \Omega \setminus E$, we have $\mu(A \cup B) = \mu(A) + \mu(B)$; moreover, the system $\mathcal{M}$ of μ -measurable sets is a σ -field containing the null sets, and the restriction of μ to $\mathcal{M}$ is countably additive on $\mathcal{M}$.

Theorem 2.8. If τ is pre-measure defined on a class $\mathcal{C}$ of subsets of a metric space Ω , and if μ is the outer measure on Ω constructed from τ via Theorem 2.4, then μ is a metric outer measure on Ω .

3. THE MEASURE λ ON $\mathbf{R}^\infty$

In this section we construct an outer measure λ on $\mathbf{R}^\infty$ such that λ has all the properties stated in Theorem I.

Let $\mathcal{R}$, τ , and λ be defined as in Theorem I. Then by Theorem 2.2, λ is an outer measure on $\mathbf{R}^\infty$. The first two theorems of this section establish the key fact that for all $R \in \mathcal{R}$, $\tau(R) = \lambda(R)$.

Theorem 3.1. Let $I = \prod_{i=1}^{\infty} [a_i, b_i]$, $-\infty < a_i \leq b_i < +\infty$, be an infinite-dimensional compact rectangle in $\mathbf{R}^{\infty}$ such that $0 \leq \prod_{i=1}^{\infty} (b_i - a_i) < +\infty$, then

$$\prod_{i=1}^{\infty} (b_i - a_i) \leq \lambda(I).$$

Proof. It will suffice to show that if $I \subseteq \bigcup_{j=1}^{\infty} R_j$, where $R_j \in \mathcal{R}$, then

$$(*) \quad \tau(I) \leq \sum_{j=1}^{\infty} \tau(R_j),$$

where $\tau(I) = \prod_{i=1}^{\infty} (b_i - a_i)$. We may assume that $0 < \tau(I)$ and $\sum_{j=1}^{\infty} \tau(R_j) < \infty$. For all $j, n \geq 1$, let

$$R_j = \prod_{i=1}^{\infty} (a_{ij}, b_{ij}), \quad R_{nj} = \prod_{i=1}^n (a_{ij}, b_{ij}) \times \prod_{i=n+1}^{\infty} \mathbf{R}.$$

Now let $0 < \varepsilon < 1$ be arbitrary, and let $\mathcal{F}$ be the family of all those R_{nj} such that $\prod_{i=n+1}^{\infty} (b_i - a_i) < 1 + \varepsilon$ and either $\prod_{i=n+1}^{\infty} (b_{ij} - a_{ij}) > 1 - \varepsilon$ or $\prod_{i=1}^n (b_{ij} - a_{ij}) < \varepsilon/2^j$. Because $0 < \tau(I) < +\infty$ and $0 \leq \tau(R_j) < +\infty$, it is clear that $\mathcal{F}$ covers I . The members of $\mathcal{F}$ are open and I is compact, hence there exists a finite subfamily $\{R_{n_p j_p} | 1 \leq p \leq k\}$ of $\mathcal{F}$ that covers I . For $n > \max\{n_1, \dots, n_k\}$ and $1 \leq p \leq k$, let

$$I_n = \prod_{i=1}^n [a_i, b_i], \quad S_{np} = \prod_{i=1}^{n_p} (a_{ij_p}, b_{ij_p}) \times \prod_{i=n_p+1}^n [a_i, b_i].$$

Because $I \subseteq \bigcup_{p=1}^k R_{n_p j_p}$, we have $I_n \subseteq \bigcup_{p=1}^k S_{np}$. Let λ_n be Lebesgue measure on $\mathbf{R}^n$, then we have

$$\begin{aligned} \prod_{i=1}^n (b_i - a_i) &= \lambda_n(I_n) \leq \sum_{p=1}^k \lambda_n(S_{np}) \\ &= \sum_{p=1}^k \left\{ \prod_{i=1}^{n_p} (b_{ij_p} - a_{ij_p}) \cdot \prod_{i=n_p+1}^n (b_i - a_i) \right\}. \end{aligned}$$

Taking the limit $n \rightarrow \infty$, we get

$$\begin{aligned} \tau(I) &\leq \sum_p^k \left\{ \prod_{i=1}^{n_p} (b_{ij_p} - a_{ij_p}) \cdot \prod_{i=n_p+1}^{\infty} (b_i - a_i) \right\} \\ &\leq (1 + \varepsilon) \sum_{p=1}^k \prod_{i=1}^{n_p} (b_{ij_p} - a_{ij_p}) \\ &\leq (1 + \varepsilon) \sum'_{1 \leq p \leq k} \prod_{i=1}^{n_p} (b_{ij_p} - a_{ij_p}) + (1 + \varepsilon) \sum''_{1 \leq p \leq k} \prod_{i=1}^{n_p} (b_{ij_p} - a_{ij_p}), \end{aligned}$$

where $\sum'$ is the sum over those p for which $\prod_{i=n_p+1}^\infty (b_{ij_p} - a_{ij_p}) > 1 - \varepsilon$ and $\sum''$ is the sum over those p for which $\prod_{i=1}^{n_p} (b_{ij_p} - a_{ij_p}) < \varepsilon/2^{j_p}$. It follows that

$$\begin{aligned} \tau(I) &\leq \left(\frac{1+\varepsilon}{1-\varepsilon} \right) \sum'_{1 \leq p \leq k} \tau(R_{j_p}) + (1+\varepsilon)\varepsilon \sum''_{1 \leq p \leq k} (1/2^{j_p}) \\ &\leq \left(\frac{1+\varepsilon}{1-\varepsilon} \right) \sum_{j=1}^\infty \tau(R_j) + (1+\varepsilon)\varepsilon. \end{aligned}$$

Because ε is arbitrary, we see that $(*)$ holds. $\square$

Theorem 3.2. For every $R \in \mathcal{R}$, we have $\tau(R) = \lambda(R)$.

Proof. Let $R \in \mathcal{R}$. It is clear that $\lambda(R) \leq \tau(R)$, and hence we may assume that $\tau(R) > 0$. Let $0 < \varepsilon < 1$ be arbitrary. Then it is not hard to see that there exists a compact rectangle $I = \prod_{i=1}^\infty [a_i, b_i] \subseteq R$ such that $\prod_{i=1}^\infty (b_i - a_i) = (1 - \varepsilon)\tau(R)$. By Theorem 3.1, $\prod_{i=1}^\infty (b_i - a_i) \leq \lambda(I)$, therefore we have $(1 - \varepsilon)\tau(R) \leq \lambda(R)$. Because ε is arbitrary, this proves the theorem. $\square$

Definition 3.3. Let ρ be the metric on $\mathbf{R}^\infty$ defined as

$$\rho(x, y) = \sum_{i=1}^\infty \frac{(1/2^i)|x_i - y_i|}{\{1 + |x_i - y_i|\}}, \quad x = (x_i), y = (y_i) \in \mathbf{R}^\infty.$$

Observe that ρ induces the product topology on $\mathbf{R}^\infty$.

Theorems 3.4 and 3.5 below, show that λ has the important property of being a metric outer measure on $\mathbf{R}^\infty$.

Theorem 3.4. Let ν be the outer measure on $\mathbf{R}^\infty$ constructed from the pair τ, ρ via Theorem 2.4. Then for all $R \in \mathcal{R}$, we have $\nu(R) = \tau(R)$.

Proof. Let $R \in \mathcal{R}$. By Theorem 3.2, $\lambda(R) = \tau(R)$. It is clear that $\lambda(R) \leq \nu(R)$, hence it will suffice to show that

$$(**) \quad \nu(R) \leq \tau(R).$$

Write

$$R = \prod_{i=1}^\infty (a_i, b_i), \quad -\infty < a_i \leq b_i < +\infty.$$

If $R = \emptyset$, then $\nu(R) = \tau(R) = 0$, hence we may assume that $R \neq \emptyset$. Then for all i , $a_i < b_i$. Therefore, if $\tau(R) = 0$, then for all n , we have $\prod_{i=n+1}^\infty (b_i - a_i) = 0$. On the other hand, if $\tau(R) > 0$, then $\lim_{n \rightarrow \infty} \prod_{i=n+1}^\infty (b_i - a_i) = 1$. Let $\delta, \varepsilon > 0$ be arbitrary, then there exists an $n \geq$ such that $\prod_{i=n+1}^\infty (b_i - a_i) < 1 + \varepsilon$ and $\sum_{i=1+n}^\infty 2^{-i} < \delta/2$. Define $R_n = \prod_{i=1}^n (a_i, b_i)$. For $x = (x_i)_{i=1}^n, y = (y_i)_{i=1}^n \in \mathbf{R}^n$, let $\rho_n(x, y)$ be defined as

$$\rho_n(x, y) = \sum_{i=1}^n \left(\frac{1}{2^i} \right) \frac{|x_i - y_i|}{\{1 + |x_i - y_i|\}}.$$

Cover the rectangle R_n by rectangles $R_{n1}, \dots, R_{nm}$ in $\mathbf{R}^n$ such that

- (a) For $1 \leq j \leq m$, $R_{nj} = \prod_{i=1}^n (a_{ij}, b_{ij})$, $-\infty < a_{ij} \leq b_{ij} < +\infty$.
- (b) For $1 \leq j \leq m$, $\sup_{x, y \in R_{nj}} \|x - y\| < \delta/2n$, where $\|\cdot\|$ is the Euclidean norm on $\mathbf{R}^n$.
- (c) $\sum_{j=1}^m \lambda_n(R_{nj}) < \prod_{i=1}^n (b_i - a_i) + \varepsilon$.

For $1 \leq j \leq m$, define $R_j = R_{nj} \times \prod_{i=n+1}^\infty (a_i, b_i)$. Let $x = (x_i)$, $y = (y_i) \in R_j$, and define $x^{(n)} = (x_i)_{i=1}^n$, $y^{(n)} = (y_i)_{i=1}^n$, then, by (b), we have

$$\begin{aligned} \rho_n(x, y) &= \rho_n(x^{(n)}, y^{(n)}) + \sum_{i=n+1}^\infty \left(\frac{1}{2^i} \right) \frac{|x_i - y_i|}{\{1 + |x_i - y_i|\}} \\ &< \rho_n(x^{(n)}, y^{(n)}) + \delta/2 \\ &\leq n \|x^{(n)} - y^{(n)}\| + \delta/2 \\ &< \delta. \end{aligned}$$

Hence, for all $1 \leq j \leq m$, we have $d(R_j) < \delta$. It is clear that $R \subseteq \bigcup_{j=1}^m R_j$, hence by definition, $\nu_\delta(R) \leq \sum_{j=1}^m \tau(R_j)$. By (c),

$$\begin{aligned} \sum_{j=1}^m \tau(R_j) &= \sum_{j=1}^m \left\{ \lambda_n(R_{nj}) \cdot \prod_{i=n+1}^\infty (b_i - a_i) \right\} \\ &\leq \left\{ \prod_{i=1}^n (b_i - a_i) + \varepsilon \right\} \cdot \prod_{i=n+1}^\infty (b_i - a_i) \\ &= \tau(R) + \varepsilon \prod_{i=n+1}^\infty (b_i - a_i) \\ &< \tau(R) + \varepsilon(1 + \varepsilon). \end{aligned}$$

Because ε is arbitrary, we get $\nu_\delta(R) \leq \tau(R)$. But δ is also arbitrary, hence $\nu(R) \leq \tau(R)$. Therefore $(**)$ holds. $\square$

Theorem 3.5. We have $\lambda(E) = \nu(E)$, for all $E \subseteq \mathbf{R}^\infty$. Hence, by Theorem 2.8, λ is a metric outer measure on $\mathbf{R}^\infty$.

Proof. For $E \subseteq \mathbf{R}^\infty$, we have $\lambda(E) \leq \nu(E)$, hence it suffices to prove that

$$(\dagger) \quad \nu(E) \leq \lambda(E), \quad E \subseteq \mathbf{R}^\infty.$$

Fix $E \subseteq \mathbf{R}^\infty$. By Theorem 2.3, we have

$$\nu(E) = \inf_{\substack{C_j \subseteq \mathbf{R}^\infty \\ \bigcup C_j \supseteq E}} \sum_{j=1}^\infty \nu(C_j).$$

Hence we see that

$$\nu(E) \leq \inf_{\substack{R_j \in \mathcal{R} \\ \bigcup R_j \supseteq E}} \sum_{j=1}^\infty \nu(R_j).$$

For all $R \in \mathcal{R}$, Theorem 3.5 implies that $\nu(R) = \tau(R)$, therefore we have $\nu(E) \leq \lambda(E)$. This proves (†). $\square$

Theorem 3.6. *The outer measure λ is translation invariant on $\mathbf{R}^\infty$.*

Proof. This follows from the fact that if $R = \prod_{i=1}^\infty (a_i, b_i) \in \mathcal{R}$ and $x \in \mathbf{R}^\infty$, then $R + x \in \mathcal{R}$ and $\tau(R + x) = \tau(R)$. $\square$

We conclude with Theorem I:

Theorem 3.7. *The outer measure τ has all the properties stated in Theorem I.*

Proof. This is the content of Theorems 2.7, 2.8, 3.3, 3.5, and 3.6. $\square$

Remarks. We conclude the paper with the following observations. These observations are presented without proof. (a) The measure λ is *Borel regular*, i.e., if $E \subseteq \mathbf{R}^\infty$, then there is a Borel subset $R \subseteq \mathbf{R}^\infty$ such that

$$E \subseteq R \quad \text{and} \quad \lambda(E) = \lambda(R).$$

For a proof of this, see [1, p. 34].

(b) The following change of variable formula holds: Let $T^n: \mathbf{R}^n \rightarrow \mathbf{R}^n$, $n \geq 1$, be a linear transformation with Jacobian $\Delta \neq 0$, and let $T^\infty: \mathbf{R}^\infty \rightarrow \mathbf{R}^\infty$ be the map defined by

$$T^\infty(x) = (T^n(x_1, \dots, x_n), x_{n+1}, x_{n+2}, \dots), \quad x = (x_i) \in \mathbf{R}^\infty.$$

Then for each $E \subseteq \mathbf{R}^\infty$, $\lambda(T^\infty(E)) = |\Delta|\lambda(E)$. The proof of this theorem is based on the same type of arguments used in the proof of Theorem 3.4.

REFERENCES

1. C. A. Rogers, *Hausdorff measures*, Cambridge Univ. Press, 1970.
2. Y. Yamasaki, *Measures on infinite dimensional spaces*, World Scientific, Singapore, 1985.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY, CALIFORNIA 94720