THE FEKETE-SZEGŐ PROBLEM FOR STRONGLY CLOSE-TO-CONVEX FUNCTIONS

H. R. ABDEL-GAWAD AND D. K. THOMAS

(Communicated by Clifford J. Earle, Jr.)

ABSTRACT. Let $K(\beta)$ denote the class of normalized analytic strongly close-to-convex functions of order $\beta \geq 0$, defined in the unit disc D and let $f \in K(\beta)$, with $f(z) = z + a_2 z^2 + a_3 z^3 + \cdots$, for $z \in D$. Sharp bounds are obtained for $|a_3 - \mu a_2^2|$ when μ is real.

INTRODUCTION

Denote by S the class of normalized analytic univalent functions f defined for $z \in D = \{z : |z| < 1\}$ by

$$(1) \quad f(z) = z + \sum_{n=2}^{\infty} a_n z^n.$$

A classical theorem of Fekete and Szegő [2] states that for $f \in S$ given by (1),

$$|a_3 - \mu a_2^2| \leq \begin{cases} 3 - 4\mu, & \text{if } \mu \leq 0 \\ 1 + 2e^{-2\mu/(1-\mu)}, & \text{if } 0 \leq \mu < 1 \\ 4\mu - 3, & \text{if } \mu \geq 1. \end{cases}$$

This inequality is sharp in the sense that for each μ there exists a function in S such that equality holds. Recently Pfluger [8] has considered the problem when μ is complex. In the case of C , S^* and K , the subclasses of convex, starlike and close-to-convex functions respectively, the above inequalities can be improved [5, 6]. In particular for $f \in K$ and given by (1), Keogh and Merkes [5] showed that

$$|a_3 - \mu a_2^2| \leq \begin{cases} 3 - 4\mu, & \text{if } \mu \leq 1/3, \\ 1/3 + 4/9\mu, & \text{if } 1/3 \leq \mu \leq 2/3, \\ 1, & \text{if } 2/3 \leq \mu \leq 1, \\ 4\mu - 3, & \text{if } \mu \geq 1. \end{cases}$$

Again, for each μ , there is a function in K such that equality holds. In this paper we extend this result to the class $K(\beta)$ of strongly close-to-convex functions of order β in the sense of Pommerenke [9]. Thus $f \in K(\beta)$ if, and

Received by the editors March 9, 1989.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 30C45.

© 1992 American Mathematical Society
0002-9939/92 \$1.00 + \$.25 per page

only if, f , given by (1), is analytic in D and is such that there exists $g \in S^*$ satisfying

$$(2) \quad \left| \arg \frac{zf'(z)}{g(z)} \right| \leq \frac{\pi\beta}{2},$$

for $z \in D$ and $\beta \geq 0$. Clearly $K(0) = C$, $K(1) = K$ and when $0 \leq \beta \leq 1$, $K(\beta)$ is a subset of K and hence contains only univalent functions. However in [4], Goodman showed that $K(\beta)$ can contain functions with unbounded valence for $\beta > 1$.

Recently, Koepf [7] has considered the Fekete-Szegő problem for $K(\beta)$ and obtained sharp results for some particular values of μ , all of which, with the exception of the case $\mu = 1$ and $\beta \geq 1$, are contained in the following result.

RESULTS

Theorem. Let $f \in K(\beta)$ and be given by (1). Then for $0 \leq \beta \leq 1$,

$$|a_3 - \mu a_2^2| \leq \begin{cases} 1 - \mu + \frac{\beta(2 - 3\mu)(\beta + 2)}{3}, & \text{if } \mu \leq \frac{2\beta}{3(\beta + 1)}, \\ 1 - \mu + \frac{2\beta}{3} + \frac{\beta(2 - 3\mu)^2}{3[2 - \beta(2 - 3\mu)]}, & \text{if } \frac{2\beta}{3(\beta + 1)} \leq \mu \leq \frac{2}{3}, \\ \frac{2\beta + 1}{3}, & \text{if } \frac{2}{3} \leq \mu \leq \frac{2(\beta + 2)}{3(\beta + 1)}, \\ \mu - 1 + \frac{\beta(3\mu - 2)(\beta + 2)}{3}, & \text{if } \mu \geq \frac{2(\beta + 2)}{3(\beta + 1)}, \end{cases}$$

whilst for $\beta > 1$, the first two inequalities hold. For each μ there are functions in $K(\beta)$ such that equality holds in all cases.

We shall require the following:

Lemma 1 ([10, p. 166]). Let $h \in P$, i.e., let h be analytic in D and satisfy $\operatorname{Re} h(z) > 0$ for $z \in D$, with $h(z) = 1 + c_1 z + c_2 z^2 + \dots$, then

$$\left| c_2 - \frac{c_1^2}{2} \right| \leq 2 - \frac{|c_1|^2}{2}.$$

Lemma 2 ([6, Lemma 3]). Let $g \in S^*$ with $g(z) = z + b_2 z^2 + b_3 z^3 + \dots$, then for μ real,

$$|b_3 - \mu b_2^2| \leq \max\{1, |3 - 4\mu|\}.$$

We note that Lemma 2 above can easily be extended to the wider class $S^*(\alpha)$ of strongly starlike functions of order $\alpha \geq 0$, i.e., g analytic and normalized in D and satisfying

$$\left| \arg \frac{zg'(z)}{g(z)} \right| \leq \frac{\alpha\pi}{2},$$

see e.g. [1]. In this case, one obtains the sharp inequality

$$|b_3 - \mu b_2^2| \leq \max\{\alpha, \alpha^2|3 - 4\mu|\},$$

for μ real.

Proof of Theorem. It follows from (2) that we can write

$$(3) \quad zf'(z) = g(z)h(z)^\beta$$

for $g \in S^*$ and $h \in P$. Equating coefficients in (3) we obtain $2a_2 = \beta c_1 + b_2$ and $3a_3 = (\beta(\beta - 1)/2)c_1^2 + \beta c_2 + \beta c_1 b_2 + b_3$, so that

$$(4) \quad \begin{aligned} a_3 - \mu a_2^2 &= \frac{1}{3} \left(b_3 - \frac{3}{4} \mu b_2^2 \right) + \frac{\beta}{3} \left(c_2 + \left(\frac{\beta(2-3\mu)}{4} - \frac{1}{2} \right) c_1^2 \right) \\ &\quad + \beta \left(\frac{1}{3} - \frac{\mu}{2} \right) c_1 b_2. \end{aligned}$$

We consider first the case $(2\beta)/(3(\beta + 1)) \leq \mu \leq 2/3$. Equation (4) gives

$$\begin{aligned} |a_3 - \mu a_2^2| &\leq \frac{1}{3} \left| b_3 - \frac{3}{4} \mu b_2^2 \right| + \frac{\beta}{3} \left| c_2 - \frac{1}{2} c_1^2 \right| + \frac{\beta^2(2-3\mu)}{12} |c_1^2| + \beta \left(\frac{1}{3} - \frac{\mu}{2} \right) |c_1| |b_2|, \\ &\leq 1 - \mu + \frac{\beta}{3} \left(2 - \frac{1}{2} |c_1^2| \right) + \frac{\beta^2(2-3\mu)}{12} |c_1^2| + \frac{\beta(2-3\mu)}{3} |c_1|, \\ &= \Phi(x) \text{ say, with } x = |c_1|, \end{aligned}$$

where we have used Lemmas 1 and 2 and the fact that $|b_2| \leq 2$ for $g \in S^*$. An elementary argument shows that the function Φ attains a maximum at $x_0 = 2(2-3\mu)/(2-\beta(2-3\mu))$, and so

$$|a_3 - \mu a_2^2| \leq \Phi(x_0),$$

which proves the theorem if $\mu \leq 2/3$ and $\beta \geq 0$. Choosing

$$c_1 = \frac{2(2-3\mu)}{2-\beta(2-3\mu)}, \quad c_2 = 2, \quad b_2 = 2, \quad \text{and} \quad b_3 = 3,$$

in (4) shows that the result is sharp. We note that $|c_1| \leq 2$, that is, $\mu \geq 2\beta/(3(\beta + 1))$.

Next consider the case $\mu \leq (2\beta)/(3(\beta + 1))$. Then

$$\begin{aligned} |a_3 - \mu a_2^2| &\leq \left| a_3 - \frac{2\beta}{3(\beta+1)} a_2^2 \right| + \left(\frac{2\beta}{3(\beta+1)} - \mu \right) |a_2|^2, \\ &\leq 1 + \frac{2\beta}{3} + \left(\frac{2\beta}{3(\beta+1)} - \mu \right) (\beta+1)^2 = 1 - \mu + \frac{\beta(2-3\mu)(\beta+2)}{3}, \end{aligned}$$

for $\beta \geq 0$, where we have used the result already proved in the case $\mu = 2\beta/(3(\beta + 1))$, and the fact that for $f \in K(\beta)$, the inequality $|a_2| \leq \beta + 1$ holds [3]. Equality is attained on choosing $\lambda = 0$, $p_1 = p_2 = b_2 = 2$, and $b_3 = 3$.

Suppose now that $2/3 \leq \mu \leq (2(\beta + 2))/(3(\beta + 1))$. Since $g \in S^*$ we can write $zg'(z) = g(z)p(z)$ for $p \in P$, with $p(z) = 1 + p_1 z + p_2 z^2 + \dots$, and so equating coefficients we have that $b_2 = p_1$ and $2b_3 = p_1^2 + p_2$.

We deal first with the case $\mu = 2(\beta + 2)/(3(\beta + 1))$. Thus (4) gives

$$\begin{aligned} a_3 - \frac{2(\beta+2)}{3(\beta+1)} a_2^2 &= \frac{1}{6} \left(p_2 - \frac{p_1^2}{2} \right) + \frac{\beta}{3} \left(c_2 - \frac{c_1^2}{2} \right) + \frac{\beta-1}{12(\beta+1)} p_1^2 \\ &\quad - \frac{\beta^2 c_1^2}{6(\beta+1)} - \frac{\beta p_1 c_1}{3(\beta+1)}, \end{aligned}$$

and so if $\beta \leq 1$,

$$\begin{aligned} \left| a_3 - \frac{2(\beta+2)}{3(\beta+1)} a_2^2 \right| &\leq \frac{1}{6} \left| p_2 - \frac{p_1^2}{2} \right| + \frac{\beta}{3} \left| c_2 - \frac{c_1^2}{2} \right| + \frac{(1-\beta)}{12(\beta+1)} |p_1^2| \\ &\quad + \frac{\beta^2 |c_1^2|}{6(\beta+1)} + \frac{\beta |p_1 c_1|}{3(\beta+1)}, \\ &\leq \frac{1}{6} \left(2 - \frac{|p_1^2|}{2} \right) + \frac{\beta}{3} \left(2 - \frac{|c_1^2|}{2} \right) + \frac{1-\beta}{12(\beta+1)} |p_1^2| \\ &\quad + \frac{\beta^2 |c_1^2|}{6(\beta+1)} + \frac{\beta |p_1 c_1|}{3(\beta+1)}, \\ &= \frac{2\beta+1}{3} - \frac{\beta}{6(\beta+1)} (|c_1| - |p_1|)^2, \\ &\leq \frac{2\beta+1}{3}, \end{aligned}$$

where we have used Lemma 1.

Now write

$$\begin{aligned} a_3 - \mu a_2^2 &= \frac{(\beta+1)(3\mu-2)}{2} \left(a_3 - \frac{2(\beta+2)}{3(\beta+1)} a_2^2 \right) \\ &\quad + \frac{3(\beta+1)}{2} \left(\frac{2(\beta+2)}{3(\beta+1)} - \mu \right) \left(a_3 - \frac{2}{3} a_2^2 \right), \end{aligned}$$

and the result follows at once on using the theorem already proved in the cases $\mu = 2/3$ and $\mu = 2(\beta+2)/(3(\beta+1))$ for $\beta \leq 1$. Equality is attained when f is given by

$$f'(z) = \frac{(1+z^2)^\beta}{(1-z^2)^{\beta+1}}.$$

We finally assume that $\mu \geq (2(\beta+2))/(3(\beta+1))$. Write

$$a_3 - \mu a_2^2 = \left(a_3 - \frac{2(\beta+2)}{3(\beta+1)} a_2^2 \right) + \left(\frac{2(\beta+2)}{3(\beta+1)} - \mu \right) a_2^2,$$

and the result follows at once on using the theorem already proved for $\mu = 2(\beta+2)/(3(\beta+1))$ in the case $\beta \leq 1$ and the inequality $|a_2| \leq \beta+1$, which was proved in [3]. Equality is attained in this last case on choosing $c_1 = b_2 = 2i$, $c_2 = -2$ and $b_3 = -3$ in (4).

We remark that the methods used in [5, 6], together with equation (4), suggest that in order to obtain sharp results for $\beta > 1$ and $\mu > 2/3$, an extension to the "area principle" may be required. Since $K(\beta)$ contains functions of unbounded valence for $\beta > 1$ establishing sharp estimates in this case may require deeper methods.

REFERENCES

1. D. A. Brannan and W. E. Kirwan, *On some classes of bounded univalent functions*, J. London Math. Soc. (2) **1** (1969), 431-443.
2. M. Fekete and G. Szegő, *Eine Bemerkung über ungerade schlichte Funktionen*, J. London Math. Soc. **8** (1933), 85-89.
3. A. W. Goodman, *On close-to-convex functions of higher order*, Ann. Univ. Sci. Budapest. Eötvös Sect. Math. **15** (1972), 17-30.

4. ———, *A note on the Noshiro-Warschawski theorem*, J. Analyse Math. **25** (1972), 401–408.
5. F. R. Keogh and E. P. Merkes, *A coefficient inequality for certain classes of analytic functions*, Proc. Amer. Math. Soc. **20** (1969), 8–12.
6. W. Koepf, *On the Fekete-Szegő problem for close-to-convex functions*, Proc. Amer. Math. Soc. **101** (1987), 89–95.
7. ———, *On the Fekete-Szegő problem for close-to-convex functions. II*, Arch. Math. **49** (1987), 420–433.
8. A. Pfluger, *The Fekete-Szegő inequality for complex parameters*, Complex Variables Theory Appl. **7** (1986), 149–160.
9. Ch. Pommerenke, *On close-to-convex analytic functions*, Trans. Amer. Math. Soc. **114** (1965), 176–186.
10. ———, *Univalent functions*, Vandenhoeck and Ruprecht, Göttingen, 1975.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, UNIVERSITY COLLEGE SWANSEA,
SWANSEA SA2 8PP, WALES