INTERPOLATING OPERATORS IN NEST ALGEBRAS

M. ANOUSSIS

(Communicated by Paul S. Muhly)

ABSTRACT. Given two families $\{x_\alpha\}, \{y_\alpha\}$ of vectors and a nest $\mathcal{N}$ in a Hilbert space H , we provide a necessary and sufficient condition for the existence of an operator T in the nest algebra satisfying $Tx_\alpha = y_\alpha$ for every α .

Given two families of vectors $\{x_\alpha\}, \{y_\alpha\}$ in a Hilbert space, an interpolating operator is a bounded operator T such that $Tx_\alpha = y_\alpha$ for every α . In [5] Lance gives a necessary and sufficient condition on two vectors x and y for the existence of an operator T lying in a given nest algebra such that $Tx = y$. Hopenwasser extended this result to the case of a *CSL* algebra [3]. Other questions about interpolating operators are investigated in [1, 4]. In this note we give a necessary and sufficient condition on two families $\{x_\alpha\}, \{y_\alpha\}$ of vectors in a Hilbert space for the existence of an operator T lying in a given nest algebra, such that $Tx_\alpha = y_\alpha$ for every α .

Throughout this note H denotes a complex Hilbert space and $B(H)$ denotes the space of all bounded operators from H to itself. By a nest we mean a complete totally ordered set of orthogonal projections. If $\mathcal{N}$ is a nest in H , then $\text{Alg } \mathcal{N}$ is the set of operators X in $B(H)$ satisfying $N^\perp X N = 0$ for every N in $\mathcal{N}$.

The main step in our proof is Proposition 3. We need two lemmas.

Lemma 1. *Let $x_1, x_2, \dots, x_m, y_1, y_2, \dots, y_m$ be vectors in H . Let V be the span of the x_i 's, $i = 1, 2, \dots, m$. Assume that there exists a real number $M > 0$ such that:*

$$\left\| \sum_{i=1}^m c_i y_i \right\| \leq M \cdot \left\| \sum_{i=1}^m c_i x_i \right\|,$$

for every m -tuple $(c_1, c_2, \dots, c_m)$ of complex numbers. Then there exists a linear operator $T: V \rightarrow H$ such that $Tx_i = y_i$, for $i = 1, 2, \dots, m$ and $\|T\| \leq M$.

Proof. We may assume that there exists an index k , $1 \leq k \leq m$ such that the set $\{x_1, x_2, \dots, x_k\}$ is a basis of V . Define $T: V \rightarrow H$ such that $Tx_i = y_i$, $1 \leq i \leq k$. Then T is a linear operator from V to H , and by the condition of the lemma, it follows that $\|T\| \leq M$. We have to show that $Tx_i = y_i$ for every

Received by the editors September 4, 1990.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 47D25.

i , $1 \leq i \leq m$. Suppose, to the contrary, there exists l , $k < l \leq m$ such that $Tx_l \neq y_l$. There are complex numbers a_i , $1 \leq i \leq k$ such that $x_l = \sum_{i=1}^k a_i x_i$. Then,

$$\left\| \sum_{i=1}^k a_i y_i - y_l \right\| = \left\| \sum_{i=1}^k a_i T x_i - y_l \right\| = \left\| T \left(\sum_{i=1}^k a_i x_i \right) - y_l \right\| = \|Tx_l - y_l\| \neq 0,$$

while $\left\| \sum_{i=1}^k a_i x_i - x_l \right\| = 0$, contradicting the hypothesis. $\square$

Lemma 2. Let A be a positive contraction in $B(H)$. We denote by B the positive semidefinite hermitian form on H defined by $B(x, y) = \langle x, Ay \rangle$ for x, y in H . Let V be a finite-dimensional subspace of H . Then there exists a closed subspace V_0 of H such that:

- (1) $V + V_0 = H$, $V \cap V_0 = \{0\}$;
- (2) $B(x, y) = 0$ for every x in V and y in V_0 .

Proof. Let $V_1 = \{x \in H : B(x, y) = 0, \forall y \in V\} = (AV)^\perp$. We show that $V + V_1 = H$. Let z be a vector in $(V + V_1)^\perp = V^\perp \cap V_1^\perp$. Since $V_1 = (AV)^\perp$, which has finite codimension, z is of the form Av for some v in V . We have:

$$\langle v, Av \rangle = 0 \Rightarrow A^{1/2}v = 0 \Rightarrow Av = 0 \Rightarrow z = 0.$$

Let $V_0 = V_1 \ominus (V \cap V_1)$. We have:

$$V + V_0 = V + V_1 = H, \quad V \cap V_0 = (V \cap V_1) \cap V_0 = \{0\}.$$

It follows that the first condition of the lemma is satisfied. The second condition holds because V_0 is contained in V_1 . $\square$

Proposition 3. Let $\mathcal{N} = \{0 = N_0, N_1, N_2, \dots, N_k = I\}$ be a finite nest in H . Let $x_1, x_2, \dots, x_m, y_1, y_2, \dots, y_m$ be vectors in H . Let $M > 0$. The following are equivalent:

- (1) There exists an operator T in $\text{Alg } \mathcal{N}$ such that $Tx_i = y_i$, for $i = 1, 2, \dots, m$, and $\|T\| \leq M$.
- (2) $\|N^\perp(\sum_{i=1}^m c_i y_i)\| \leq M \cdot \|N^\perp(\sum_{i=1}^m c_i x_i)\|$, for every m -tuple of complex numbers $(c_1, c_2, \dots, c_m)$, and every projection N in $\mathcal{N}$.

Proof. The implication (1) $\Rightarrow$ (2) is easy:

$$\begin{aligned} \left\| N^\perp \left(\sum_{i=1}^m a_i y_i \right) \right\| &= \left\| N^\perp T \left(\sum_{i=1}^m a_i x_i \right) \right\| = \left\| N^\perp T N^\perp \left(\sum_{i=1}^m a_i x_i \right) \right\| \\ &\leq \|N^\perp T\| \cdot \left\| N^\perp \left(\sum_{i=1}^m a_i x_i \right) \right\| \leq M \cdot \left\| N^\perp \left(\sum_{i=1}^m a_i x_i \right) \right\|. \end{aligned}$$

The converse is shown by induction on the length k of the nest $\mathcal{N}$. We may assume that $M = 1$.

First suppose that $k = 1$. It follows from Lemma 1 that there exists a contraction T_0 from the span V of the x_i 's, $i = 1, 2, \dots, m$ into H such that $T_0 x_i = y_i$, $i = 1, 2, \dots, m$. We may extend T_0 to an operator T on H by defining T to be 0 on $V^\perp$. Now assume that the length of $\mathcal{N}$ is $k > 1$. Let H_0 be the Hilbert space $N_1^\perp H$, and let $\mathcal{N}_0$ be the nest in H_0 consisting

of all projections of the form $N_1^\perp N$ with N in $\mathcal{N}$. It is easy to see that if L belongs to $\mathcal{N}_0$ then:

$$\left\| L^\perp \left(\sum_{i=1}^m c_i N_1^\perp y_i \right) \right\| \leq M \cdot \left\| L^\perp \left(\sum_{i=1}^m c_i N_1^\perp x_i \right) \right\|,$$

for every m -tuple $(c_1, c_2, \dots, c_m)$ of complex numbers. Hence, by the induction hypothesis, there exists a contraction A_0 in $\text{Alg } \mathcal{N}_0$ such that $A_0 N_1^\perp x_i = N_1^\perp y_i$ for every i , $1 \leq i \leq m$. We denote by T_0 the operator defined by $T_0 x = A_0 x$ if $x \in N_1^\perp H$, $T_0 x = 0$ if $x \in N_1 H$. It is easy to see that T_0 belongs to $\text{Alg } \mathcal{N}$. For each $(c_1, c_2, \dots, c_m)$ in $\mathbb{C}^m$ we have:

$$\left\| N_1 \left(\sum_{i=1}^m c_i y_i \right) \right\| \leq \left\| \sum_{i=1}^m c_i y_i \right\| \leq 1 \cdot \left\| \sum_{i=1}^m c_i x_i \right\|.$$

Thus by Lemma 1, there exists a contraction $A_1: V \rightarrow H$ such that $A_1 x_i = N_1 y_i$ for every i , $1 \leq i \leq m$. For each $(c_1, c_2, \dots, c_m)$ in $\mathbb{C}^m$, we have:

$$\begin{aligned} & \left\| N_1 \left(\sum_{i=1}^m c_i y_i \right) \right\|^2 + \left\| N_1^\perp \left(\sum_{i=1}^m c_i y_i \right) \right\|^2 \leq \left\| \sum_{i=1}^m c_i x_i \right\|^2 \\ & \Rightarrow \left\| A_1 \left(\sum_{i=1}^m c_i x_i \right) \right\|^2 \leq \left\| \sum_{i=1}^m c_i x_i \right\|^2 - \left\| N_1^\perp \left(\sum_{i=1}^m c_i y_i \right) \right\|^2 \\ & = \left\| \sum_{i=1}^m c_i x_i \right\|^2 - \left\| N_1^\perp T_0 N_1^\perp \left(\sum_{i=1}^m c_i x_i \right) \right\|^2, \end{aligned}$$

since $T_0 N_1^\perp x_i = N_1^\perp y_i$ for $1 \leq i \leq m$ by construction. It follows that for every v in V , we have:

$$\|A_1 v\|^2 \leq \|v\|^2 - \|N_1 T_0 N_1^\perp v\|^2.$$

Rewrite this as:

$$(*) \quad \|A_1 v\|^2 \leq B(v, v), \quad \text{where } B(x, y) = \langle x, (I - N_1^\perp T_0^* N_1^\perp T_0 N_1^\perp) y \rangle$$

for x, y in H . Since $\|T_0\| \leq 1$, $I - N_1^\perp T_0^* N_1^\perp T_0 N_1^\perp$ is a positive contraction. By Lemma 2, there exist a closed subspace V_0 of H such that $V + V_0 = H$, $V \cap V_0 = \{0\}$, and $B(x, y) = 0$ for every x in V and y in V_0 . We define an operator $T_1: H \rightarrow H$ by: $T_1 x = A_1 x$ if $x \in V$, $T_1 x = 0$ if $x \in V_0$.

Claim. $\|T_1 x\|^2 \leq B(x, x)$, for every x in H .

Indeed, let x be in H . Then x is of the form $x = v + v_0$ with $v \in V$ and $v_0 \in V_0$. We have:

$$\begin{aligned} \|T_1 x\|^2 &= \|T_1(v + v_0)\|^2 = \|A_1 v\|^2 \leq B(v, v), \quad \text{by } (*) \\ &\leq B(v, v) + B(v_0, v_0) = B(v + v_0, v + v_0) = B(x, x). \end{aligned}$$

The range of T_1 being contained in $N_1 H$, T_1 belongs to $\text{Alg } \mathcal{N}$. We may now define $T = T_1 + T_0$. Then $T \in \text{Alg } \mathcal{N}$ and $T x_i = y_i$ for every i , $1 \leq i \leq m$. It remains to see that T is a contraction. Indeed, for every x in H , we have:

$$\begin{aligned} \|T x\|^2 &= \|T_1 x + T_0 x\|^2 = \|T_1 x\|^2 + \|T_0 x\|^2 \\ &\leq B(x, x) + \|T_0 x\|^2 = \|x\|^2 - \|N_1^\perp T_0 N_1^\perp x\|^2 + \|T_0 x\|^2 = \|x\|^2, \end{aligned}$$

where we have used the fact that $T_0 = N_1^\perp T_0 N_1^\perp$. $\square$

Theorem 4. Let $\mathcal{N}$ be a nest in H . Let $\{x_\alpha\}, \{y_\alpha\}$ be two families of vectors in H . Let $M \geq 0$. The following are equivalent:

- (1) There exist T in $\text{Alg } \mathcal{N}$ such that $Tx_\alpha = y_\alpha$ for every α and $\|T\| \leq M$.
- (2) $\|N^\perp(\sum c_\alpha y_\alpha)\| \leq M \cdot \|N^\perp(\sum c_\alpha x_\alpha)\|$, for every N in $\mathcal{N}$ and for every family $\{c_\alpha\}$ of complex numbers, that is 0 except for finitely many indices.

Proof. The implication (1) $\Rightarrow$ (2) is as easy as in the finite nest case.

For the converse, first assume that the families $\{x_\alpha\}, \{y_\alpha\}$ are finite.

Let $\mathcal{L}$ be the set of finite subnests of $\mathcal{N}$ directed by inclusion. It follows from the proposition that for every $\mathcal{F}$ in $\mathcal{L}$ there exists an operator $T_{\mathcal{F}}$ in $\text{Alg } \mathcal{F}$ such that $T_{\mathcal{F}}x_\alpha = y_\alpha$ for every α and $\|T_{\mathcal{F}}\| \leq M$. The net $\{T_{\mathcal{F}} : \mathcal{F} \in \mathcal{L}\}$ is contained in the set $\{T \in B(H) : \|T\| \leq M\}$, which is compact in the weak operator topology. Let T be a weak operator limit of the net $\{T_{\mathcal{F}} : \mathcal{F} \in \mathcal{L}\}$. Clearly, $\|T\| \leq M$ and $Tx_\alpha = y_\alpha$ for every α . Also it is easy to see that T leaves every N in $\mathcal{N}$ invariant.

For the general case, we may assume that the families $\{x_\alpha\}, \{y_\alpha\}$ are indexed by an infinite set A . Then it follows from above that for every finite subset U of A there exists an operator T_U in $\text{Alg } \mathcal{N}$ such that $T_U x_\alpha = y_\alpha$ for every α in U and $\|T_U\| \leq M$. Let T be a weak operator limit point of the net $\{T_U\}$. Then $\|T\| \leq M$, $Tx_\alpha = y_\alpha$ for every α in A and T belongs to $\text{Alg } \mathcal{N}$ since $\text{Alg } \mathcal{N}$ is closed in the weak operator topology. $\square$

The following corollary generalizes Theorem 4.3 of [2].

Corollary 5. Let A and B be operators in $B(H)$, and let $\mathcal{N}$ be a nest in H . Let $M \geq 0$. The following are equivalent:

- (1) There exists an operator X in $\text{Alg } \mathcal{N}$ such that $A = XB$ and $\|X\| \leq M$.
- (2) $\|N^\perp Ax\| \leq M \cdot \|N^\perp Bx\|$, for every N in $\mathcal{N}$ and x in H .

Proof. Let $\{e_\alpha\}$ be an orthonormal basis of H . We set $y_\alpha = Ae_\alpha$, $x_\alpha = Be_\alpha$ and apply the Theorem. $\square$

Remark. After this work was completed, it came to our attention (private communication with E. Katsoulis) that E. Katsoulis, R. Moore, and T. Trent have arrived at similar results.

REFERENCES

1. M. Anoussis, E. Katsoulis, R. Moore, and T. Trent, *Interpolation problems for ideals in certain reflexive operator algebras*, preprint.
2. W. Arveson, *Interpolation problems in nest algebras*, J. Funct. Anal. **20** (1975), 208–233.
3. A. Hopenwasser, *The equation $Tx = y$ in a reflexive operator algebra*, Indiana Univ. Math. J. **29** (1980), 121–126.
4. —, *Hilbert-Schmidt interpolation in CSL algebras*, Illinois J. Math. **33** (1989), 657–672.
5. E. Lance, *Some properties of nest algebras*, Proc. London Math. Soc. (3) **19** (1969), 45–68.