LAYERS OF COMPONENTS OF $\beta([0, 1] \times N)$ ARE INDECOMPOSABLE

MICHEL SMITH

(Communicated by James E. West)

ABSTRACT. We examine the structure of certain subcontinua of the Stone-Čech compactification of the reals. Let N denote the integers, let $X = [0, 1] \times N$, and let C be a component of $X^* = \beta X - X$. It is known that C admits an upper semicontinuous decomposition G into maximal nowhere dense subcontinua of C so that C/G is a Hausdorff arc. The elements of G are called layers. It has been shown that the layers of C that contain limit points of a countable increasing or decreasing sequence of cut points of C are nondegenerate indecomposable continua (various forms of this fact have been proven by Bellamy and Rubin, Mioduszewski, and Smith). We show that all the layers of C are indecomposable. *

Let $A = [0, \infty)$. We wish to examine certain subcontinua of $A^* = \beta A - A$ the remainder of the Stone-Čech compactification of A . Let N denote the integers. Then $X = [0, 1] \times N$ is a subspace of A and X^* is a subspace of A^* . It has been shown that certain layers of components of X^* are indecomposable. (See Mioduszewski [M], Bellamy and Rubin [BR], and Smith [S4].) It is the purpose of this paper to show that every "layer" of components of X^* is indecomposable. This answers a question posed by Mioduszewski in [M]. A layer of a component C of X^* is a maximal nowhere dense (in C) subcontinuum of C .

First we introduce some notation. Let $Z(X)$ denote the zero sets of X (the closed subsets of X). Elements of βX are identified with ultrafilters of zero sets of X (closed sets of X). Let I_n denote $[0, 1] \times \{n\}$ for each $n \in N$. If $x \in X^*$, let $u_x = \{\{n | I_n \cap H \neq \emptyset\} | H \in x\}$. It can be verified that if $x \in X^*$ then $u_x \in N^*$ and if $u \in N^*$ then $C(u) = \{x | u_x = u\}$ is a component of X^* .

We wish to state some properties of the layers of components of X^* and a useful characterization.

There is a natural order induced by the topology on the layers of $C(u)$. Let $p, q \in C(u)$; then we say that $p <_u q$ if and only if there exist $H \in p$, $K \in q$, and $D \in u$ so that every element of $H \cap I_n$ precedes every element of $K \cap I_n$ with respect to the usual order on I_n for all $n \in D$ (i.e., $H \cap I_n <_n K \cap I_n \ \forall n \in D$).

Received by the editors August 15, 1988 and, in revised form, October 19, 1990.

1991 *Mathematics Subject Classification*. Primary 54D40, 54F20.

Key words and phrases. Stone-Čech compactifications, nonmetric continua, indecomposable continua.

Presented at the Spring Topology Conference, Sacramento, CA, April 11–13, 1991.

* The theorem was recently discovered independently by J. Zhu.

The subscript “ u ” of the symbol “ $<_u$ ” may be omitted if the meaning is clear from the context. Let Ω denote the set of all sequences $S = \{S(n)\}_{n=1}^{\infty}$ so that $S(n) \in I_n$. If $u \in N^*$ and $S \in \Omega$ then define $A(u, S) = \{H \in Z(\bar{X}) \mid \exists D \in u \text{ so that } \{S(n) \mid n \in D\} \subset H\}$. $A(u, S)$ is an ultrafilter and hence an element of X^* . In fact $A(u, S) \in C(u)$. If $\{i \mid S(i) = 1\} \in u$ then $A(u, S)$ is an end point of $C(u)$. Similarly if $\{i \mid S(i) = 0\} \in u$ then $A(u, S)$ is an end point of $C(u)$. Except for these two cases, if $S \in \Omega$ then $A(u, S)$ is a cut point of $C(u)$. (The converse cannot be assumed. See Baldwin and Smith [BS].) Note that $\{A(u, S) \mid S \in \Omega\}$ is dense in $C(u)$.

The following observations are not difficult to verify (see Baldwin and Smith [BS], Mioduszewski [M], and Smith [S4]).

Observation 1. If p and q are two points of $C(u)$ and $p >_u q$ then there is a cut point $A(u, S)$ with $S \in \Omega$ so that $p <_u A(u, S) <_u q$.

Observation 2. If $p = A(u, S)$ for some $S \in \Omega$ and $q \in C(u)$, with $q \neq p$, then either $q <_u p$ or $p <_u q$.

Observation 3. If S_1 and S_2 are two elements of Ω , $u \in N^*$, and $A(u, S_1) <_u A(u, S_2)$ then $\{p \in C(u) \mid A(u, S_1) <_u p <_u A(u, S_2)\}$ is open in $C(u)$ and $\{p \in C(u) \mid A(u, S_1) \leq_u p \leq_u A(u, S_2)\}$ is a subcontinuum of $C(u)$, and hence is not nowhere dense.

Lemma 1. Every cut point of $C(u)$ is a layer of $C(u)$.

Proof. If $x \in C(u)$ then define L_x and R_x as follows:

$$L_x = \{p \in C(u) \mid p <_u x\} \quad R_x = \{p \in C(u) \mid x <_u p\}.$$

If p and q are points of $C(u)$ and $p <_u q$ then there is a sequence $S \in \Omega$ so that $p <_u A(u, S) <_u q$. Therefore we have

Claim 1.1. If $x \in C(u)$ then L_x and R_x are connected.

Also since $\{A(u, S) \mid S \in \Omega\}$ is dense in $C(u)$ we have

Claim 1.2. $\text{Cl}_{C(u)}(R_x \cup L_x) = C(u)$.

We also need to prove

Claim 1.3. If $x \in M$ and M is a nowhere dense subcontinuum of $C(u)$ containing x then $M \cap L_x = \emptyset$ and $M \cap R_x = \emptyset$.

Proof. Suppose $x \in C(u)$ and M is a subcontinuum of $C(u)$ that contains x and $M \cap R_x \neq \emptyset$. Then there exists $y \in M \cap R_x$ so that $x <_u y$. By Observation 1 there exist S_1 and S_2 in Ω so that $x <_u A(u, S_1) <_u A(u, S_2) <_u y$. Since $A(u, S_1)$ is a cut point of $C(u)$ and separates x from y , it follows that $A(u, S_1) \in M$. Similarly $A(u, S_2) \in M$; and in fact, if $A(u, S_1) < A(u, S) < A(u, S_2)$ for $S \in \Omega$ then $A(u, S) \in M$. Therefore since M is compact, $\{z \in C(u) \mid A(u, S_1) \leq z \leq A(u, S_2)\} \subset M$. So by Observation 3, M is not nowhere dense. This is a contradiction. Therefore $M \cap R_x = \emptyset$ and similarly $M \cap L_x = \emptyset$, which verifies Claim 1.3.

Suppose now that x is a cut point of $C(u)$ and $\{x\}$ is not a layer of $C(u)$. Then there is a nowhere dense nondegenerate subcontinuum M of $C(u)$ containing x . Let $y \in M - \{x\}$. Since x is a cut point of $C(u)$, $C(u) - \{x\}$ is the union of two mutually separated point sets H and K . Since

$L_x \cup R_x \subset C(u) - \{x\}$, by Claims 1.1 and 1.2 either $L_x \subset H$ and $R_x \subset K$ or $R_x \subset H$ and $L_x \subset K$. Without loss of generality assume $R_x \subset H$ and $L_x \subset K$. Furthermore by Claim 3 $M \cap (L_x \cup R_x) = \emptyset$.

Claim 1.4. y is a limit point of R_x .

Proof. Let O be an open set containing y . Let $A \in y$, $B \in x$, and U be a basic open set in βX so that

- (i) $A \cap B = \emptyset$,
- (ii) $A \subset U$,
- (iii) $\text{Cl}_{\beta X} U \subset 0$,
- (iv) $B \cap \text{Cl}_{\beta X} U = \emptyset$,
- (v) the number of components of $U \cap I_n$ is finite for all $n \in N$ (this can be done since $A \cap I_n$ is compact). Furthermore choose A so that $\{(0, i)\} \notin A$ for all i .

Let $N' = \{i | U \cap I_i \neq \emptyset\}$. Then $N' \in u$.

Let v_i be the rightmost component of $U \cap I_i$ (with respect to the usual order on $I_i = [0, 1] \times \{i\}$). Then if $i \in N'$ let $p_i \in v_i$ be a point such that $p_i > A \cap I_i$. If $P \in \Omega$ is such that $P(i) = p_i$ then $A(u, P) \in C(u)$, $A(u, P) > y$, and $A(u, P) \in U$.

Note that $p_i | i \in N'\} \cap B = \emptyset$ and $\{B \cap I_j | j \in N'\} \in x$.

Consider $B' = \{B \cap ([1, p_j] \times \{j\}) | j \in N'\}$. If $B' \in x$ then $x >_u A(u, P)$ so $x >_u y$. So $y \in L_x$, which contradicts Claim 1.3. Therefore $B'' = \{B \cap ([p_j, 0] \times \{j\}) | j \in N'\}$ is in x . Therefore $A(u, P) > x$. Therefore O contains a point of R_x . So y is a limit point of R_x . Thus Claim 1.4 is established.

Similarly it can be shown that y is a limit point of L_x . So y is a limit point of both H and K . But H and K are mutually separated. This is a contradiction so the lemma must be true.

Lemma 2. If $p \in C(u)$ and $L = \{q \in C(u) | \text{neither } p <_u q \text{ nor } q <_u p\}$ then L is a layer of $C(u)$.

Proof. Let $p \in L$ and let $L = \{q \in C(u) | \text{neither } p <_u q \text{ nor } q <_u p\}$. Then $L = \bigcap \{ \{x \in C(u) | A(u, s) \leq_u x \leq_u A(u, r)\} | A(u, s) < p < A(u, r) \}$. So L is a continuum. In order to verify that L is nowhere dense we only need to observe that if L is not nowhere dense in $C(u)$ then L would contain at least two cut points q_1 and q_2 of $C(u)$. But then one of them, q_i , is not p so either $q_i < p$ or $p < q_i$, which is a contradiction. In order to verify that L is a maximal nowhere dense subcontinuum we only need to observe that if L' is a subcontinuum of $C(u)$ with $L \subset L'$ then L' contains a point q so that either $p < q$ or $q < p$. In either case L' contains at least two cut points of $C(u)$ and hence is not nowhere dense in $C(u)$. Therefore L is a layer of $C(u)$.

Lemma 3. If L is a maximal nowhere dense subcontinuum of $C(u)$ and $p \in L$ then $L = \{q | \text{neither } p <_u q \text{ nor } q <_u p\}$.

Proof. If p is a cut point or an end point of $C(u)$ then the lemma follows from the observations above and Lemma 1.

Suppose that L is a nondegenerate maximal nowhere dense subcontinuum of $C(u)$. Let $p \in L$. Then since L is nondegenerate, it follows from the above observations that L contains no cut point of $C(u)$. Therefore if $q \in L$ then

$p \not\prec q$ and $q \not\prec p$. Therefore $\hat{L} = \{x | p \not\prec x \text{ and } x \not\prec p\}$ contains L . But $\hat{L}$ is nowhere dense, so since L is maximal, it follows that $L = \hat{L}$.

Notice that if G is the collection of layers of $C(u)$ then G is upper semi-continuous and $C(u)/G$ is a Hausdorff arc.

Lemmas 1, 2, and 3 (with a slight modification for the end points of $C(u)$) establish that L is a layer in $C(u)$ if and only if for each point $p \in L$,

$$L = \bigcap \{ \{x \in C(u) | A(u, s) \leq x \leq A(u, r)\} | A(u, s) < p < A(u, r); r, s \in \Omega \}.$$

We are now prepared to prove the main result.

Theorem. *Let $u \in N^*$ and let L be a layer of $C(u)$. Then L is indecomposable.*

Proof. Suppose that M is an arbitrary property subcontinuum of L . We will prove that M is nowhere dense in L , which implies that L is indecomposable. Let $\hat{U}$ and $\hat{V}$ be disjoint open sets in βX so that $\text{Cl}_{\beta X} \hat{U} \cap \text{Cl}_{\beta X} \hat{V} = \emptyset$, $M \cap \hat{V} \neq \emptyset$, $M \cap \text{Cl}_{\beta X} \hat{U} = \emptyset$, and $\hat{U} \cap L \neq \emptyset$. Let U and V be open sets in βX so that

$$\begin{aligned} U \subset \text{Cl}_{\beta X} U \subset \hat{U}, & \quad V \subset \text{Cl}_{\beta X} V \subset \hat{V}, \\ M \cap V \neq \emptyset, & \quad L \cap U \neq \emptyset, \end{aligned}$$

and for each n the number of components of $I_n \cap U$ and of $I_n \cap V$ is finite.

Let $p \in M \cap V$.

Claim 1. If k is an integer, $H \in p$, $H \subset V \cap X$, and J is the set of all integers n such that there is a sequence $U_1^n, U_2^n, \dots, U_{i_n}^n$ of components of $U \cap I_n$ and a sequence $V_1^n, V_2^n, \dots, V_{i_n}^n$ of components of $V \cap I_n$ that intersect H such that $U_1^n < V_1^n < U_2^n < V_2^n < \dots < U_{i_n}^n < V_{i_n}^n$ and $k \leq i_n$, then $J \in u$.

Proof. Let k be a positive integer and suppose the hypothesis of the claim and that $J \notin u$. So there is a sequence $\{s^i\}_{i=0}^{2k+3}$ of elements of Ω such that for each $n \in N - J$, $0 = s^0(n) \leq s^1(n) < s^2(n) < \dots < s^{2k+1}(n) < s^{2k+2}(n) \leq s^{2k+3}(n) = 1$ and

$$\begin{aligned} U \cap I_n &\subset \{x \in I_n | s^i(n) \leq x \leq s^{i+1}(n) \text{ and } i \text{ is even}\}, \\ V \cap I_n &\subset \{x \in I_n | s^i(n) \leq x \leq s^{i+1}(n) \text{ and } i \text{ is odd}\}. \end{aligned}$$

Note that if the first component of $V \cap I_n$ precedes the first component of $U \cap I_n$ it may be ignored. Let $H^i = \bigcup_{n \in N-J} \{x \in I_n | s^i(n) \leq x \leq s^{i+1}(n)\}$. Since $J \notin u$ we have $N - J \in u$ and then $\hat{H} = \{x \in H \cap I_n | n \in N - J\} \in p$. Then $\hat{H} \subset \bigcup_{i=1}^{2k+1} H^i$, so $H^1 \cup H^3 \cup \dots \cup H^{2k+1} \in p$, and hence $H^i \in p$ for some odd integer i . But then $A(u, s^i) \leq_u p \leq_u A(u, s^{i+1})$ and $U \cap \{z \in C(u) | A(u, s^i) \leq_u z \leq_u A(u, s^{i+1})\} = \emptyset$. This contradicts the fact that $U \cap L \neq \emptyset$, because $L \subset \{z \in C(u) | A(u, s^i) \leq_u z \leq_u A(u, s^{i+1})\}$. This establishes the claim.

For each n let $B_1^n, B_2^n, \dots, B_{k_n}^n$ be the components of $I_n - U$ that intersect V listed in order. If $H \in p$ let $B(H) = \{B_i^n | B_i^n \cap H \neq \emptyset\}$ and let $\mathcal{B} = \{B(H) | H \in p\}$. Define $\text{Ls}(\mathcal{B}) = \bigcap_{B \in \mathcal{B}} \text{Cl}_{\beta X}(\bigcup B)$.

Claim 2. $\text{Ls}(\mathcal{B})$ is a continuum, $p \in \text{Ls}(\mathcal{B})$, and $\text{Ls}(\mathcal{B})$ is a component of $X^* - U$.

Various forms of this claim have appeared elsewhere (see Blass [Bl], Smith [S1], and van Douwen [vD]). We will outline a proof for completeness.

Proof. Assume the hypothesis of the claim. Then the collection $\{\text{Cl}_{\beta X}(\bigcup B(H)) \mid H \in p\}$ is a collection of compact sets in βX so the common part $\text{Ls}(\mathcal{B})$ is compact. Since $p \in \text{Cl}_{\beta X}(\bigcup B(H))$ for each $H \in p$ we have $p \in \text{Ls}(\mathcal{B})$. Suppose that $\text{Ls}(\mathcal{B})$ is not a continuum but is the union of two mutually exclusive compact sets C_1 and C_2 . Let W_1 and W_2 be open sets in βX containing C_1 and C_2 respectively with disjoint closures. Then there exists an element $B(H)$ in $\mathcal{B}$ so that $\text{Cl}_{\beta X}(\bigcup B(H)) \subset W_1 \cup W_2$. But $X \cap W_1 \cap (\bigcup B(H))$ and $X \cap W_2 \cap (\bigcup B(H))$ are both compact and $(W_1 \cap (\bigcup B(H))) \cup (W_2 \cap (\bigcup B(H))) \in p$ so $W_i \cap (\bigcup B(H)) \in p$ for $i = 1$ or $i = 2$. (Note that each element of $B(H)$ is connected and lies in $W_1 \cup W_2$.) Assume $i = 1$. Let $\hat{H} = W_1 \cap (\bigcup B(H))$. Then $\hat{H} \in p$ and $\text{Cl}_{\beta X}(\bigcup B(\hat{H})) \subset \text{Cl}_{\beta X}(W_1)$, and hence $\text{Cl}_{\beta X}(\bigcup B(\hat{H})) \cap W_2 = \emptyset$ and so $\text{Ls}(\mathcal{B}) \cap W_2 = \emptyset$, which is a contradiction. It is easy to verify that $\text{Ls}(\mathcal{B})$ is a component of $(\bigcup_{n=1}^{\infty} \bigcup_{i=1}^{k_n} B_i^n)^* = X^* - U$.

Let $\widehat{M}$ denote $\text{Ls}(\mathcal{B})$. Since $\widehat{M}$ is a component of $X^* - U$, $p \in \widehat{M}$, $M \subset X^* - U$, and $p \in M$, it follows that $M \subset \widehat{M}$.

For each B_i^n let $p_i^n \in B_i^n \cap V$, and let $D = \{p_i^n\}_{i=1, n=1}^{k_n, \infty}$. If $d \in D$ then let $B(d)$ denote B_i^n where $d = p_i^n$. Let $D^* = \text{Cl}_{\beta X}(D) - D$.

Claim 3. Every component of $X^* \cap \text{Cl}_{\beta X}(\bigcup \{B_i^n \mid n \in N, 1 \leq i \leq k_n\})$ contains exactly one element of D^* .

Proof. Let $R = \{(n, i) \mid n \in N, 1 \leq i \leq k_n\}$. If $r \in R$ with $r = (n, i)$ let $Br = B_i^n$ and $dr = p_i^n$. If $\alpha \in R^*$ then let

$$d\alpha = \bigcap \{\text{Cl}_{\beta X}\{dr \mid r \in J\} \mid J \in \alpha\}.$$

Then $d\alpha$ is a single point of D^* and $d\alpha = dy$ iff $\alpha = y$. Furthermore C is a component of $X^* \cap \text{Cl}_{\beta X}(\bigcup \{B_i^n \mid n \in N, 1 \leq i \leq k_n\})$ if and only if there is an $\alpha \in R^*$ so that $C = B\alpha$ where

$$B\alpha = \bigcap \{\text{Cl}_{\beta X}\{Br \mid r \in J\} \mid J \in \alpha\}.$$

(See the argument to Claim 2 above and also [S1].) Furthermore $B\alpha = By$ if and only if $\alpha = y$. Therefore $d\alpha \in By$ if and only if $\alpha = y$; and this verifies Claim 3.

We have $D^* \subset V$ by construction. Consider the sets $\text{Even} = \bigcup \{B_i^n \mid i \text{ is even}\}$, $\text{Odd} = \bigcup \{B_i^n \mid i \text{ is odd}\}$. Then either $\text{Even} \in p$ or $\text{Odd} \in p$. Without loss of generality assume $\text{Even} \in p$. For i even, $1 \leq i \leq k_n$, and n a positive integer let $E_i^n = \{x \in I_n \mid x \in B_{i-1}^n \cup B_i^n \text{ or } B_{i-1}^n <_n x <_n B_i^n\}$. Then $B_i^n \subset E_i^n$. Let $E(J) = \{E_i^n \mid E_i^n \cap J \neq \emptyset\}$ and let $\mathcal{E} = \{E(J) \mid J \in p\}$. Let $\text{Ls}(\mathcal{E}) = \bigcap_{E \in \mathcal{E}} \text{Cl}_{\beta X}(\bigcup E)$, which is a continuum containing p by the same argument used to prove Claim 2. Since $\text{Even} \in p$ and $\text{Even} \subset E(\text{Even})$ we have $\widehat{M} = \text{Ls}(\mathcal{B}) \subset \text{Ls}(\mathcal{E})$.

Claim 4. $\text{Ls}(\mathcal{E}) \subset L$.

Proof. Suppose that the claim is not true. Since $p \in \text{Ls}(\mathcal{E})$ and $\text{Ls}(\mathcal{E}) \not\subset L$, there exists an element $s \in \Omega$ so that $A(u, s) \in \text{Ls}(\mathcal{E})$ and $A(u, s) \in V$.

But by Claim 1 $E(\{s_n | n \in N\}) \notin p$. So $p \notin \text{Cl}_{\beta X}(\bigcup E(\{s(n) | n \in N\}))$, and hence $p \notin \text{Cl}_{\beta X}(\bigcup \{E_i^n | s(n) \in E_i^n, n \in N, i \text{ even}, i \leq i \leq k_n\})$. So $E = \{E_i^n | s(n) \in E_i^n, n \in N, i \text{ even}, i \leq i \leq k_n\}$ is an element of $\mathcal{E}$ and $A(u, s) \notin \text{Cl}_{\beta X}(\bigcup E)$, which is a contradiction.

Let $z = d\alpha$ be the element of D^* that lies in $\widehat{M}$. Let $\hat{\alpha}$ be the element of R^* defined as follows: $\hat{\alpha} = \{G \in R | \text{there is a } J \in \alpha \text{ so that } \{(n, i - 1) | (n, i) \in J\} \subset G\}$. It is not difficult to verify that $\hat{\alpha} \in R^*$. But since $\{(n, i) | i \text{ is even}\} \in \alpha$ and $\{(n, i - 1) | i \text{ is even}\} \in \hat{\alpha}$, it follows that $\hat{\alpha} \neq \alpha$, so $d\hat{\alpha} \neq d\alpha$. Yet by construction since $B_i^n \cup B_{i-1}^n \subset E_i^n$ for $i \text{ even}, 1 \leq i \leq k_n$, it is easy to see that $d\hat{\alpha} \in \text{Ls}(\mathcal{E})$. Therefore $d\hat{\alpha}$ is a point of D^* distinct from $d\alpha$, $d\hat{\alpha} \in \text{Ls}(\mathcal{E}) \subset L$ so $d\hat{\alpha} \in L$, and since $\widehat{M} \cap D^* = \{d\alpha\}$, it follows that $d\hat{\alpha} \notin \widehat{M}$, and hence $d\hat{\alpha} \notin M$. So $d\hat{\alpha}$ is a point of $L \cap V$ that is not in M . So it follows that M is nowhere dense in L , and hence L is indecomposable.

REFERENCES

- [Be] D. P. Bellamy, *A non-metric indecomposable continuum*, Duke Math. J. **38** (1971), 15–20.
- [B1] A. Blass, *Near coherence of filters, II: Applications to operator ideals, the Stone-Ćech remainder of a half-line, order ideals of sequences, and slenderness of groups*, Trans. Amer. Math. Soc., **300** (1987), no. 2, 557–581.
- [BR] D. P. Bellamy and L. R. Rubin, *Indecomposable continua in Stone-Ćech compactifications*, Proc. Amer. Math. Soc. **39** (1973), 427–432.
- [BS] S. Baldwin and M. Smith, *On a possible property of far points of $\beta[0, \infty)$* , Topology Proc. **11** (1986), 239–245.
- [vD] E. K. van Douwen, *The number of subcontinua of the remainder of the plane*, Pacific J. Math. **97** (1981), no. 2, 349–355.
- [GJ] L. Gillman and M. Jerison, *Rings of continuous functions*, van Nostrand, Princeton, NJ, 1960.
- [J] T. Jech, *Set theory*, Academic Press, NY, 1968.
- [K] K. Kuratowski, *Topology I & II*, Academic Press, NY, 1968.
- [M] J. Mioduszewski, *On composants of $\beta R - R$* , Proc. Conf. Topology and Measure I (Zinnowitz, 1974) (J. Flachsmeier, Z. Fralik, and F. Terpe, eds.) Ernst-Meritz-Arndt-Universität zu Grifwald, 1978, pp. 257–283.
- [Mo] R. L. Moore, *Foundations of point-set theory*, Amer. Math. Soc. Colloq. Publ., vol. XII, Amer. Math. Soc., Providence, RI, 1962.
- [S1] M. Smith, *$\beta([0, \infty))$ does not contain non-degenerate hereditarily indecomposable continua*, Proc. Amer. Math. Soc. **101** (1987), 377–384.
- [S2] —, *$\beta(X - \{x\})$ for X not locally connected*, Topology Appl. **26** (1987), 239–250.
- [S3] —, *No arbitrary product of $\beta([0, \infty)) - [0, \infty)$ contains a nondegenerate hereditarily indecomposable continuum*, Topology Appl. **28** (1988), 23–28.
- [S4] —, *The subcontinua of $\beta[0, \infty) - [0, \infty)$* , Topology Proc. **11** (1986), 385–413.

DEPARTMENT OF FOUNDATIONS, ANALYSIS, AND TOPOLOGY, AUBURN UNIVERSITY, AUBURN, ALABAMA 36849-5310