DEFEAT OF THE FP^2F CONJECTURE: HUCKABA'S EXAMPLE

CARL FAITH

(Communicated by Louis J. Ratliff, Jr.)

ABSTRACT. A commutative ring R is FP^2F (resp. FPF) provided that all finitely presented (resp. finitely generated) faithful modules generate the category $\text{mod-}R$ of all R -modules. A conjecture of the author dating to the middle 1970s states that any FP^2F ring R has FP-injective classical quotient ring $Q = Q_{cl}(R)$. It was shown by the author (*Injective quotient rings*, II, Lecture Notes in Pure and Appl. Math., vol. 72, Dekker, New York, 1982, pp. 71–105) that FPF rings R have injective Q and by the author and P. Pillay (*Classification of commutative FPF rings*, Notas Math., vol. 4, Univ. de Murcia, Murcia, Spain, 1990) that CFP^2F local rings (defined below) have FP-injective Q .

The counterexample is a difficult example due to Huckaba of a strongly Prüfer ring without "Property A." (A ring with Property A was labelled a McCoy ring by the author.)

This counterexample is CFP^2F in the sense that every factor ring of R is FP^2F .

1. Theorem. (1) A ring R is CFP^2F iff (2) R_M is a valuation ring ($= VR$) for each maximal ideal M .

Proof. In [F1] (also [FP]) CFP^2F is characterized by the statement: R is locally a VR, i.e., R_P is a VR for all prime ideals P . However, a look at the proof in [F1, Corollary 5E, p. 176] establishes the equivalence of (1) and (2).

A ring R is said to be *McCoy* (see [F2]) or have Property A (see [H]) provided that every finitely generated dense ($=$ faithful) ideal is regular, i.e., contains a regular element. This is equivalent to the statement that $Q = Q_{cl}(R)$ is McCoy, i.e., that Q is the only finitely generated dense ideal. A sufficient condition for R to be McCoy is for Q to be FP-injective.

(Every FP-injective ring Q has the property that finitely generated ideals are annihilator ideals.)

A ring R is *Prüfer* if every finitely generated regular ideal is invertible (see [H, p. 29]), equivalently projective. A ring R is *strongly Prüfer* iff every finitely generated dense ideal is locally principal [H, p. 115].

Huckaba's example. There exists a strongly Prüfer ring R with the properties that R is not McCoy; hence $Q = Q_{cl}(R)$ is not FP-injective, but R_M is a VR, in fact a domain ($= VD$), for all maximal ideals. (See [H, Example 17, p. 191].)

Received by the editors February 12, 1991.

1991 *Mathematics Subject Classification*. Primary 13B20, 13C11, 13F05; Secondary 16A52.

© 1992 American Mathematical Society
0002-9939/92 \$1.00 + \$.25 per page

This provides the counterexample to the FP^2F conjecture since R is CFP^2F by Theorem 1, yet Q is not FP-injective.

We also remark the following:

Theorem (Huckaba and Keller). *A reduced coherent ring R is McCoy iff Q is VNR (= von Neumann regular).*

Proof. See [H, Theorem 4.7, p. 20].

REFERENCES

- [F1] C. Faith, *Injective quotient rings of commutative rings*, Module Theory, Lecture Notes in Math., vol. 700, Springer-Verlag, Berlin, Heidelberg, and New York, 1979, pp. 150–203.
- [F2] —, *Associated primes, annihilator ideals, and Kasch-McCoy commutative rings* (in memory of Robert Warfield), *Comm. Algebra* **119** (1991).
- [F3] —, *Injective quotient rings. II*, Lecture Notes in Pure and Applied Math., vol. 72, Dekker, New York, 1982, pp. 71–105.
- [FP] C. Faith and P. Pillay, *Classification of commutative FPF rings*, *Notas Mat.*, vol. 4, Univ. de Murcia, Murcia, Spain, 1990.
- [H] J. Huckaba, *Commutative rings with zero divisors*, Dekker, New York and Basel, 1988.
- [HK] J. Huckaba and Keller, *Annihilator ideals in commutative rings*, *Pacific J. Math.* **83** (1979), 375–379.

DEPARTMENT OF MATHEMATICS, RUTGERS UNIVERSITY, NEW BRUNSWICK, NEW JERSEY 08903

Current address: 199 Longview Drive, Princeton, New Jersey 08540