ON SOME PROPERTY OF FUNCTIONS DEFINED ON R^2 THAT ARE $\mathcal{I}$ -APPROXIMATELY CONTINUOUS WITH RESPECT TO ONE VARIABLE

R. CARRESE AND E. LAZAROW

(Communicated by Andrew M. Bruckner)

ABSTRACT. Balcerzak, Lazarow, and Wilczyński proved that every separately $\mathcal{I}$ -approximately continuous function is Baire 2. In this paper we shall prove that if f is a function $\mathcal{I}$ -approximately continuous with respect to one of its variables and of the α -class of Baire with respect to the other one, then f is of the $(\alpha + 1)$ -class of Baire in R^2 .

In this paper $\mathcal{I}$ denotes the σ -ideal of sets of the first category on the line and $\mathcal{B}$ denotes the σ -algebra of subsets of R having the Baire property. We shall say that 0 is an $\mathcal{I}$ -density point of a set $A \in \mathcal{B}$ if and only if for each increasing sequence of natural numbers $\{n_m\}_{m \in N}$ there exists a subsequence $\{n_{m_p}\}_{p \in N}$ such that the set $\{x: \chi_{n_{m_p} \cdot A \cap [-1, 1]} \rightarrow 1\} \in \mathcal{I}$, where χ_C denotes the characteristic function of the set C . We shall say that x_0 is an $\mathcal{I}$ -density point $A \in \mathcal{B}$ if and only if 0 is an $\mathcal{I}$ -density point of the set $A - x_0 = \{x - x_0: x \in A\}$. We shall say that x_0 is an $\mathcal{I}$ -dispersion point of $A \in \mathcal{B}$ if and only if x_0 is an $\mathcal{I}$ -density point of $R - A$. The family of all sets $A \in \mathcal{B}$ such that each point of A is its $\mathcal{I}$ -density point forms a topology called the $\mathcal{I}$ -density topology. The functions that are continuous with respect to the $\mathcal{I}$ -density topology are called $\mathcal{I}$ -approximately continuous (see [4]).

We shall say that $f: R^2 \rightarrow R$ is $\mathcal{I}$ -approximately continuous in the direction x (resp. y) at (x_0, y_0) if the function $f(x, y_0)$ (resp. $f(x_0, y)$) is $\mathcal{I}$ -approximately continuous at x_0 (resp. y_0) as a function of x (resp. y). We shall say that $f: R^2 \rightarrow R$ is separately approximately continuous if and only if f is $\mathcal{I}$ -approximately continuous in the direction x and y simultaneously.

In the sequel we shall need the following

Proposition. *Let G be an open set of the real line; then 0 is an $\mathcal{I}$ -dispersion point of G if and only if for every natural number n there exist a natural number k and a real number $\delta > 0$ such that for each $h \in (0, \delta)$ and for each $i \in \{1, \dots, n\}$ there exist two natural numbers $j, j' \in \{1, \dots, k\}$ such that*

$$G \cap \left(\left(\frac{i-1}{n} + \frac{j-1}{nk} \right) h, \left(\frac{i-1}{n} + \frac{j}{nk} \right) h \right) = \emptyset$$

Received by the editors February 25, 1991.

1991 *Mathematics Subject Classification.* Primary 26B05, 26A21.

©1992 American Mathematical Society
0002-9939/92 \$1.00 + \$.25 per page

and

$$G \cap \left(- \left(\frac{i-1}{n} + \frac{j'}{nk} \right) h, \left(\frac{i-1}{n} + \frac{j'-1}{nk} \right) h \right) = \emptyset.$$

For the proof see [2, Theorem 1].

We shall say that x_0 is a deep $\mathcal{F}$ -density point of the set $A \in \mathcal{B}$ if there exists an open set $B \supset R - A$ such that x_0 is an $\mathcal{F}$ -dispersion point of B . The family of all sets $A \in \mathcal{B}$ such that each point of A is its deep $\mathcal{F}$ -density point forms a topology called the deep $\mathcal{F}$ -density topology. The $\mathcal{F}$ -approximately continuous functions are continuous with respect to this topology (see [5, 3]).

Lemma. Let $f: R^2 \rightarrow R$ be a function $\mathcal{F}$ -approximately continuous at (x_0, y_0) in the direction x . Then $f(x_0, y_0) \leq \alpha$ if and only if for every $n \in N$ there exist $k \in N$, $k > n$, and n rational numbers $r_1^n, \dots, r_n^n$ such that for each $i \in \{1, \dots, n\}$, $r_i^n \in (x_0 + (i-1)/nk, x_0 + i/nk)$, and $f(r_i^n, y_0) < \alpha + 1/n$.

Proof. For simplicity assume that $(x_0, y_0) = (0, 0)$. Let $n \in N$; if $f(0, 0) \leq \alpha$ then $f(0, 0) < \alpha + 1/n$ and $(0, 0)$ is a deep $\mathcal{F}$ -density point of the set $\{(x, 0) : f(x, 0) < \alpha + 1/n\}$ in the direction x . Therefore there exist $p \in N$ and $\delta > 0$ such that for each $i \in \{1, \dots, n\}$ and for each $h < \delta$ there exists $j \in \{1, \dots, k\}$ such that

$$\left[\frac{(i-1)p + j - 1}{np} h, \frac{(i-1)p + j}{np} h \right] \times \{0\} \subset \left\{ (x, 0) : f(x, 0) < \alpha + \frac{1}{n} \right\}.$$

Let $k > \max\{n, 1/\delta\}$. Then for each $i \in \{1, \dots, n\}$ there exists a rational number

$$r_i^n \in \left(\frac{(i-1)p + j - 1}{np} \frac{1}{k}, \frac{(i-1)p + j}{np} \frac{1}{k} \right) \subset \left[\frac{i-1}{nk}, \frac{i}{nk} \right]$$

such that $f(r_i^n, 0) < \alpha + 1/n$.

Conversely, for each $n \in N$ let k be a natural number greater than n , and let $r_1^n, \dots, r_n^n$ be rational numbers such that for each $i \in \{1, \dots, n\}$ we have $r_i^n \in ((i-1)/nk, i/nk)$ and $f(r_i^n, 0) < \alpha + 1/n$. We shall show that $f(0, 0) \leq \alpha$.

Let $A = \bigcup_{n=1}^{\infty} \{r_1^n, \dots, r_n^n\}$. We shall show that for each $p \in N$ and for each $\delta > 0$ there exists $h < \delta$ such that for each $j \in \{1, \dots, p\}$, $((j-1)h/p, jh/p) \cap A \neq \emptyset$. Let $p \in N$ and $\delta > 0$. Choose $n \in N$ such that $n > \max\{2p, \frac{1}{\delta}\}$. Then there exist $k > n$ and $r_1^n, \dots, r_n^n$ such that for each $i \in \{1, \dots, n\}$, $r_i^n \in ((i-1)/nk, i/nk)$. Put $h = 1/k < 1/n < \delta$.

Since $n > 2p$, we observe that for each $j \in \{1, \dots, p\}$ there exists $i \in \{1, \dots, n\}$ such that

$$\left[\frac{i-1}{nk}, \frac{i}{nk} \right] \subset \left(\frac{j-1}{p} \frac{1}{k}, \frac{j}{p} \frac{1}{k} \right).$$

Therefore for each $j \in \{1, \dots, p\}$,

$$\left(\frac{j-1}{p} \frac{1}{k}, \frac{j}{p} \frac{1}{k} \right) \cap A \neq \emptyset.$$

If $f(0, 0) > \alpha$ then there exists $s \in N$ such that $f(0, 0) > \alpha + 1/s$.

Since f is an $\mathcal{F}$ -approximately continuous function in the direction x at $(0, 0)$, then there exists a closed set $F \subset \{(x, 0) : f(x, 0) > \alpha + 1/s\}$ such

that $(0, 0)$ is a point of $\mathcal{F}$ -density of F in the direction x . Thus there exist $p_0 \in N$ and $\delta > 0$ such that for each $h < \delta$ there exists $j \in \{1, \dots, p_0\}$ such that $[(j-1)h/p_0, jh/p_0] \times \{0\} \subset F$.

Let $h_1 < \delta$ such that for each $j \in \{1, \dots, p_0\}$, $((j-1)h_1/p_0, jh_1/p_0) \cap A \neq \emptyset$.

Then there exists $r_1 \in N$ such that for each $j \in \{1, \dots, p_0\}$ we have

$$\left(\frac{j-1}{p_0}h_1, \frac{j}{p_0}h_1\right) \cap \{r_1^{n_1}, \dots, r_{n_1}^{n_1}\} \neq \emptyset.$$

Now let $h_2 < r_1$ such that for each $j \in \{1, \dots, p_0\}$ we have $((j-1)h_2/p_0, jh_2/p_0) \cap A \neq \emptyset$. Then there exists $n_2 > n_1$, $n_2 \in N$, such that for each $j \in \{1, \dots, p_0\}$,

$$\left(\frac{j-1}{p_0}h_2, \frac{j}{p_0}h_2\right) \cap \{r_1^{n_2}, \dots, r_{n_2}^{n_2}\} \neq \emptyset.$$

By induction we define sequences $\{h_m\}_{m \in N}$ and $\{n_m\}_{m \in N}$ such that for each $m \in N$ we have $h_m < \delta$,

$$\left(\frac{j-1}{p_0}h_m, \frac{j}{p_0}h_m\right) \cap \{r_1^{n_m}, \dots, r_{n_m}^{n_m}\} \neq \emptyset$$

and $\lim_m h_m = 0$, $\lim_m n_m = \infty$.

Thus, if m_0 is such that $n_{m_0} > s$ then for each $j \in \{1, \dots, n_{m_0}\}$ we have $f(r_i^{n_{m_0}}, 0) < \alpha + 1/n_{m_0} < \alpha + 1/s$. Therefore for each $j \in \{1, \dots, p_0\}$,

$$\left(\frac{j-1}{p_0}h_{m_0}, \frac{j}{p_0}h_{m_0}\right) \cap \left\{(x, 0) : f(x, 0) < \alpha + \frac{1}{s}\right\} \neq \emptyset,$$

a contradiction.

Then $f(0, 0) \leq \alpha$ and the proof is completed.

An analogous proposition obviously holds for $(x_0, y_0) \in R^2$ such that $f(x_0, y_0) \geq \alpha$ and for f $\mathcal{F}$ -approximately continuous at (x_0, y_0) in the direction y .

Theorem. Let $f: R^2 \rightarrow R$ be a function such that for each $(x_0, y_0) \in R^2$, f is $\mathcal{F}$ -approximately continuous at (x_0, y_0) in the direction x and of the Baire class α (with respect to the natural topology) in the direction y . Then f is a function of the Baire class $\alpha + 1$ on the plane.

Proof. By the previous lemma we have that for each $\lambda \in R$

$$\begin{aligned} & \{(x, y) : f(x, y) \leq \lambda\} \\ &= \bigcap_{n=1}^{\infty} \bigcup_{k=n+1}^{\infty} \bigcup_{\{r_1^n, \dots, r_n^n\} \in Q} \bigcap_{i=1}^n \left(\left\{ x : \frac{i-1}{nk} < |x - r_i^n| < \frac{i}{nk} \right\} \times R \right) \\ & \quad \cap \left(R \times \left\{ y : f(r_i^n, y) < \lambda + \frac{1}{n} \right\} \right), \end{aligned}$$

where Q denotes the set of rational numbers.

If f is of the Baire class α in the direction y at every $(x, y) \in R^2$ then the set $\{y : f(r_i^n, y) < \lambda + 1/n\}$ belongs to F_α if α is odd and to G_α if α is even.¹

¹For each ordinal α such that $0 < \alpha < \omega_1$, F_α and G_α denote the Borel classes with respect to the natural topology.

Therefore $\{(x, y) : f(x, y) \leq \lambda\}$ belongs to $F_{\alpha+1}$ if α is odd and to $G_{\alpha+1}$ if α is even.

REFERENCES

1. M. Balcerzak, E. Lazarow, and W. Wilczyński, *On one and two-dimensional $\mathcal{F}$ -densities and related kinds of continuity*, Real Anal. Exchange **13** (1987–88), 80–120.
2. E. Lazarow, *On the Baire class of $\mathcal{F}$ -approximate derivatives*, Proc. Amer. Math. Soc. **100** (1987), 669–674.
3. —, *The coarsest topology for $\mathcal{F}$ -approximately continuous functions*, Comment. Math. Univ. Carolin. **27** (1986), 695–704.
4. W. Poreda, E. Wagner-Bojakowska, and W. Wilczyński, *A category analogue of the density topology*, Fund. Math. **128** (1985), 167–173.
5. E. Wagner-Bojakowska and W. Poreda, *The topology of $\mathcal{F}$ -approximately continuous functions*, Rad. Mat. **2** (1986), 263–277.

DIPARTIMENTO DI MATEMATICA E APPLICAZIONI "R. CACCIOPOLI" VIA CINTIA NAPOLI, ITALY
INSTITUT OF MATHEMATICS, UL. STEFANA BANACHA, 22 90-238 ŁÓDŹ, POLAND