WEIGHTED NORM INEQUALITIES FOR BOCHNER-RIESZ OPERATORS AND SINGULAR INTEGRAL OPERATORS

XIANLIANG SHI AND QIYU SUN

(Communicated by J. Marshall Ash)

ABSTRACT. Weighted norm inequalities for the Bochner-Riesz operator at the critical index $\frac{1}{2}(n-1)$ are investigated. We also give some weighted norm inequalities for a class of singular integral operators introduced by Fefferman and Namazi.

1. INTRODUCTION AND STATEMENTS

The Bochner-Riesz operators in R^n are defined as

$$(T_\lambda^R f)^\wedge(x) = (1 - R^2|x|^2)_+^\lambda \hat{f}(x)$$

and the associated maximal operator is defined as

$$T_\lambda^* f(x) = \sup_{R>0} |T_\lambda^R f(x)|$$

for $\lambda > 0$, where $\wedge$ denotes the Fourier transform. It is well known by the works of Carleson and Sjölin [4], Fefferman [8, 9], Tomas [19], and Christ [6] that T_λ^f is bounded on $L^p(R^n)$ if and only if $|1/p - 1/2| < (1 + 2\lambda)/2n$ provided $\lambda > 0$ in dimension 2 and $\lambda \geq (n-1)/2(n+1)$ in dimension greater than two. Rubio [16] and Hirschman [12] studied the weighted norm inequality for the Bochner-Riesz operator T_λ^f and showed that T_λ^f is bounded on $L^2(|x|^a)$ provided $|a| < 1 + 2\lambda < n$. In 1988 Andersen [1] gave a sufficient condition and a necessary condition on radial weight $w(|x|)$ such that the inequality

$$\int_{R^n} |T_\lambda^1 f(x)|^2 w(|x|) dx \leq C \int_{R^n} |f(x)|^p w(|x|) dx$$

holds for all radial functions f in $L^p(w(|x|))$.

Notice that the Bochner-Riesz operator is a summation operator and $T_\lambda^R f(x)$ tends to $f(x)$ as R tends to infinity for all Schwartz functions f . Hence it is meaningful to consider the almost everywhere convergence of $T_\lambda^R f$ as R tends to infinity for some appropriate function f . In 1986 Lu [14] proved that

$$T_\lambda^{R_j} f(x) \rightarrow f(x) \quad \text{a.e. as } j \rightarrow \infty$$

Received by the editors March 2, 1990 and, in revised form, June 25, 1990 and September 7, 1990.

1991 *Mathematics Subject Classification.* Primary 42B15, 42B25.

for all $f \in L^2(|x|^a)$ provided $0 < a < \min(2, 2\lambda) < n-1$ and $\{R_j\}_{j=1}^\infty$ being a Hadamard lacunary sequence, i.e., $\lim_{j \rightarrow \infty} (R_{j+1}/R_j) > 1$. In addition we can reduce almost everywhere convergence of the operator T_λ^R to some maximal inequality. In [14] Lu proved

Theorem L [14]. *Let $0 < \lambda < \frac{1}{2}(n-1)$, $0 < a < \min(2, 2\lambda)$, and $\{R_j\}_{j=1}^\infty$ be a Hadamard lacunary sequence. Then*

$$\int \left(\sup_j |T_\lambda^{R_j} f(x)| \right)^2 |x|^a dx \leq C \int |f(x)|^2 |x|^a dx.$$

In 1988 Carbery, Rubio, and Vega proved

Theorem CRV [3]. *Let $|a| < 1 + 2\lambda < n$. Then*

$$\int |T_\lambda^* f(x)|^2 |x|^a dx \leq C \int |f(x)|^2 |x|^a dx.$$

Hence they improved Theorem L.

On other hand, we observe that if λ exceeds the critical index $\frac{1}{2}(n-1)$ then $T_\lambda^* f$ is dominated by a multiple of the Hardy-Littlewood maximal function Mf defined by

$$Mf(x) = \sup_{x \in Q} |Q|^{-1} \int_Q |f(y)| dy,$$

where supremum is taken over all cubes with center x and sides parallel to the coordinate axes. Hence a result of Muckenhoupt [11] showed that T_λ^* is bounded on $L^p(w)$ provided $w \in A_p$, i.e.,

$$\left(|Q|^{-1} \int_Q w(x) dx \right) \left(|Q|^{-1} \int_Q w(x)^{-(p-1)^{-1}} dx \right)^{p-1} \leq C$$

holds for all cubes $Q \subset \mathbb{R}^n$ with sides parallel to the coordinate axes and some C independent of Q . Then a natural question is whether $T_{(n-1)/2}^*$ is bounded on $L^p(w)$ provided $w \in A_p$ and $1 < p < \infty$. In this paper we prove

Theorem 1. *Let $1 < p < \infty$ and $w \in A_p$. Then*

$$\int |T_{(n-1)/2}^* f(x)|^p w(x) dx \leq C \int |f(x)|^p w(x) dx.$$

We also observe that $T_{(n-1)/2}^1 f$ can be written as

$$\int h(|y|) |y|^{-n} f(x-y) dy,$$

where

$$h(t) = (2\pi)^{n/2} 2^{(n-1)/2} \Gamma(\tfrac{1}{2}(n-1)) J_{n-1/2}(t) t^{1/2},$$

$\Gamma(t)$ denotes the Gamma function and $J_v(t)$ denotes the Bessel function, which is defined by

$$J_v(t) = \frac{(t/2)^v}{\Gamma(v+1/2)\Gamma(v)} \int_0^{\pi/2} \cos(t \sin u) (\cos u)^{2v} du.$$

Therefore the weighted norm inequality for $T_{(n-1)/2}^*$ is closely related to the one for the operator introduced by Fefferman [9] and Namazi [15].

Now let us write the operator introduced by Fefferman and Namazi precisely. Let $h \in L^\infty([0, \infty))$ and Ω be an integrable function on the unit sphere S^{n-1} having mean zero, i.e., $\int_{S^{n-1}} \Omega(x) d\sigma(x) = 0$, $d\sigma$ is the standard measure on S^{n-1} .

Define

$$T_\varepsilon f(x) = \int_{|y|>\varepsilon} h(|y|) \Omega\left(\frac{y}{|y|}\right) |y|^{-n} f(x-y) dy$$

for $\varepsilon > 0$,

$$T_0 f(x) = \lim_{\varepsilon \rightarrow 0} T_\varepsilon f(x),$$

and the associated maximal operator

$$T^* f(x) = \sup_{\varepsilon > 0} |T_\varepsilon f(x)|.$$

There are many works about the operators T_0 and T^* (see [5, 7, 15, 17, 18], etc.). Namazi [15] proved that T_0 is a bounded operator on $L^p(\mathbb{R}^n)$ for all $1 < p < \infty$ when $\Omega \in L^q(S^{n-1})$ for some $q > 1$, Chen [5] proved that T^* is also a bounded operator on $L^p(\mathbb{R}^n)$ when $\Omega \in L^q(S^{n-1})$ for some $q > 1$, and the second author [18] proved that $\Omega \in L \log^+ L(S^{n-1})$ (resp. $L(\log^+ L)^{3/2}(S^{n-1})$) is a sufficient condition such that T_0 (resp. T^*) is a bounded operator on $L^2(\mathbb{R}^n)$. As weighted norm inequalities for the operators T_0 and T^* , Duoandikoetxea and Rubio [7] proved that T^* and T_0 are bounded operators on $L^p(w)$ provided $1 < p < \infty$, $w \in A_p$, and $\Omega \in L^\infty(S^{n-1})$.

In this paper we prove the following with the complex interpolation method.

Theorem 2. Let $1 < q \leq \infty$, $\Omega \in L^q(S^{n-1})$, $q(q-1)^{-1} < p < \infty$, and $w \in A_{p(1-1/q)}$. Then

$$\int |T_0 f(x)|^p w(x) dx \leq C \int |f(x)|^p w(x) dx$$

holds for all f in $L^p(w)$.

Theorem 3. Let $\Omega \in L^\infty(S^{n-1})$ and $w \in A_p$. Then

$$\int |T^* f(x)|^p w(x) dx \leq C \int |f(x)|^p w(x) dx$$

holds for all f in $L^p(w)$.

Hence we prove Duoandikoetxea and Rubio's result in another way. The above results are still interesting even when $h \equiv 1$ because in [13] Ω satisfies an L' -Dini condition for some $r > 1$.

2. SOME LEMMAS

Define the Bochner-Riesz operator T_z^R by

$$(T_z^R f)^\wedge(x) = (1 - R^2|x|^2)_+^z f^\wedge(x)$$

for $\operatorname{Re} z > 0$. Then we have

Lemma 1 [3]. *Let $\operatorname{Re} z > 0$ and $k = 0, 1$. Then*

$$\int \sup_{R>0} \left| \left(\frac{\partial}{\partial z} \right)^k T_z^R f(x) \right|^2 dx \leq C e^{c|\operatorname{Im} z|} (|\operatorname{Re} z|^{-c} + 1) \int |f(x)|^2 dx$$

holds for a constant C independent of z .

To prove Theorem 1, we will also use

Lemma 2. *Let $\operatorname{Re} z > \frac{1}{2}(n-1)$. Then the inequality*

$$\sup_{R>0} |T_z^R f(x)| \leq C (\operatorname{Re} z - \frac{1}{2}(n-1))^{-c} e^{c|\operatorname{Im} z|} M f(x)$$

holds for all $f \in L_{\text{loc}}^1(\mathbb{R}^n)$ and an absolutely positive constant C .

Proof. Write

$$T_z^R f(x) = \frac{2^{z+1-n/2} \Gamma(z+1)}{\Gamma(n/2)} R^{-n/2+z} \int J_{n/2+z}(R^{-1}|y|) \cdot |y|^{-z-n/2} f(x-y) dy.$$

Also we observe from the asymptotic properties of the Bessel function $J_{n/2+z}(t)$ that

$$|J_{n/2+z}(t)| \leq C t^{n/2+\operatorname{Re} z}$$

when $t \leq 1$ and

$$|J_{n/2+z}(t)| \leq C t^{-1/2}$$

when $t > 1$. Therefore for $\operatorname{Re} z > \frac{1}{2}(n-1)$ we have

$$\begin{aligned} |T_z^R f(x)| &\leq C R^{-n/2+\operatorname{Re} z} \int_{|y| \leq R} |f(x-y)| R^{-n/2-\operatorname{Re} z} dy \\ &\quad + C R^{n/2+\operatorname{Re} z} \int_{|y| > R} |f(x-y)| |y|^{-\operatorname{Re} z-(n+1)/2} R^{-1/2} dy \\ &\leq C M f(x) \end{aligned}$$

and Lemma 2 holds.

Lemma 3 [11]. *For $s \in (1, \infty)$ and $w \in A_s$, there exists a positive number δ such that $w^{1+\delta} \in A_s$.*

Lemma 4 (Three-Circles Theorem). *Suppose F is a bounded continuous complex-valued function on the closed strip $S = \{x+iy; 0 \leq x \leq 1\}$ that is analytic in the interior of S . If $|F(iy)| \leq m_0$ and $|F(1+iy)| \leq m_1$ for all y , then $|F(x+iy)| \leq m_0^{1-x} m_1^x$ for all $x+iy \in S$.*

To prove Theorems 2 and 3, we need to introduce some notation and use some lemmas. Let $\Omega \subset L^q(S^{n-1})$ for some $q > 1$.

Define

$$T_{z,\varepsilon} f = \frac{\pi^{(z-1)/2} \Gamma((n-z)/2)}{\Gamma(z/2)} |y|^{-n-z} h(|y|) \Omega\left(\frac{y}{|y|}\right) \chi_{|y|>\varepsilon} * |y|^{-n+z} * f, \quad \varepsilon > 0,$$

for $-\frac{1}{2}(1-1/q) < \operatorname{Re} z < 1$, where $*$ is the convolution operation. Denote the kernel function of the operator $T_{z,0}$ by K_z . Therefore we have

Lemma 5 [18]. For $-\frac{1}{2}(1 - 1/q) < \operatorname{Re} z < 1$ and $k = 0, 1$, the inequality

$$\int \sup_{\varepsilon > 0} \left| \left(\frac{\partial}{\partial z} \right)^k T_{z, \varepsilon} f(x) \right|^2 dx \leq C \left(|\operatorname{Re} z - 1|^{-c} + \left| \operatorname{Re} z + \frac{1}{2} \left(1 - \frac{1}{q} \right) \right|^{-c} \right) e^{c|\operatorname{Im} z|} \int |f(x)|^2 dx$$

holds.

Lemma 6. For $0 < \operatorname{Re} z < 1$ and $1 < p < \infty$, the inequality

$$\int \sup_{\varepsilon > 0} |T_{z, \varepsilon} f(x)|^p dx \leq C(|\operatorname{Re} z|^{-c} + |\operatorname{Re} z - 1|^{-c}) e^{c|\operatorname{Im} z|} \int |f(x)|^p dx$$

holds.

Lemma 7. For $0 < \operatorname{Re} z < 1$, the inequality

$$\left(R^{-n} \int_{R < |x| < 2R} |K_z(x+y) - K_z(x)|^q dx \right)^{1/q} \leq C(z) R^{-n} \left(\frac{|y|}{R} \right)^{\operatorname{Re} z}$$

holds for all $R > 0$ and $|y| < \frac{1}{2}R$, where

$$C(z) \leq C(|\operatorname{Re} z|^{-C} + |\operatorname{Re} z - 1|^{-C}) \exp(C|\operatorname{Im} z|).$$

Lemma 8 [13]. Let $K \in L^1_{\text{loc}}(R^n \setminus \{0\})$, $q > 1$, and $\delta > 0$. Suppose

$$\left(R^{-n} \int_{R < |x| < 2R} |K(x+y) - K(x)|^q dx \right)^{1/q} \leq C R^{-n} \left(\frac{|y|}{R} \right)^\delta$$

holds for all $R > 0$ and $|y| < \frac{1}{2}R$. Suppose $\bar{T}$ be defined as $\bar{T}f = K * f$ and $\bar{T}$ be bounded on $L^2(R^n)$. Then the operator $\bar{T}^-$ is bounded on $L^p(w)$ for all $q(q-1)^{-1} < p < \infty$ and $w \in A_{p(1-1/q)}$.

Lemma 9. Let $\Omega \in L^q(S^{n-1})$ for some $q > n$. For $n/q < \operatorname{Re} z < 1$, the inequalities

$$(i) \quad |K_z(x)| \leq C(z)|x|^{-n},$$

$$(ii) \quad |K_z(x+y) - K_z(x)| \leq C(z)|x|^{-n-(\operatorname{Re} z - n/q)}|y|^{\operatorname{Re} z - n/q}$$

hold for all $x \neq 0$ and $|y| \leq \frac{1}{2}|x|$, where

$$C(z) \leq C(|\operatorname{Re} z - n/q|^{-c} + |\operatorname{Re} z - 1|^{-c}) \exp(C|\operatorname{Im} z|).$$

Lemma 10 [11]. Let $K \in L^1_{\text{loc}}(R^n \setminus \{0\})$. Suppose

$$(i) \quad |K(x)| \leq C|x|^{-n},$$

$$(ii) \quad |K(x+y) - K(x)| \leq C|y|^\delta |x|^{-n-\delta}$$

hold for all $x \neq 0$, $|y| < \frac{1}{2}|x|$, and some $\delta > 0$. Suppose an operator $\bar{T}$ defined as $\bar{T}f = K * f$ is bounded on $L^2(R^n)$. Then the operator $\bar{T}^*$ defined as

$$\bar{T}^* f(x) = \sup_{\varepsilon > 0} \left| \int_{|y| > \varepsilon} k(y) f(x-y) dy \right|$$

is bounded on $L^p(w)$ provided $1 < p < \infty$ and $w \in A_p$.

The proofs of Lemmas 7 and 9 are elementary and the proof of Lemma 6 is similar to the one in [5]. We omit the details here.

3. PROOFS OF THEOREMS

Let $p \in (1, \infty)$, $s \in (1, \infty)$, $0 < \lambda \leq \frac{1}{2}(n-1)$, and $q > n$. Denote $\alpha(p) = \frac{1}{p} - \frac{1}{2}$, $p^1 = p(p-1)^{-1}$,

$$\theta_1(p, s, \lambda) = \begin{cases} \frac{2\lambda}{n-1}, & \text{when } p = s = 2, \\ \frac{\alpha(p)}{\alpha(s)} \cdot \frac{p}{s}, & \text{when } 0 < \frac{\alpha(p)}{\alpha(s)} < \frac{2\lambda}{n-1}, p \neq 2, \\ 0, & \text{otherwise;} \end{cases}$$

$$\theta_2(p, s, q) = \begin{cases} \left(1 + \frac{2n}{q-1}\right)^{-1}, & \text{when } p = s = 2, \\ \frac{\alpha(p)}{\alpha(s)} \cdot \frac{p}{s}, & \text{when } 0 < \frac{\alpha(p)}{\alpha(s)} \leq \left(1 + \frac{2n}{q-1}\right)^{-1}, p \neq 2, \\ \frac{1}{2} \left(1 - \frac{\alpha(p)}{\alpha(s)}\right) \left(1 - \frac{1}{q}\right) \frac{p}{s} \cdot \frac{q}{n}, & \\ & \text{when } \left(1 + \frac{2n}{q-1}\right)^{-1} < \frac{\alpha(p)}{\alpha(s)} < 1, \\ 0, & \text{otherwise.} \end{cases}$$

Then the following theorems are general versions of Theorems 1 and 3.

Theorem 4. Let $0 < \lambda \leq \frac{1}{2}(n-1)$, $|\alpha(p)| < \lambda/(n-1)$, and $w \in A_s$. Then

$$\int |T_\lambda^* f(x)|^p w^{\theta_1(p, s, \lambda)}(x) dx \leq C \int |f(x)|^p w^{\theta_1(p, s, \lambda)}(x) dx.$$

Theorem 5. Let $\Omega \in L^q(S^{n-1})$ for some $q > n$ and $w \in A_s$. Then

$$\int |T^* f(x)|^p w^{\theta_2(p, s, q)}(x) dx \leq C \int |f(x)|^p w^{\theta_2(p, s, q)}(x) dx.$$

By Lemma 3 and the fact $|x|^a \in A_s$ for $-n < a < n(s-1)$, we improve Theorem L.

Proof of Theorem 4. Let f, h be two nonnegative smooth functions with compact support and $R(x)$ be any arbitrary positive measurable function bounded below and above, i.e., $R(x)^{-1}$ and $R(x)$ bounded. Suppose $w \in A_s$ and ε_1 and ε_2 are sufficiently small positive constants chosen later, without loss of generality we assume $\theta_1(p, s, \lambda) > 0$. Let

$$p^{-1} = \frac{1}{2}(1 - \theta) + s^{-1}\theta, \quad \lambda = (1 + \theta)\varepsilon_1 + \theta\left(\frac{1}{2}(n-1) + \varepsilon_2\right), \quad 0 < \theta < 1.$$

Denote $s(z)^{-1} = \frac{1}{2}(1 - z) + s^{-1}z$ for $0 < \operatorname{Re} z < 1$ and $f_0(x) = \exp(-|x|^2)$.

Define

$$\begin{aligned} f_{\delta_1, \delta_2}^z(x) &= (f(x) + \delta_1 f_0(x))^{ps(z)^{-1}-1} (w(x) + \delta_2)^{-s^{-1}z} f(x), \\ h_{\delta_3}^z(x) &= (h(x) + \delta_3 f_0(x))^{p'(1-s(z)^{-1})-1} h(x), \\ w_N(x) &= \begin{cases} w(x), & \text{when } w(x) \leq N, \\ N, & \text{when } w(x) > N, \end{cases} \end{aligned}$$

and

$$\tilde{T}_{z, \varepsilon_1, \varepsilon_2, \delta_3, N}^{R(x)}(f)(x) = T_{\varepsilon_1 + z((n-1)/2 + \varepsilon_2 - \varepsilon_1)}^{R(x)} f(x) (w_N(x) + \delta_2)^{zs^{-1}} e^{-z^2}$$

where N^{-1} , δ_i ($i = 1, 2, 3$) are small positive numbers.

Hence by Lemma 1 we can show easily

$$g(z) = \int \tilde{T}_{z, \varepsilon_1, \varepsilon_2, \delta_2, N}^{R(x)}(f_{\delta_1, \delta_2}^z)(x) h_{\delta_3}^z(x) dx$$

is analytic in the strip $0 < \operatorname{Re} z < 1$ and continuous on the closed strip $0 \leq \operatorname{Re} z \leq 1$. In addition $g(z)$ is bounded function, hence by Lemma 4 we have

$$|g(\theta)| \leq C \left(\sup_{t \in \mathbb{R}} |g(it)| \right)^{1-\theta} \left(\sup_{t \in \mathbb{R}} |g(1+it)| \right)^\theta.$$

On the other hand by Lemma 1 we get

$$\begin{aligned} \sup_{t \in \mathbb{R}} |g(it)| &\leq C_{\varepsilon_1, \varepsilon_2} \sup_{t \in \mathbb{R}} e^{-t^2} e^{c|t|} \left(\int |f(x)|^p dx \right)^{1/2} \left(\int |h(x)|^{p'} dx \right)^{1/2} \\ &\leq C_{\varepsilon_1, \varepsilon_2} \|f\|_p^{p/2} \|h\|_{p'}^{p'/2} \end{aligned}$$

and by Lemma 2

$$\begin{aligned} \sup_{t \in \mathbb{R}} |g(1+it)| &\leq C_{\varepsilon_1, \varepsilon_2} \sup_{t \in \mathbb{R}} e^{-t^2} \left(\int |T_{1+it, \varepsilon_1, \varepsilon_2, \delta_2, N}^{R(x)}(f_{\delta_1, \delta_2}^{1+it})(x)|^s dx \right)^{1/s} \\ &\quad \times \left(\int |h_{\delta_3}^{1+it}(x)|^{s(s-1)^{-1}} dx \right)^{s-1/3} \\ &\leq C_{\varepsilon_1, \varepsilon_2} \|f\|_p^{ps^{-1}} \|h\|_{p'}^{p's^{-1}(s-1)}. \end{aligned}$$

Therefore

$$(1) \quad |g(\theta)| \leq C_{\varepsilon_1, \varepsilon_2} \|f\|_p \|h\|_{p'}$$

where $C_{\varepsilon_1, \varepsilon_2}$ is independent of $\delta_1, \delta_2, \delta_3$, and N .

Write $g(\theta)$ as

$$\begin{aligned} (2) \quad g(\theta) &= \int \tilde{T}_{\theta, \varepsilon_1, \varepsilon_2, \delta, N}^{R(x)}(f(w + \delta_2)^{-\theta s^{-1}})(x) h(x) dx \\ &= \int T_{\lambda}^{R(x)}(f(w + \delta_2)^{-\theta s^{-1}})(x) (w_N(x) + \delta_2)^{\theta s^{-1}} h(x) dx. \end{aligned}$$

Notice that for every $f \in L^{p'}$ there exist two nonnegative smooth function sequences $\{f_n^1\}$ and $\{f_n^2\}$ with compact supports such that

$$\|f_n^i\|_{p'} \leq C \|f\|_{p'}, \quad i = 1, 2, \quad n \in \mathbb{N},$$

and

$$\|f_n^1 - f_n^2 - f\|_{p'} \rightarrow 0, \quad \text{as } n \rightarrow \infty.$$

Hence by (1) and (2) we get

$$\int |T_\lambda^{R(x)}(f(w + \delta_2)^{-\theta s^{-1}})(x)|^p (w_N(x) + \delta_2)^{p\theta s^{-1}} dx \leq C \int |f(x)|^p dx.$$

Let N tend to infinity in the inequality above. Then we have

$$\int |T_\lambda^{R(x)}(f(w + \delta_2)^{-\theta s^{-1}})(x)|^p (w(x) + \delta_2)^{p\theta s^{-1}} dx \leq C \int |f(x)|^p dx.$$

Notice that C is independent of δ_2 and $R(x)$ being a measurable function bounded below and above. Therefore

$$(3) \quad \int |T_\lambda^* f(x)|^p w^{p\theta s^{-1}}(x) dx \leq C \int |f(x)|^p w^{p\theta s^{-1}}(x) dx$$

holds for all $\theta < \theta_1(p, s, \lambda) \neq 0$ by choosing appropriate ε_1 and ε_2 . By Lemma 3 we can replace w in (3) by $w^{1+\delta}$. Hence Theorem 4 holds by choosing $\theta = \theta_1(p, s, x)/(1 + \delta) \neq 0$.

Proof of Theorem 2. By Lemmas 7 and 8, T_z is bounded on $L^p(w)$ provided $w \in A_{p(1-1/q)}$ and $\operatorname{Re} z > 0$. In addition T_z is bounded on $L^2(R^n)$ when $|\operatorname{Re} z| < \frac{1}{2}(1 - 1/q)$. By the complex interpolation theorem (see [10]) and Lemma 3, T_0 is bounded on $L^p(w)$ provided $p > q(q - 1)^{-1}$ and $w \in A_{p(1-1/q)}$. Therefore Theorem 2 holds.

Proof of Theorem 5. Define

$$T_z^* f(x) = \sup_{\varepsilon > 0} \left| \int_{|y| > \varepsilon} K_z(y) f(x - y) dy \right|.$$

Then by Lemmas 9 and 10 we get

$$\int |T_z^* f(x)|^p w(x) dx \leq C(z) \int |f(x)|^p w(x) dx$$

provided $w \in A_p$ and $n/q < \operatorname{Re} z < 1$. In addition the pointwise estimate

$$\sup_{\varepsilon > 0} |T_{z, \varepsilon} f(x)| \leq C(z) \mu f(x) + T_z^* f(x)$$

holds for all x in R^n , $n/q < \operatorname{Re} z < 1$, and

$$C(z) \leq C(|\operatorname{Re} z - 1|^{-c} + |\operatorname{Re} z - n/q|)^{-c} e^{c|\operatorname{Im} z|}.$$

Hence we have

$$(4) \quad \int \sup_{\varepsilon > 0} |T_{z, \varepsilon} f(x)|^p w(x) dx \leq C(z) \int |f(x)|^p w(x) dx$$

for $n/q < \operatorname{Re} z < 1$ and $w \in A_p$. By (4), Lemmas 3, 5, and 6 we can prove the following in such a way as we prove Theorem 4,

$$\begin{aligned} & \int \sup_{\varepsilon > 0} |T_{z, \varepsilon} f(x)|^s w^{(\operatorname{Re} z)n/q}(x) dx \\ & \leq C(z) \int |f(x)|^s w^{(\operatorname{Re} z)n/q}(x) dx \end{aligned}$$

provided $0 < \operatorname{Re} z < n/q$, $w \in A_s$, and $C(z) \leq C|\operatorname{Re} z|^{-C} \exp(2|\operatorname{Im} z|^2)$.

Hence

$$\int \sup_{\varepsilon > 0} |T_{s,\varepsilon} f|^p(x) w^{\theta_2(p,s,q)}(x) dx \leq C \int |f(x)|^p w^{\theta_2(p,s,q)}(x) dx$$

holds for $1 < p < \infty$, $q > n$, and $w \in A_s$ and Theorem 5 holds.

ACKNOWLEDGMENTS

The authors would like to thank the referee for some corrections.

REFERENCES

1. K. F. Andersen, *Weighted norm inequalities for Bochner-Riesz spherical summation multipliers*, Proc. Amer. Math. Soc. **103** (1988), 165–170.
2. A. Carbery, *A weighted inequality for the maximal Bochner-Riesz operator on $\mathbb{R}^2$* , Trans. Amer. Math. Soc. **289** (1985), 673–680.
3. A. Carbery, J. L. Rubio de Francia, and L. Vega, *Almost everywhere summability of Fourier integrals*, J. London Math. Soc. (2) **38** (1988), 513–524.
4. L. Carleson and P. Sjolin, *Oscillatory integrals and a multiplier problem for the disc*, Studia Math. **44** (1972), 287–299.
5. L.-K. Chen, *On a singular integral*, Studia Math. **85** (1987), 61–72.
6. M. Christ, *Weak type endpoint bounds for Bochner-Riesz multiplier*, Rev. Mat. Iberoamericana **3** (1987), 25–31.
7. J. Duoandikoetxea and J. L. Rubio de Francia, *Maximal and singular integral operators via Fourier transform estimate*, Invent. Math. **84** (1986), 541–561.
8. C. Fefferman, *Inequalities for strongly singular convolution operators*, Acta Math. **124** (1970), 9–36.
9. R. Fefferman, *A note on singular integral*, Proc. Amer. Math. Soc. **74** (1979), 266–270.
10. J. Garcia-Cuerva and J. L. Rubio de Francia, *Weighted norm inequalities and related topics*, North-Holland, Amsterdam, 1985.
11. B. Muckenhoupt, *Weighted norm inequalities for the Hardy maximal function*, Trans. Amer. Math. Soc. **165** (1972), 207–226.
12. I. I. Hirschman, *Multiplier transforms. II*, Duke Math. J. **28** (1961), 45–50.
13. D. S. Kurtz and R. L. Wheeden, *Results on weighted norm inequalities for multiplier*, Trans. Amer. Math. Soc. **255** (1979), 343–362.
14. S.-Z. Lu, *A class of maximal functions on spherical Riesz means*, Sci. Sinica **29** (1986), 113–137.
15. J. Namazi, *On a singular integral*, Proc. Amer. Math. Soc. **96** (1986), 421–424.
16. J. L. Rubio de Francia, *Weighted norm inequalities and vector valued inequalities*, Lecture Notes in Math., vol. 908, Springer-Verlag, Berlin and New York, 1982, pp. 86–101.
17. X. Shi, *Some remarks on singular integrals*, Indiana Univ. Math. J. **35** (1986), 103–116.
18. Q. Sun, *Two problems about singular integral operators*, Approx. Theory Appl. **7** (1991), 83–98.
19. P. Tomas, *Restriction theorem for the Fourier transform*, Proc. Sympos. Pure Math., vol. 35, Amer. Math. Soc., Providence, RI, 1979, pp. 111–114.

DEPARTMENT OF MATHEMATICS, HANGZHOU UNIVERSITY, HANGZHOU, ZHEJIANG 310028, PEOPLE'S REPUBLIC OF CHINA

Current address, X. Shi: Department of Mathematics, Texas A & M University, College Station, Texas 77843-3368

E-mail address: shi@wavelet1.math.tamu.edu