A NOTE ON THE LARGEST REGULAR SUBALGEBRA OF A BANACH ALGEBRA

JYUNJI INOUE AND SIN-EI TAKAHASI

(Communicated by Paul S. Muhly)

ABSTRACT. A simple proof of Albrecht's result on the existence of the largest closed regular subalgebra of a semisimple commutative Banach algebra is given.

Albrecht [1] proved that any semisimple commutative Banach algebra with identity has the largest closed regular subalgebra, using the theory of decomposable operators. In this short note, we give a simple proof of Albrecht's result without the assumption of semisimplicity. Our proof is quite different from Albrecht's, and based on the consideration of the hull-kernel topology of the carrier space of the given Banach algebra.

Theorem. *If A is a commutative Banach algebra with identity, then there exists a closed regular subalgebra of A which contains all closed regular subalgebras of A .*

Our proof is essentially based on the following

Lemma. *Let X be a commutative Banach algebra with identity and B a Banach subalgebra of X . If B is regular, then for any $b \in B$ the Gelfand transform of b as an element of X is continuous on the carrier space Φ_X of X in the hull-kernel topology.*

Proof. We can assume without loss of generality that B contains the identity of X . Then it is sufficient to show that the restriction map $\theta: \Phi_X \rightarrow \Phi_B; \varphi \rightarrow \varphi|_B$ is continuous in the hull-kernel topology. To do this let F be a closed subset of Φ_B in the hull-kernel topology. Then $\{\varphi \in \Phi_X: \varphi|_{\ker F} = 0\} = \theta^{-1}(F)$. Also since $\ker F \subset \ker \theta^{-1}(F)$, it follows that $\text{hul}(\ker \theta^{-1}(F)) \subset \{\varphi \in \Phi_X: \varphi|_{\ker F} = 0\}$. Therefore $\theta^{-1}(F)$ is closed in the hull-kernel topology. In other words, θ is continuous in this topology. $\square$

Proof of Theorem. Let $\text{reg}(A)$ be the closed subalgebra of A generated by the class of all closed regular subalgebras of A . It is sufficient to show that $\text{reg}(A)$ is regular. Note that $\text{reg}(A)$ is a commutative Banach algebra with identity. If

Received by the editors November 29, 1989.

1980 *Mathematics Subject Classification* (1985 Revision). Primary 46H05.

Key words and phrases. Commutative Banach algebra, regular, hull-kernel topology.

Supported in part by a Grant-in-Aid for Scientific Research from the Japanese Ministry of Education.

b is an element of A which belongs to a certain regular Banach subalgebra B of A , then $B \subset \text{reg}(A)$. Hence by the preceding lemma the Gelfand transform of b as an element of $\text{reg}(A)$ is continuous on $\Phi_{\text{reg}(A)}$ in the hull-kernel topology. Since $\text{reg}(A)$ is generated by such b 's, the Gelfand transform of an arbitrary element of $\text{reg}(A)$ is also continuous on $\Phi_{\text{reg}(A)}$ in the hull-kernel topology. Consequently $\text{reg}(A)$ is regular. $\square$

REFERENCES

1. E. Albrecht, *Decomposable systems of operators in harmonic analysis*, Toeplitz Centennial (I. Gohberg, ed.), Birkhäuser, Basel, 1982, pp. 19–35.

DEPARTMENT OF MATHEMATICS, HOKKAIDO UNIVERSITY, SAPPORO 060, JAPAN

DEPARTMENT OF BASIC TECHNOLOGY, YAMAGATA UNIVERSITY, YONEZAWA 992, JAPAN