TOEPLITZ OPERATORS ON CARTAN DOMAINS ESSENTIALLY COMMUTE WITH A BILATERAL SHIFT

MIROSLAV ENGLIŠ

(Communicated by Theodore W. Gamelin)

ABSTRACT. For bounded symmetric domains $\Omega \subset \mathbb{C}^N$, a bilateral shift operator U is shown to exist on the Bergman space $A^2(\Omega)$ such that $UT_f - T_fU$ is a compact operator for all Toeplitz operators T_f . This may be viewed as an extension of the well-known fact that $S^*TS - T = 0$ whenever T is a Toeplitz operator on H^2 , S being the unilateral shift. It also follows that the C^* -algebra generated by Toeplitz operators on $A^2(\Omega)$ does not contain all bounded operators.

Let Ω be a bounded symmetric (Cartan) domain with its standard (Harish-Chandra) realization in $\mathbb{C}^N$ [6], dv the $2n$ -dimensional Lebesgue measure on Ω , and $L^2(\Omega, dv)$ the Hilbert space of square-integrable complex-valued functions on Ω . The Bergman space, $A^2(\Omega)$, is the closed subspace of $L^2(\Omega, dv)$ consisting of functions analytic on Ω . Denote by P the orthogonal projection from L^2 onto A^2 . For $f \in L^\infty(\Omega)$, the Toeplitz operator $T_f: A^2 \rightarrow A^2$ and the Hankel operator $H_f: A^2 \rightarrow L^2 \ominus A^2$ are given by

$$T_fx = P(fx), \quad H_fx = (I - P)(fx).$$

These operators generalize, in an obvious way, the well-known Toeplitz and Hankel operators on H^2 [5, Chapter 25].

The main result of the present note is the following theorem.

Theorem. *There is a bilateral shift U on $A^2(\Omega)$ such that for all $f \in L^\infty(\Omega)$ the commutator $[U, T_f] \equiv UT_f - T_fU$ is a compact operator.*

Proof. We first introduce some notation and terminology from [1, 2]. Let $\beta(\cdot, \cdot)$ be the Bergman metric on Ω [6] and $\text{dist}(\cdot, \cdot)$ the usual euclidean metric in $\mathbb{C}^N = \mathbb{R}^{2N}$. Denote by $\text{BC}(\Omega)$ the algebra of bounded continuous complex-valued functions on Ω , with $C_\partial(\Omega)$ the subalgebra of functions for which $f(z) \rightarrow 0$ as $\text{dist}(z, \partial\Omega) \rightarrow 0$. Define

$$\text{Osc}(f, z) = \sup\{|f(w) - f(z)|: \beta(z, w) < 1\},$$

the oscillation of f at z , and

$$\text{VO}_\partial(\Omega) = \{f \in \text{BC}(\Omega): \text{Osc}(f, \cdot) \in C_\partial(\Omega)\},$$

Received by the editors May 30, 1991.

1991 *Mathematics Subject Classification.* Primary 47B35; Secondary 32M15.

Key words and phrases. Toeplitz operators, bounded symmetric domains.

the algebra of functions with *vanishing oscillation at the boundary*.

Consider the function Φ on Ω given by

$$\Phi(z) = \exp(i\sqrt{\beta(0, z)}).$$

It is well known [1] that $\Phi \in \text{VO}_\partial(\Omega)$. Indeed,

$$\begin{aligned} |\Phi(z) - \Phi(w)| &\leq |\beta(0, z)^{1/2} - \beta(0, w)^{1/2}| \\ &\leq \frac{|\beta(0, z) - \beta(0, w)|}{\beta(0, z)^{1/2} + \beta(0, w)^{1/2}} \\ &\leq \frac{\beta(z, w)}{\beta(0, z)^{1/2} + \beta(0, w)^{1/2}} \end{aligned}$$

and the assertion is immediate since $\beta(0, x) \rightarrow +\infty$ as $\text{dist}(x, \partial\Omega) \rightarrow 0$. Hence, by [2, Theorem B], H_Φ and $H_{\bar{\Phi}}$ are compact operators. According to a well-known commutator identity for Toeplitz operators,

$$T_f T_g - T_g T_f = H_g^* H_f - H_f^* H_g$$

for arbitrary $f, g \in L^\infty(\Omega)$. Taking $g = \Phi$, we see that the commutator $T_f T_\Phi - T_\Phi T_f \equiv [T_f, T_\Phi]$ is compact $\forall f \in L^\infty(\Omega)$. Thus the proof will be accomplished if we find a compact operator K such that $T_\Phi + K$ is a bilateral shift with respect to some basis. Since, in particular, $[T_{\bar{\Phi}}, T_\Phi]$ is compact, T_Φ is essentially normal. Hence, by the Brown-Douglas-Fillmore theory [3], it suffices to show that the essential spectrum of T_Φ is the unit circle, $\mathbb{T}$, and that $\text{ind } T_\Phi = 0$.

Denote

$$\mathcal{Q} = \{f \in L^\infty(\Omega) : H_f \text{ and } H_{\bar{f}} \text{ are compact}\},$$

and let $\tau(\mathcal{Q})$ be the C^* -algebra generated by $\{T_f : f \in \mathcal{Q}\}$. By [1, Theorem B], there is a C^* -isomorphism

$$(*) \quad \tau(\mathcal{Q})/\text{Compacts} \simeq \text{VO}_\partial/C_\partial(\Omega)$$

which maps T_f into the coset $[f]$ of f in $\text{VO}_\partial/C_\partial$. It follows that, for $f \in \mathcal{Q}$, $\sigma_e(T_f)$ coincides with the spectrum of $[f]$ in $\text{VO}_\partial/C_\partial(\Omega)$. Since $[f]$ is invertible in $\text{VO}_\partial/C_\partial$ iff it is invertible in L^∞/C_∂ , the latter spectrum is easily seen to coincide with the set of all cluster values of f at $\partial\Omega$,

$$\sigma_{\text{VO}_\partial/C_\partial}([f]) = \bigcap_{R>0} \overline{\{f(z) : \beta(0, z) > R\}}.$$

Taking $f = \Phi$, we conclude that

$$\sigma_e(T_\Phi) = \bigcap_{R>0} \overline{\{\Phi(z) : \beta(0, z) > R\}} = \mathbb{T},$$

which proves the first claim.

To compute the Fredholm index of T_Φ , consider the functions

$$\Phi_m(z) = \exp\left(\frac{i}{m}\sqrt{\beta(0, z)}\right),$$

where m is a positive integer (thus, $\Phi_1 = \Phi$). Everything that was said about Φ is readily seen to apply to Φ_m as well: Φ_m is a bounded continuous function that belongs to VO_∂ , so that H_{Φ_m} , $H_{\bar{\Phi}_m}$ are compact operators and $\Phi_m \in \mathcal{Q}$;

further, $\sigma_e(T_{\Phi_m}) = \mathbf{T}$, so T_{Φ_m} are Fredholm operators. Since $\Phi_m^m = \Phi$, it follows from (*) that $T_{\Phi_m}^m = T_\Phi$ modulo the compacts. Thus, we have

$$\operatorname{ind} T_\Phi = \operatorname{ind} T_{\Phi_m}^m = m \cdot \operatorname{ind} T_{\Phi_m}$$

for all positive integers m , which is only possible when $\operatorname{ind} T_\Phi = 0$. The proof is complete.

For Toeplitz operators on the N -dimensional Fock space $A^2(\mathbb{C}^N)$, $N \geq 1$, a similar result was obtained by the present author in [4]. It is also shown there that for arbitrary bounded planar domain $\Omega \subset \mathbb{C}$ there exists a *unilateral* shift operator S on the Bergman space $A^2(\Omega)$ such that, for all Toeplitz operators T_f , $ST_f - T_fS$ is compact. The assertion was also proved for the one-dimensional Fock space $A^2(\mathbb{C})$. These results may be compared with the classical characterization [5, Problem 242] of Toeplitz operators on H^2 :

$$T \text{ is a Toeplitz operator} \Leftrightarrow S^*TS = T,$$

S being the unilateral shift on H^2 .

As a direct consequence of the above theorem, we have

Corollary. *The C^* -algebra generated by all Toeplitz operators $\{T_f: f \in L^\infty(\Omega)\}$, where Ω is a bounded symmetric domain in $\mathbb{C}^N$, does not contain all bounded operators on $A^2(\Omega)$.*

Proof. Let $U = T_\Phi + K$ be the bilateral shift obtained above. Since

$$\begin{aligned} [A + B, U] &= [A, U] + [B, U], & [cA, U] &= c[A, U], \\ [A^*, U] &= U[A, U]^*U, & [AB, U] &= A[B, U] + [A, U]B, \end{aligned}$$

and

$$\|A_n - A\| \rightarrow 0 \text{ implies } \|[A_n, U] - [A, U]\| \rightarrow 0,$$

we see that the essential commutant, U^{ess} , of U is a C^* -algebra. According to the preceding theorem, it contains all operators T_f , $f \in L^\infty(\Omega)$. It follows that U^{ess} contains the C^* -algebra generated by them as well, and we only have to find an operator not belonging to U^{ess} . Let $\{e_n\}_{n \in \mathbb{Z}}$ be a basis with respect to which U is the bilateral shift, i.e.,

$$Ue_n = e_{n+1}, \quad n \in \mathbb{Z},$$

and let J be given by

$$Je_n = (-1)^n e_n, \quad n \in \mathbb{Z}.$$

Then $JU - UJ = 2JU$ is not compact, and so $J \notin U^{\text{ess}}$.

REFERENCES

1. C. A. Berger, L. A. Coburn, and K. H. Zhu, *Function theory on Cartan domains and the Berezin-Toeplitz symbol calculus*, Amer. J. Math. **110** (1988), 921–953.
2. D. Békollé, C. A. Berger, L. A. Coburn, and K. H. Zhu, *BMO in the Bergman metric on bounded symmetric domains*, J. Funct. Anal. **93** (1990), 310–350.
3. L. G. Brown, R. G. Douglas, and P. A. Fillmore, *Unitary equivalence modulo the compact operators and extensions of C^* -algebras*, Lecture Notes in Math., vol. 345, Springer-Verlag, Berlin and Heidelberg, 1973, pp. 58–128.

4. M. Engliš, *Density of algebras generated by Toeplitz operators on Bergman spaces*, Ark. Mat. (to appear).
5. P. R. Halmos, *A Hilbert space problem book*, Graduate Texts in Math., Springer-Verlag, New York, 1967.
6. S. Helgason, *Differential geometry, Lie groups and symmetric spaces*, Academic Press, New York, 1978.

MATHEMATICAL INSTITUTE, CZECHOSLOVAK ACADEMY OF SCIENCES, ŽITNÁ 25, 115 67 PRAGUE 1, CZECHOSLOVAKIA

E-mail address: englis@cspgas11.bitnet

Current address: Department of Mathematics, Kansas State University, Manhattan, Kansas 66506

E-mail address: englis@ksuvm.ksu.edu