TWO THEOREMS OF JOSEFSON-NISSENZWEIG TYPE FOR FRÉCHET SPACES

J. BONET, M. LINDSTRÖM, AND M. VALDIVIA

(Communicated by William J. Davis)

ABSTRACT. We characterize the Fréchet-Montel (respectively, Fréchet-Schwartz) spaces by sequences in their dual spaces.

It is well known that a Fréchet space E is Montel (respectively, Schwartz) if and only if it is separable and every weak*-null sequence in E' also converges strongly to zero (respectively, converges uniformly to zero on some zero-neighbourhood in E). Because of the Josefson-Nissenzweig theorem [3], Jarcho asked in [5] if these two results are true without the assumption of the separability condition on the Fréchet space E .

In this short note we give a positive answer to both of these questions. These results were discovered during the Conference of Function Theory on Infinite-Dimensional Spaces held in the Universidad Complutense de Madrid from December 11–16, 1989. The Schwartz case had already been discovered independently by Lindström and Schlumprecht in [7] and by Bonet in [1].

We start by recalling the following result from [7], which is based on a very deep result due to Schlumprecht [8].

Theorem 1. *Let E be a Fréchet space such that every weak*-null sequence in E' is also strongly null-convergent. Then E is reflexive.*

Now we are ready to prove the announced results.

Theorem 2. *A Fréchet space E is Montel (respectively, Schwartz) if and only if weak*-null sequence in E' is also strongly null-convergent (respectively, converges uniformly to zero on some zero-neighbourhood in E).*

Proof. Only the Montel case needs a proof because every Fréchet-Montel space is separable.

If E is Montel then the condition is clearly necessary.

Let us now assume that every weak*-null sequence in E' is strongly null-convergent. By the theorem above [7] E must then be reflexive. Now we present two different proofs to show that E is Montel.

First, by [6, Remark 1] it is enough to show that U° is $\sigma(E', E)$ -sequentially compact for every zero-neighbourhood U in E . To do this observe that E'_β

Received by the editors January 25, 1991.

1991 *Mathematics Subject Classification.* Primary 46A04, 46A11.

©1993 American Mathematical Society
0002-9939/93 \$1.00 + \$.25 per page

is a DF-space whose strong dual is E by reflexivity. Hence $\sigma(E', (E'_\beta)') = \sigma(E', E)$. Now according to [2, Theorem 11, Example 1.2(C)], every DF-space is weakly angelic (see, e.g., [4]), and since U° is always $\sigma(E', E)$ -compact, it is also $\sigma(E', E)$ -sequentially compact.

To give the second proof assume that E is not Montel. Since E is reflexive by the theorem above [7], there exists a weakly compact separable subset A of E that is not compact. Let F be the closed linear span of A . By a recent result of Valdivia in [9], given F we can find a closed separable subspace G of E that contains F and has a topological complement L in E . Certainly G is separable but not Montel. Hence there is a $\sigma(G', G)$ -null sequence in $L^\perp = G'$ that is not $\beta(G', G)$ -null convergent. Since G is complemented in E , this contradicts the assumption on E .

REFERENCES

1. J. Bonet, *A question of Valdivia on quasi-normable Fréchet spaces*, *Canad. Math. Bull.* **34** (1991), 301–304.
2. B. Cascales and J. Orihuela, *On compactness in locally convex spaces*, *Math. Z.* **195** (1987), 365–381.
3. J. Diestel, *Sequences and series in Banach spaces*, Graduate Texts in Math., vol. 92, Springer, Berlin, Heidelberg and New York, 1984.
4. K. Floret, *Weakly compact sets*, Lecture Notes in Math., vol. 801, Springer-Verlag, New York, 1980.
5. H. Jarchow, *Locally convex spaces*, Teubner, Stuttgart, 1981.
6. M. Lindström and Th. Schlumprecht, *On limitedness in locally convex spaces*, *Arch. Math.* **53** (1989), 65–74.
7. —, *A Josefson-Nissenzweig theorem for Fréchet spaces*, *Bull. London Math. Soc.* (to appear).
8. Th. Schlumprecht, *On dual spaces with bounded sequences without weak* convergent convex blocs*, *Proc. Amer. Math. Soc.* **107** (1989), 395–408.
9. M. Valdivia, *Resolution of the identity operator in certain metrizable locally convex spaces*, *Rev. Real Acad. Ci. Madrid* **83** (1989), 75–96.

DEPARTAMENTO DE MATEMÁTICA APLICADA, E.T.S. ARQUITECTURA, UNIVERSIDAD POLITÉCNICA DE VALENCIA, E-46071 VALENCIA, SPAIN
E-mail address: JBONET@MAT.UPV.ES

DEPARTMENT OF MATHEMATICS, ABO AKADEMI, SF-20500 ABO, FINLAND
E-mail address: MLINDSTROM@FINABO.ABO.FI

DEPARTAMENTO DE ANÁLISIS MATEMÁTICO, FACULTAD DE MATEMÁTICAS, UNIVERSIDAD DE VALENCIA, 46100 BURJASOT, SPAIN