UPPER BOUND OF $\sum 1/(a_i \log a_i)$ FOR PRIMITIVE SEQUENCES

PAUL ERDÖS AND ZHENXIANG ZHANG

(Communicated by William W. Adams)

ABSTRACT. A sequence $A = \{a_i\}$ of positive integers $a_1 < a_2 < \dots$ is said to be primitive if no term of A divides any other. The senior author conjectures that, for any primitive sequence A ,

$$\sum_{a \leq n, a \in A} \frac{1}{a \log a} \leq \sum_{p \leq n} \frac{1}{p \log p} \quad \text{for } n > 1,$$

where p is a variable prime. In our two previous papers we partially proved this conjecture. The main result of this paper is: for any primitive sequence A ,

$$\sum_{a \in A} \frac{1}{a \log a} < 1.84.$$

We also give a necessary and sufficient condition for this conjecture, i.e.,

$$\sum_{b \in B} \frac{1}{b \log b} \leq \sum_{p \log p}$$

for any primitive sequence B .

1. INTRODUCTION

A sequence $A = \{a_i\}$ of positive integers $a_1 < a_2 < \dots$ is said to be primitive if no term of A divides another [3, 4]. We denote by p_m the m th prime and by p a variable prime. We define the degree of an integer a , denoted by $\Omega(a)$, to be the number of prime factors of a , counted with multiplicity. The degree of a primitive sequence A , denoted by $d^\circ(A)$, is defined as the maximum degree of its terms. We take $d^\circ(A) = 0$ if $A = \{1\}$ or $\emptyset$.

We define $f(A) = \sum_{a \in A} 1/(a \log a)$. We take $f(A) = 0$ if $d^\circ(A) = 0$. Erdős [1] (cf. [4]) proved that there exists an absolute constant C such that $f(A) < C$ for any primitive sequence, using the inequality

$$(1) \quad \sum_{a \in A} \frac{1}{a} \prod_{p \leq P(a)} \left(1 - \frac{1}{p}\right) \leq 1,$$

where $P(a)$ is the largest prime factor of a . It is significant to get a reasonable upper bound for $f(A)$. Using (1) we first got $f(A) < 2.886$; then Robin

Received by the editors January 16, 1991 and, in revised form, July 9, 1991.

1991 *Mathematics Subject Classification*. Primary 11B83.

Key words and phrases. Primitive sequences.

improved it to 2.77 (neither is published). It seems to us that we could not do better by only using (1).

Erdős [2] has asked if the inequality

$$(2) \quad \sum_{a \leq n, a \in A} \frac{1}{a \log a} \leq \sum_{p \leq n} \frac{1}{p \log p} \quad \text{for } n > 1$$

is always true for any A . Zhang [7] proved that if A is primitive with $d^\circ(a) \leq 4$ then (2) is true. For a given primitive sequence A and $m \geq 1$, let $A_m = \{a : a \in A, \text{ all prime factors of } a \text{ are } \geq p_m\}$ and $A'_m = \{a : a \in A_m, p_m | a\}$. Zhang [8] proved that (2) is true for any primitive sequence A with the property that each A_m is "homogeneous." That is, for each m , there is some integer s_m such that either $A_m = \emptyset$ or $\Omega(a) = s_m$ for all $a \in A_m$.

In this paper we prove the following two theorems.

Theorem 1. $f(A) < 1.84$ for any primitive sequence $A = \{a_i\}$.

Theorem 2. The necessary and sufficient condition for (2) being true for any primitive sequence A is that $f(B) \leq \sum 1/(p \log p)$ for any primitive sequence B .

The basic idea for proving Theorem 1 is the same as that used in [7, 8]. That is, we consider the least prime factors of terms of A . For a given primitive sequence A and $m \geq 1$, with A_m and A'_m already defined, let $A''_m = \{a/p_m : a \in A'_m\}$. Clearly all of them are primitive. We study upper bounds F_m of $f(A_m)$. To find F_m we can suppose A_m is finite by Lemma 3. At first we find F_m for $m \geq N + 1$, where $N = 10^5$, by induction on $d^\circ(A_m)$. It is based on the fact that

$$f(A_m) = \sum_{i \geq m} f(A'_i) \quad \text{and} \quad f(A'_i) \leq \frac{f(A''_i)}{p_i}.$$

Then we find F_m from F_{m+1} for $m = N, N-1, \dots, 2, 1$. It is based on the fact that

$$f(A_m) = f(A_{m+1}) + f(A'_m) \quad \text{and} \quad f(A'_m) \leq f(A''_m)/p_m.$$

At last

$$f(A) = f(A_1) \leq F_1 < 1.84.$$

2. PROOFS OF THE THEOREMS

We have three constants: $N = 10^5$, $\alpha = 1.0072629$, and $k = 0.0072847$. We need five lemmas.

Lemma 1. We have $p_n \geq n(\log n + \log \log n - \alpha)$ for $n \geq 2$ [5].

Lemma 2. We have

$$\sum_{i \geq m} \frac{1}{p_i(\log i + \log \log i - k)} < \frac{1}{\log m + \log \log m - k}$$

for $m \geq N + 1$.

Proof. Put $h(m) = \sum_{i \geq m} 1/\{p_i(\log i + \log \log i - k)\}$,

$$g(m) = 1/(\log m + \log \log m - k),$$

$$c = (1 + 1/\log N)/N, \\ H(y) = (y + \log y - \alpha)(y + \log y - k), \text{ and} \\ G(y) = (y + \log y + c - k)^2.$$

Then by Lemma 1,

$$h(m+1) < \int_m^\infty \frac{dx}{x(\log x + \log \log x - \alpha)(\log x + \log \log x - k)} = \int_{\log m}^\infty \frac{dy}{H(y)}, \\ g(m+1) > \frac{1}{\log m + 1/m + \log(\log m + 1/m) - k} \\ > \frac{1}{\log m + \log \log m + (1 + 1/\log m)/m - k} \\ \geq \frac{1}{\log m + \log \log m + c - k} \\ = \int_{y=\log m}^\infty \frac{d(y + \log y + c - k)}{G(y)} = \int_{\log m}^\infty \frac{1 + 1/y}{G(y)} dy \quad \text{for } m \geq N.$$

Since $k - \alpha - 2c + 1 > 10^{-8} > 0$,

$$H(y)(1 + 1/y) - G(y) = (k - \alpha - 2c + 1)y + (k - \alpha - 2c + 2)\log y \\ + (\log y - \alpha)(\log y - k)/y + \alpha k - \alpha - k - (c - k)^2 \\ > \log y + \alpha k - \alpha - k - (c - k)^2 > 1.4 > 0$$

for $y \geq \log m \geq \log N$; thus $h(m+1) < g(m+1)$ for $m \geq N$, i.e., $h(m) < g(m)$ for $m \geq N + 1$.

Lemma 3. If c is a constant such that $f(S) \leq c$ for any finite subsequence S of a given primitive sequence $A = \{a_i\}$, then $f(A) \leq c$.

Proof. Put $c_m = \sum_{1 \leq i \leq m} 1/(a_i \log a_i)$. Then $c_m \leq c$ and $c_m < c_{m+1}$ for $m \geq 1$; thus $f(A) = \lim_{m \rightarrow \infty} c_m \leq c$.

Lemma 4. We have $f(A_m) \leq 1/(\log m + \log \log m - k)$ for $m \geq N + 1$ for any given primitive sequence $A = \{a_i\}$.

Proof. By Lemma 3, we suppose A_m is finite. Induction on $d^\circ(A_m)$. If $d^\circ(A_m) \leq 1$, then

$$f(A_m) \leq \sum_{i \geq m} \frac{1}{p_i \log p_i} < \sum_{i \geq m} \frac{1}{p_i(\log i + \log \log i - k)} \\ < \frac{1}{\log m + \log \log m - k}$$

by Lemmas 1 and 2.

Now suppose $d^\circ(A_m) = s > 1$. Since $A_m = \bigcup_{i \geq m} A'_i$ is disjoint, $f(A_m) = \sum_{i \geq m} f(A'_i)$. If $d^\circ(A'_i) \leq 1$ then

$$f(A'_i) \leq \frac{1}{p_i \log p_i} < \frac{1}{p_i(\log i + \log \log i - k)}.$$

If $d^\circ(A'_i) > 1$ then

$$f(A'_i) < \frac{f(A''_i)}{p_i} < \frac{1}{p_i(\log i + \log \log i - k)}$$

since $d^\circ(A''_i) = d^\circ(A'_i) - 1 < s$. Thus by Lemma 2,

$$\begin{aligned} f(A_m) &= \sum_{i \geq m} f(A'_i) \leq \sum_{i \geq m} \frac{1}{p_i(\log i + \log \log i - k)} \\ &< \frac{1}{\log m + \log \log m - k}. \end{aligned}$$

Lemma 5. *If $f(A_{m+1}) \leq F_{m+1}$ for any primitive sequence A and*

$$F_m = \max\{F_{m+1}/(1 - 1/p_m), F_{m+1} + 1/(p_m \log p_m)\},$$

then $f(A_m) \leq F_m$ for any primitive sequence A .

Proof. Since $A_m = A_{m+1} \cup A'_m$ is disjoint,

$$f(A_m) = f(A_{m+1}) + f(A'_m) \leq F_{m+1} + f(A'_m).$$

By Lemma 3, we can suppose A_m is finite. Induction on $d^\circ(A_m)$. If $d^\circ(A_m) \leq 1$ or $d^\circ(A'_m) \leq 1$ then

$$f(A'_m) \leq 1/(p_m \log p_m)$$

and

$$f(A_m) \leq F_{m+1} + 1/(p_m \log p_m) \leq F_m.$$

Now suppose $d^\circ(A_m) = s > 1$ and $d^\circ(A'_m) > 1$. Then $d^\circ(A''_m) < s$; and thus

$$f(A''_m) \leq F_m \quad \text{and} \quad f(A'_m) < f(A''_m)/p_m \leq F_m/p_m.$$

Therefore,

$$f(A_m) \leq F_{m+1} + f(A'_m) \leq F_{m+1} + F_m/p_m \leq F_m(1 - 1/p_m) + F_m/p_m = F_m.$$

Theorem 1. *$f(A) < 1.84$ for any primitive sequence A .*

Proof. Put $F_{N+1} = 1/(\log(N+1) + \log \log(N+1) - k)$. Then

$$f(A_{N+1}) \leq F_{N+1}$$

by Lemma 4. Let

$$F_m = \max\{F_{m+1}/(1 - 1/p_m), F_{m+1} + 1/(p_m \log p_m)\}.$$

Then

$$f(A_m) \leq F_m \quad \text{for } m = N, N-1, \dots, 2, 1$$

by Lemma 5. Thus $f(A) = f(A_1) \leq F_1 < 1.84$. This completes the proof.

Theorem 2. *The necessary and sufficient condition for (2) to be true for any primitive sequence A is that $f(B) \leq \sum 1/(p \log p)$ for any primitive sequence B .*

Proof. (Necessity) If there exists a primitive sequence B such that $f(B) > \sum 1/(p \log p)$, then there exists an integer n such that

$$\sum_{b \in B, b \leq n} \frac{1}{b \log b} > \sum \frac{1}{p \log p},$$

thus

$$\sum_{b \in B, b \leq n} \frac{1}{b \log b} > \sum_{p \leq n} \frac{1}{p \log p};$$

this contradicts (2).

(Sufficiency) Suppose there exists a primitive sequence $A = \{a_i\}$ and an integer n such that

$$\sum_{a \in A, a \leq n} \frac{1}{a \log a} > \sum_{p \leq n} \frac{1}{p \log p}.$$

Let $B = \{a : a \in A, a \leq n\} \cup \{p : p > n\}$. Then B is primitive and $f(B) > \sum 1/(p \log p)$. This is a contradiction.

Remark. From Lemma 1 and the inequality

$$p_n \leq n(\log n + \log \log n - 0.5) \quad \text{for } n > 20 \text{ [6]}$$

we can get $1.63 < \sum 1/(p \log p) < 1.64$. From this fact and the above theorems we can see how far we are now from the complete proof of the conjecture.

REFERENCES

1. P. Erdős, *Note on sequences of integers no one of which is divisible by any other*, J. London Math. Soc. (2) **10** (1935), 126–128.
2. —, Seminar at the University of Limoges, 1988.
3. P. Erdős, A. Sarközi, and E. Szemerédi, *On divisibility properties of sequences of integers*, Colloq. Math. Soc. Janos Bolyai, vol. 2, Debrecen, Hungary, 1968.
4. H. Halberstam and K. F. Roth, *Sequences*, Chapter V, Oxford Univ. Press, 1966.
5. Guy Robin, *Estimation de la fonction de Tchebychef θ sur le k -ième nombre premier et grandes valeurs de la fonction $\omega(n)$ nombre de diviseurs premiers de n* , Acta Arith. **52** (1983), 367–389.
6. J. Barkley Rosser and L. Schoenfeld, *Approximate formulas for some functions of prime numbers*, Illinois J. Math. **6** (1962), 64–94.
7. Zhang Zhenxiang, *On a conjecture of Erdős on the sum $\sum_{p \leq n} 1/(p \log p)$* , J. Number Theory **39** (1991), 14–17.
8. —, *On a problem of Erdős concerning primitive sequences*, submitted.

MATHEMATICAL INSTITUTE, HUNGARIAN ACADEMY OF SCIENCES, BUDAPEST, HUNGARY

DEPARTMENT OF MATHEMATICS, ANHUI NORMAL UNIVERSITY, 241000 WUHU, ANHUI, PEOPLE'S REPUBLIC OF CHINA