A NOTE ON THE JACOBSON RADICAL

YU-LEE LEE

(Communicated by Maurice Auslander)

ABSTRACT. In this paper we solve a problem of Divinsky to represent the Jacobson radical as a lower radical class.

The purpose of this paper is to represent the Jacobson radical as a lower radical class.

For each class of rings M , there exists a radical class $U(M)$, called the upper radical class determined by M , such that $U(M)$ is the largest radical class with every ring in M $U(M)$ -semisimple [2, 5, 8]. For each class of rings L , there exists a radical class $Y(L)$, called the lower radical class determined by L , such that $Y(L)$ is the smallest radical class containing L [1, 3, 4, 6–8]. It is clear that every radical class is the lower radical class determined by itself; the Jacobson radical can be characterized as the upper radical of the class of all primitive rings [1, 7, 8]. Divinsky raised the problem of identifying the Jacobson radical as the lower radical of some reasonable class of rings [1, 8]. We have the following solution.

Theorem 1. *Let J be the Jacobson radical class. For each ring A , let $J(A)$ be the Jacobson radical of A and $J_2 = \{(J(A))^2 : A \text{ is a ring}\}$. Let N be the class of all nilpotent rings. Then the Jacobson radical class is the lower radical class determined by N and J_2 .*

Proof. It is clear that $Y(J_2 \cup N) \subseteq J$. Let $A \in J$, $I \not\subseteq A$ and A/I be any nonzero quotient ring of A . If $I \supseteq (J(A))^2$, then $A \supsetneq (J(A))^2$ and $A/I = (A/(J(A))^2)/(I/(J(A))^2)$. We have $(A/(J(A))^2)^2 = (J(A)/(J(A))^2)^2 = 0$. Hence $A/(J(A))^2 \in N$ and $0 \neq A/I \in N$.

If $I \not\supseteq (J(A))^2$ then $A/I \geq I + (J(A))^2/I \cong (J(A))^2/I \cap (J(A))^2 \neq 0$.

$$\begin{aligned} (J(A))^2/I \cap (J(A))^2 &= (J(A)/I \cap (J(A))^2)^2 \\ &= [J(A/I \cap (J(A))^2)]^2 \in J_2. \end{aligned}$$

Hence we have that $A \in Y(J_2 \cup N)$.

Let S be any radical class and $S_n = \{(S(A))^n : A \text{ is a ring}\}$. By the same proof, we have the following more general result.

Received by the editors March 29, 1991 and, in revised form, September 24, 1991.

1991 *Mathematics Subject Classification.* Primary 16N20, 16N80.

Key words and phrases. Jacobson radical, lower radicals.

Theorem 2. *If S is a radical class containing all nilpotent rings N , then S is the lower radical class determined by N and S_n for any positive integer n .*

We wish to give an example to show that $N \not\subseteq J_2$.

Example. Let A be the ring generated by $\{a\}$ with $2a = a^3 = 0$. Then A is a nilpotent ring. If $A \in J_2$ then there is a ring $B \in J$ with $A = B^2$ and B is nilpotent. Suppose that there exists b in B such that the ideal C generated by $\{A, b\}$ satisfies $C^2 = A$. Then $b^2 \neq a$; otherwise, $b^3 = ab \notin A$. Also, $ba \neq a$, $ba \neq a^2 + a$, $ab \neq a$, $ab \neq a^2 + a$, $ba^2 \neq a$, $ba^2 \neq a^2 + a$, $a^2b \neq a$, and $a^2b \neq a^2 + a$. For if $a = ba$ then $a = b^na = 0$. The other inequalities can be verified in a similar way. Hence $b^2 = a^2 + a$. Now $ba \neq 0$; otherwise, $0 = b^2a = a^3 + a^2 = a^2$. Hence $ba = a^2$ and $b^2 = a^2 + a = ba + a$, $a = b(b - a)$, and $a^2 = b(b - a)a = b(ba - a^2) = 0$. Hence $C^2 \not\subseteq A$ and $C^2 = \{0, a^2\}$.

Hence there are b and c in B such that the ideal D generated by $\{A, b, c\}$ satisfies $D^2 = A$, while the square of the ideal generated by A and b or c is $\{0, a^2\}$. If $bc = a$ then $ca = a^2$; otherwise, $ca = 0$ and $0 = bca = a^2$. However, $a^2 = bca = ba^2 = b^2a^2 = b^na^2 = 0$ and hence $bc = a + a^2$. Then $ca = a^2$; otherwise, $ca = 0$, $a^2 = (a + a^2)a = bca = 0$. However, $a^2 = (a + a^2)a = bca = ba^2 = b^na^2 = 0$; which is a contradiction. Similarly $cb \neq a$ and $cb \neq a^2 + a$. Hence $D^2 \subsetneq A$ and $B^2 \subsetneq A$ and A is not in J_2 . Therefore $N \not\subseteq J_2$ and $J_2 \neq J$.

This example also shows that $N \not\subseteq S_n$ and $S_n \neq S$ for any radical class S containing all nilpotent rings and $n \geq 2$. Because of the lack of a mechanism, it is tedious to verify a ring not in J_2 . It would be interesting to find more examples of rings in $J \setminus J_2$ and to establish general properties.

ACKNOWLEDGMENT

The author wishes to thank the referee for valuable suggestions.

REFERENCES

1. N. J. Divinsky, *Rings and radicals*, Univ. of Toronto Press, Ontario, 1965.
2. B. J. Gardner, *Radical theory*, Longman Scientific and Technical, England, 1989.
3. A. G. Kurosh, *Radicals of rings and algebra*, Mat. Sb. **33** (1953), 13–26; English transl., *Rings, modules, and radicals*, Colloq. Math. Soc. János Bolyai, vol. 6, North-Holland, Amsterdam, 1973, pp. 297–312.
4. W. G. Leavitt, *The general theory of radicals*, Univ. of Nebraska, Lincoln, NE, 1972.
5. Y. L. Lee, *On the construction of upper radical properties*, Proc. Amer. Math. Soc. **19** (1968), 1165–1166.
6. —, *On the construction of lower radical properties*, Pacific J. Math. **28** (1969), 393–395.
7. F. A. Szasz, *Radicals of rings*, Wiley, New York, 1981.
8. R. Wiegandt, *Radical and semisimple classes of rings*, Queen's Univ., Kingston, Ontario, Canada, 1974.