C^* -EXTREME POINTS OF SOME COMPACT C^* -CONVEX SETS

D. R. FARENICK AND PHILLIP B. MORENZ

(Communicated by Palle E. T. Jorgensen)

ABSTRACT. In the C^* -algebra M_n of complex $n \times n$ matrices, we consider the notion of noncommutative convexity called C^* -convexity and the corresponding notion of a C^* -extreme point. We prove that each irreducible element of M_n is a C^* -extreme point of the C^* -convex set it generates, and we classify the C^* -extreme points of any C^* -convex set generated by a compact set of normal matrices.

1. INTRODUCTION

For C^* -algebras and, more generally, for bimodules over C^* -algebras there is a notion of convexity that incorporates algebra-valued convex coefficients in a natural way. This form of convexity, called C^* -convexity, was studied in its own right by Loeb and Paulsen in [6] and by Hopenwasser, Moore, and Paulsen in [5]. There is an extremal theory associated with C^* -convexity and it is our aim in this paper to further its development by studying the C^* -extreme points of specific compact C^* -convex sets in the C^* -algebra M_n of complex $n \times n$ matrices.

Sections 3 and 4 contain the main results. In §3 it is proved that every irreducible element of M_n is a C^* -extreme point of the C^* -convex set it generates. In §4 conditions are presented that must be satisfied by the C^* -extreme points of a C^* -convex set generated by a family of matrices; these necessary conditions lead to a classification of the C^* -extreme points of compact C^* -convex sets generated by normals. Auxiliary results include an extension of Stampfli's theorem concerning the extreme points of the classical numerical range of a hyponormal operator and a result to the effect that irreducible elements $x \in M_n$ can be determined up to unitary equivalence among all elements of M_n by the norms of linear polynomials in x with coefficients taken from M_n .

We begin by recalling the definitions [6] of C^* -convexity and C^* -extreme point. A subset K of a unital C^* -algebra A is said to be C^* -convex if K is closed under the formation of finite sums of the type $\sum_i t_i^* x_i t_i$, where each

Received by the editors November 22, 1991; presented at GPOTS, College Station, TX, and at the Canadian Operator Symposium, Montreal, Québec, May 1991.

1991 *Mathematics Subject Classification.* Primary 15A60, 47A12, 47D20.

Key words and phrases. Matricial range, C^* -convex set, C^* -extreme point.

Work supported in part by NSERC of Canada and FCAR du Québec.

$x_i \in K$ and the elements $t_i \in A$ are under the proviso that $\sum_i t_i^* t_i = 1$. The t_i are called C^* -convex coefficients and the C^* -convex combination $\sum_i t_i^* x_i t_i$ is called proper if each coefficient t_i is invertible in A . An element x in a C^* -convex set $K \subset A$ is said to be a C^* -extreme point of K if the only possible representations of x as proper C^* -convex combinations of elements $x_i \in K$ are those for which each x_i comes from the unitary orbit of x . To distinguish extreme points from C^* -extreme points, the former are sometimes referred to as linear extreme points.

In [6] it is shown that the C^* -extreme points of C^* -convex subsets of M_n are linear extreme points, and in [5] an example in M_2 of a C^* -convex set having some linear extreme points that are not C^* -extreme is provided. In §4 we present a new class of examples, occurring in every matrix algebra, that show that linear extreme points need not be C^* -extreme. Although a compact C^* -convex subset of M_n always possesses C^* -extreme points [4], the Krein-Milman type problem of whether such extreme points exist in sufficient numbers to recover the original set remains open.

Complete information about the C^* -extreme points of certain C^* -convex sets is available in a few special cases: the closed unit ball of the algebra of operators acting on a Hilbert space, the closed interval $\{x : 0 \leq x \leq 1\}$ of operators, and the set of $x \in M_2$ with numerical radius $w(x) \leq 1$ (see [5]). We augment these results by completing the classification begun in [4] of the C^* -extreme points of C^* -convex sets generated by compact sets of normal matrices (Corollary 4.2) and by determining all of the C^* -extreme points of the C^* -convex set generated by a single 2×2 matrix (Theorem 4.3).

One source for interest in C^* -convexity is the theory of matricial ranges [1, §2.4; 3; 9, §4]. The n th matricial range of an element a in a unital C^* -algebra A is the compact C^* -convex subset $W^n(a) \subset M_n$ of elements of the form $\phi(a)$, where ϕ denotes a unital completely positive map of A into M_n . If x is an element of $B(H)$, the C^* -algebra of bounded linear operators on a complex Hilbert space H , then the numerical range $W^1(x)$ of x is the closure of the classical numerical range $\{(x\xi, \xi) : \xi \in H, \|\xi\| = 1\}$. In §5 some of our results on C^* -extreme points are applied to the matricial ranges of hyponormal operators.

The C^* -convex hull of a subset $\mathcal{S} \subset M_n$ is the smallest C^* -convex set containing $\mathcal{S}$ and is denoted by $C^*\text{-conv } \mathcal{S}$. An essential fact is that $C^*\text{-conv } \mathcal{S}$ is compact whenever $\mathcal{S}$ is a compact subset of M_n [3, 2.4].

The unitary equivalence of x and z is denoted by $x \sim z$ and is taken to mean that there is a unitary u such that $x = u^* z u$.

2. ON A COMPLETE UNITARY INVARIANT OF IRREDUCIBLE MATRICES

The (finite) order of an irreducible operator $x \in B(H)$ is the least positive integer k , if it exists, for which the values of $\|\sum_{i=0}^k x^i \otimes c_i\|$, where $c_0, c_1, \dots, c_k$ are complex $n \times n$ matrices with n arbitrary, determine x among the irreducible elements of $B(H)$ uniquely up to unitary equivalence. An important theorem of Arveson asserts that for irreducible compact operators acting on a complex separable Hilbert space, the order exists and is equal to 1 [1, Chapter 2]; in particular, irreducible elements in the C^* -algebra M_n are of first order. The purpose of this section is to show that for the algebra M_n the situation is

somewhat more special in that irreducible elements $x \in M_n$ can be determined up to unitary equivalence among all elements of M_n by the norms of linear matrix polynomials $x \otimes c_1 + 1 \otimes c_0$ in just one algebra only— $M_n \otimes M_n$.

2.1. Theorem. *Suppose that $x, z \in M_n$ satisfy $\|x \otimes b + 1 \otimes m\| = \|z \otimes b + 1 \otimes m\|$ for every $b, m \in M_n$. If x is irreducible, that is, if the only selfadjoint idempotents commuting with x are 0 and 1, then z must be irreducible and, moreover, unitarily equivalent to x .*

As in Arveson's paper [1], the proof of Theorem 2.1 involves the use of his boundary theorem and the comparison of the matricial ranges of z with those of x . It is in this latter aspect that the connection with C*-convexity theory arises.

Every unital completely positive map $\phi : M_n \rightarrow M_n$ has the form $\phi(\cdot) = \sum_i t_i^*(\cdot) t_i$, where the sum is a finite sum and the matrices t_i satisfy $\sum_i t_i^* t_i = 1$ [2]. Hence, the C*-convex set generated by $x \in M_n$ is precisely the n th matricial range of x , and so we are able to make use of some of the results of [3; 9] in a manner summarized by the

2.2. Theorem. *For $x, z \in M_n$, the following statements are equivalent.*

- (1) $C^*\text{-conv}\{x\} \subset C^*\text{-conv}\{z\}$;
- (2) $W^1(x \otimes b + 1 \otimes m) \subset W^1(z \otimes b + 1 \otimes m)$ for every $b, m \in M_n$;
- (3) $\|x \otimes b + 1 \otimes m\| \leq \|z \otimes b + 1 \otimes m\|$ for every $b, m \in M_n$.

Proof. In identifying $C^*\text{-conv}\{x\}$ with $W^n(x)$ and $C^*\text{-conv}\{z\}$ with $W^n(z)$, the implication (3) $\Rightarrow$ (1) is given by [3, 2.1] and the equivalence of (1) and (2) by [9, 4.3]. To prove the implication (1) $\Rightarrow$ (3), observe that because $x \in W^n(x) \subset W^n(z)$, there is a unital completely positive $\phi : M_n \rightarrow M_n$ with $x = \phi(z)$. The induced map $\phi \otimes \text{id}$ on $M_n \otimes M_n$, where $\text{id} : M_n \rightarrow M_n$ is the identity map, is a contraction, and therefore for each $b, m \in M_n$, $\|x \otimes b + 1 \otimes m\| = \|\phi(z) \otimes b + 1 \otimes m\| \leq \|z \otimes b + 1 \otimes m\|$. $\square$

We now provide the

Proof of Theorem 2.1. Using Theorem 2.2, the hypothesis is that $W^n(x) = W^n(z)$. There exist, therefore, unital completely positive maps $\phi, \psi : M_n \rightarrow M_n$ such that $x = \psi(z)$ and $z = \phi(x)$, whence $x = \psi \circ \phi(x)$. Let $\mathcal{L}$ denote the subspace spanned by x, x^* , and 1, and let ω denote the identity map $\mathcal{L} \rightarrow M_n$: namely, $\omega = \psi \circ \phi|_{\mathcal{L}}$. Because $\mathcal{L}$ is irreducible and the C*-algebra $\mathcal{L}$ generates is M_n , the boundary theorem of Arveson [1, 2.1.1] states that ω has a unique completely positive extension to M_n . The identity map and $\psi \circ \phi$, being two such extensions of ω , must therefore coincide.

From $\text{id} = \psi \circ \phi$, it follows that, for every $a \in M_n$,

$$\|a\| = \|\psi(\phi(a))\| \leq \|\phi(a)\| \leq \|a\|;$$

that is, ϕ is a unital isometry on M_n and, therefore, has the form $x \mapsto u^* x u$ or $x \mapsto u^* x^t u$ for some unitary $u \in M_n$ [7]. Complete positivity rules out the latter possibility, so that $z = \phi(x) = u^* x u$ as desired. $\square$

3. C*-CONVEX SETS WITH A SINGLE, IRREDUCIBLE GENERATOR

Using the main result of §2, we will prove that each irreducible element of M_n is C*-extreme in the C*-convex set it generates.

3.1. Theorem. *If $x \in M_n$ is irreducible, then x is a C^* -extreme point of its C^* -convex hull.*

Proof. Suppose that $x = \sum_i t_i^* x_i t_i$ for some elements $x_i \in C^*\text{-conv}\{x\}$ and for some invertible C^* -convex coefficients $t_i \in M_n$. Because each $x_i \in C^*\text{-conv}\{x\}$, it follows from (2) of Theorem 2.2 that $W^1(x_i \otimes b + 1 \otimes m) \subset W^1(x \otimes b + 1 \otimes m)$ for every i and for every $b, m \in M_n$. We now show that the reverse inclusions $W^1(x \otimes b + 1 \otimes m) \subset W^1(x_i \otimes b + 1 \otimes m)$ hold.

Fix $b, m \in M_n$. Let λ be an arbitrary extreme point of $W^1(x \otimes b + 1 \otimes m)$; so λ is determined by some unit vector $\xi \in \mathbb{C}^n \otimes \mathbb{C}^n$. By using the unit vectors

$$\eta_i = \|(t_i \otimes 1)\xi\|^{-1}(t_i \otimes 1)\xi$$

and the induced elements

$$\lambda_i = ((x_i \otimes b + 1 \otimes m)\eta_i, \eta_i),$$

the extreme point λ can be expressed as a (proper) convex combination of the $\lambda_i \in W^1(x_i \otimes b + 1 \otimes m)$ as follows:

$$\begin{aligned} \lambda = ((x \otimes b + 1 \otimes m)\xi, \xi) &= \sum_i ((x_i \otimes b + 1 \otimes m)(t_i \otimes 1)\xi, (t_i \otimes 1)\xi) \\ &= \sum_i \|(t_i \otimes 1)\xi\|^2 \lambda_i. \end{aligned}$$

But each λ_i is an element of $W^1(x \otimes b + 1 \otimes m)$ and therefore, because λ is an extreme point of $W^1(x \otimes b + 1 \otimes m)$, we must have $\lambda_i = \lambda$ for every i . This proves that the extreme points of $W^1(x \otimes b + 1 \otimes m)$ lie in each of the sets $W^1(x_i \otimes b + 1 \otimes m)$. Hence, $W^1(x \otimes b + 1 \otimes m)$ is contained in each $W^1(x_i \otimes b + 1 \otimes m)$ and so, by Theorem 2.2, $C^*\text{-conv}\{x_i\} = C^*\text{-conv}\{x\}$ for every i . Equivalently, $\|x_i \otimes b + 1 \otimes m\| = \|x \otimes b + 1 \otimes m\|$ for every $b, m \in M_n$ and every i . But x is irreducible, and so as Theorem 2.1 asserts, each x_i is unitarily equivalent to x ; that is, x is a C^* -extreme point of its C^* -convex hull. $\square$

4. THE SEARCH FOR C^* -EXTREME POINTS; NARROWING DOWN THE POSSIBILITIES

Before we can hope to answer the Krein-Milman question of whether the C^* -extreme points are sufficient to recover the original C^* -convex set, it would help to be able to describe all the C^* -extreme points. The purpose of this section is to present necessary conditions for points to be C^* -extreme in various cases in order to narrow the search considerably. The main results are Theorem 4.1 and its applications.

We begin by presenting techniques for rewriting C^* -convex combinations.

Technique A. Let $x = \sum_{i=1}^m t_i^* x_i t_i$ with $\sum_i t_i^* t_i = 1$ and $\|t_i\| < 1$ for each i . If this is not already a proper C^* -convex combination, then we can rewrite it as one in the following manner. Let

$$a_i = (m-1)^{-1/2}(1 - t_i^* t_i)^{1/2}$$

(invertible since $\|t_i\| < 1$),

$$\kappa_{ij} = \frac{1 - \delta_{ij}}{m-1}, \quad h_{ij} = \sqrt{k_{ij}} t_{ij} a_i^{-1}.$$

For each i ,

$$\begin{aligned}\sum_j h_{ij}^* h_{ij} &= \sum_j \sqrt{\kappa_{ij}} (a_i^{-1})^* t_j^* t a_i^{-1} \sqrt{\kappa_{ij}} \\ &= (1 - t_i^* t_i)^{-1/2} \left(\sum_j (1 - \delta_{ij}) t_j^* t_j \right) (1 - t_i^* t_i)^{-1/2} \\ &= (1 - t_i^* t_i)^{-1/2} (1 - t_i^* t_i) (1 - t_i^* t_i)^{-1/2} = 1,\end{aligned}$$

and so $z_i = \sum_{j=1}^m h_{ij}^* x_j h_{ij} \in C^*\text{-conv}\{x_1, \dots, x_m\}$ by the above computation. Further, $x = \sum_i a_i^* z_i a_i$ is a representation of x as a proper C^* -convex combination of the z_i . An important subtlety that will be essential later is that each $h_{ii} = 0$ and so each z_i is a C^* -convex combination of the x_j using fewer than m nonzero coefficients.

Technique B. If $p_1^* p_1 + p_2^* p_2 = 1$ and if

$$x = \begin{pmatrix} 1 & 0 \\ 0 & p_1^* \end{pmatrix} x_1 \begin{pmatrix} 1 & 0 \\ 0 & p_1 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & p_2^* \end{pmatrix} x_2 \begin{pmatrix} 0 & 0 \\ 0 & p_2 \end{pmatrix},$$

then for every scalar $\lambda \in (0, 1)$, x can be rewritten as

$$\begin{aligned}x &= \begin{pmatrix} \lambda 1 & 0 \\ 0 & \lambda p_1^* \end{pmatrix} x_1 \begin{pmatrix} \lambda 1 & 0 \\ 0 & \lambda p_1 \end{pmatrix} \\ &\quad + \begin{pmatrix} \sqrt{1 - \lambda^2} 1 & 0 \\ 0 & (1 - \lambda^2 p_1^* p_1)^{1/2} \end{pmatrix} x' \begin{pmatrix} \sqrt{1 - \lambda^2} 1 & 0 \\ 0 & (1 - \lambda^2 p_1^* p_1)^{1/2} \end{pmatrix},\end{aligned}$$

where

$$\begin{aligned}x' &= \begin{pmatrix} 1 & 0 \\ 0 & \sqrt{1 - \lambda^2} (1 - \lambda^2 p_1^* p_1)^{-1/2} p_1^* \end{pmatrix} x_1 \begin{pmatrix} 1 & 0 \\ 0 & p_1 \sqrt{1 - \lambda^2} (1 - \lambda^2 p_1^* p_1)^{-1/2} \end{pmatrix} \\ &\quad + \begin{pmatrix} 0 & 0 \\ 0 & (1 - \lambda^2 p_1^* p_1)^{-1/2} p_2^* \end{pmatrix} x_2 \begin{pmatrix} 0 & 0 \\ 0 & p_2 (1 - \lambda^2 p_1^* p_1)^{-1/2} \end{pmatrix}.\end{aligned}$$

Observe that in order for x to be a proper C^* -convex combination of x_1 and x' , it is sufficient that p_1 be invertible.

Technique C. If $\sum_{i=1} p_i^* p_i$ is invertible, then

$$\sum_i \begin{pmatrix} 0 & 0 \\ 0 & p_i^* \end{pmatrix} x_i \begin{pmatrix} 0 & 0 \\ 0 & p_i \end{pmatrix}$$

can be rewritten as

$$\begin{pmatrix} 0 & 0 \\ 0 & (\sum_j p_j^* p_j)^{1/2} \end{pmatrix} z \begin{pmatrix} 0 & 0 \\ 0 & (\sum_j p_j^* p_j)^{1/2} \end{pmatrix},$$

where $z \in C^*\text{-conv}\{x_i\}_i$ and is given by

$$z = \sum_i \begin{pmatrix} m^{-1/2} 1 & 0 \\ 0 & (\sum_j p_j^* p_j)^{-1/2} p_i^* \end{pmatrix} x_i \begin{pmatrix} m^{-1/2} 1 & 0 \\ 0 & p_i (\sum_j p_j^* p_j)^{-1/2} \end{pmatrix}.$$

Finally, we will make use of the following two elementary facts. One is that if $K \subset M_n$ is C^* -convex, and if $p \in M_n$ is a projection, then the compression pKp of K to the image of p is clearly C^* -convex. The other is that $x = x_1 \oplus x_2$

is C^* -extreme in K only if x_1 and x_2 are C^* -extreme in the corresponding compressed images of K . To prove the second fact, suppose that x_1 , say, is not C^* -extreme in K_1 , the corresponding compression of K . Then x_1 can be expressed as a proper C^* -convex combination $\sum_{i=1}^m s_i^* z_i s_i$ of elements $z_i \in K_1$, with at least one $z_i \not\sim x_1$. Thus,

$$x = \sum_{i=1}^m (s_i \oplus m^{-1/2} 1)^* (z_i \oplus x_2) (s_i \oplus m^{-1/2} 1)$$

is a representation of x as a proper C^* -convex combination of elements $z_i \oplus x_2 \in K$, where at least one $z_i \oplus x_2 \not\sim x_1 \oplus x_2$.

4.1. Theorem. Suppose that $K = C^*\text{-conv}\{x_\alpha : \alpha \in I\} \subset M_n$, where I is any index set. If x is C^* -extreme in K , then x must be unitary equivalent to some x_α or reducible. Moreover, there exist projections q_i such that $\sum_i q_i = 1$, $x = \sum_i q_i x'_{\alpha_i} q_i$, and each $x'_{\alpha_i} \sim x_{\alpha_i}$.

Proof. Let $x = \sum_{i=1}^m t_i^* x_{\alpha_i} t_i$ be a representation of x as a C^* -convex combination of the x_α using the fewest number of coefficients possible. (This least integer m does not change if x is replaced by some $x' \sim x$.) Because $m = 1$ if and only if $x \sim x_{\alpha_1}$, it is assumed henceforth that $m \geq 2$ and that x is not unitarily equivalent to some x_α . We are to prove that x is reducible.

We will assume without loss of generality that $\|t_1\| = 1$. Such an assumption is valid for the following reasons. If, on the contrary, $\|t_i\| < 1$ for every i , then x can be rewritten using Technique A as a proper C^* -convex combination of elements z_i . Because x is a C^* -extreme point of K , each z_i must be unitarily equivalent to x . However, the construction in Technique A shows that each z_i is a C^* -convex combination of fewer than m of the x_α ; hence, x must possess this property of the z_i as well, in contradiction of the minimality of m . Thus, at least one coefficient t_i has unit norm.

By expressing each coefficient t_i in its polar decomposition $t_i = u_i a_i$, where u_i is unitary and $a_i \geq 0$, and by absorbing the unitary part u_i of t_i into x_{α_i} , we have that $x = \sum_i a_i x'_{\alpha_i} a_i$, where $a_i \geq 0$, $\sum_i a_i^2 = 1$, and $x'_{\alpha_i} = u_i^* x_{\alpha_i} u_i$. There is a unitary $u \in M_n$ for which $u^* a_1 u$ has a representation as a 2×2 operator matrix given by

$$u^* a_1 u = \begin{pmatrix} 1 & 0 \\ 0 & y_1 \end{pmatrix}$$

with $y_1 \geq 0$ and $\|y_1\| < 1$. (This follows from the fact that $a_1 \geq 0$ and $\|a_1\| = 1$.) Because $a_i \geq 0$ for all $i \geq 2$ and because $\sum_i a_i^2 = 1$, the same unitary u gives

$$u^* a_i u = \begin{pmatrix} 0 & 0 \\ 0 & y_i \end{pmatrix}$$

for $i \geq 2$.

Again, each $y_i \geq 0$. Hence,

$$x' = \begin{pmatrix} 1 & 0 \\ 0 & y_1 \end{pmatrix} x''_{\alpha_1} \begin{pmatrix} 1 & 0 \\ 0 & y_1 \end{pmatrix} + \sum_{i \geq 2} \begin{pmatrix} 0 & 0 \\ 0 & y_i \end{pmatrix} x''_{\alpha_i} \begin{pmatrix} 0 & 0 \\ 0 & y_i \end{pmatrix},$$

where $x' = uxu^*$ and $x''_{\alpha_i} = u^*x'_{\alpha_i}u$; for simplicity of notation, let us assume that

$$a_1 = \begin{pmatrix} 1 & 0 \\ 0 & y_1 \end{pmatrix}, \quad a_i = \begin{pmatrix} 0 & 0 \\ 0 & y_i \end{pmatrix},$$

et cetera, so that $x = \sum_i a_i x_{\alpha_i} a_i$.

Now because $\|y_1\| < 1$, $1 - y_1^2$ is invertible; therefore, from

$$\sum_{i \geq 2} a_i^2 = \sum_{i \geq 2} \begin{pmatrix} 0 & 0 \\ 0 & y_i^2 \end{pmatrix} = 1 - a_1^2 = \begin{pmatrix} 0 & 0 \\ 0 & 1 - y_1^2 \end{pmatrix},$$

we conclude that $\sum_{i \geq 2} y_i^* y_i$ is invertible. Hence, Technique C allows us to write $\sum_{i \geq 2} a_i x_{\alpha_i} a_i$ as a single term $t_0^* x_0 t_0$, for some $x_0 \in K$. By passing to the polar decomposition of t_0 and absorbing the unitary part of t_0 into x_0 , we may assume that $\sum_{i \geq 2} a_i x_{\alpha_i} a_i = a_0 x_0 a_0$ for some $a_0 \geq 0$ and $x_0 \in K$.

From $a_0^2 + a_1^2 = 1$, we see that

$$a_0 = \begin{pmatrix} 0 & 0 \\ 0 & y_0 \end{pmatrix}.$$

In fact, a_0 and a_1 admit a further decomposition so that x is given by

$$x = \begin{pmatrix} 1 & & \\ & p_1 & \\ & & 0 \end{pmatrix} x_{\alpha_1} \begin{pmatrix} 1 & & \\ & p_1 & \\ & & 0 \end{pmatrix} + \begin{pmatrix} 0 & & \\ & p_0 & \\ & & 1 \end{pmatrix} x_0 \begin{pmatrix} 0 & & \\ & p_0 & \\ & & 1 \end{pmatrix},$$

where $p_1 \geq 0$ is invertible and is of norm $\|p_1\| < 1$. We are now in a position to make repeated use of Technique B.

For each $\lambda \in (0, 1)$, we rewrite x (using Technique B) as

$$x = \begin{pmatrix} \lambda 1 & & \\ & \lambda p_1 & \\ & & 0 \end{pmatrix} x_{\alpha_1} \begin{pmatrix} \lambda 1 & & \\ & \lambda p_1 & \\ & & 0 \end{pmatrix} + \begin{pmatrix} \sqrt{1-\lambda^2} 1 & & \\ & (1-\lambda^2 p_1^2)^{1/2} & \\ & & 1 \end{pmatrix} x(\lambda) \begin{pmatrix} \sqrt{1-\lambda^2} 1 & & \\ & (1-\lambda^2 p_1^2)^{1/2} & \\ & & 1 \end{pmatrix},$$

where

$$x(\lambda) = \begin{pmatrix} 1 & & \\ & \sqrt{1-\lambda^2} (1-\lambda^2 p_1^2)^{-1/2} p_1 & \\ & & 0 \end{pmatrix} x_{\alpha_1} \begin{pmatrix} 1 & & \\ & \sqrt{1-\lambda^2} p_1 (1-\lambda^2 p_1^2)^{1/2} & \\ & & 0 \end{pmatrix} + \begin{pmatrix} 0 & & \\ & (1-\lambda^2 p_1^2)^{-1/2} p_0 & \\ & & 1 \end{pmatrix} x_0 \begin{pmatrix} 0 & & \\ & p_0 (1-\lambda^2 p_1^2)^{-1/2} & \\ & & 1 \end{pmatrix}.$$

In the expression for x , the (matrix) coefficients on $x(\lambda)$ are invertible for all $\lambda \in (0, 1)$. By grouping together the 2×2 upper left corner blocks of the coefficients in the expression for x , that is,

$$x = \begin{pmatrix} a(\lambda) & 0 \\ 0 & 0 \end{pmatrix} x_{\alpha_1} \begin{pmatrix} a(\lambda) & 0 \\ 0 & 0 \end{pmatrix} + \begin{pmatrix} b(\lambda) & 0 \\ 0 & 1 \end{pmatrix} x(\lambda) \begin{pmatrix} b(\lambda) & 0 \\ 0 & 1 \end{pmatrix},$$

we see that Technique B applies once again (though cosmetically modified since 1 appears in the bottom right corner rather than in the top left) to yield x as a proper C^* -convex combination

$$x = t(\lambda)^* x'' t(\lambda) + s(\lambda)^* x(\lambda) s(\lambda).$$

(This is a proper combination because $b(\lambda)$ is invertible for all $\lambda \in (0, 1)$.) Because x is C^* -extreme in K , $x(\lambda) \sim x$ for all $\lambda \in (0, 1)$. Moreover, because $x(\lambda)$ depends continuously on λ and because the unitary orbit of x is closed, $x(1) \sim x$. This proves that x is reducible, for $x(1)$ is given by

$$x(1) = \begin{pmatrix} 1 & & \\ & 0 & \\ & & 0 \end{pmatrix} x_{\alpha_1} \begin{pmatrix} 1 & & \\ & 0 & \\ & & 0 \end{pmatrix} + \begin{pmatrix} 0 & & \\ & 1 & \\ & & 1 \end{pmatrix} x_0 \begin{pmatrix} 0 & & \\ & 1 & \\ & & 1 \end{pmatrix},$$

or equivalently, $x \sim x(1) = q_1 x_{\alpha_1} q_1 + q_1^\perp x_0 q_1^\perp$, where q_1 is a projection.

Recall that Technique C has x_0 as a C^* -convex combination of $x_{\alpha_2}, \dots, x_{\alpha_m}$ and so

$$x(1) = q_1 x_{\alpha_1} q_1 + \sum_{i \geq 2} s_i^* x_{\alpha_i} s_i,$$

where $q_1 + \sum_{i \geq 2} s_i^* s_i = 1$ and $q_1 \perp s_i$ for all $i \geq 2$. Observe that $\sum_{i \geq 2} s_i^* s_i = 1|_{\text{Im } q_1^\perp}$ and that the restriction of $x(1)$ to $\text{Im } q_1^\perp$ is C^* -extreme in $q_1^\perp K q_1^\perp$ (by the second of our two elementary facts stated prior to the theorem); hence, we may repeat our earlier arguments to conclude that one of the s_i is of norm 1, say s_2 , and to conclude further that there exists a projection q_2 such that

$$x \sim q_1 x_{\alpha_1} q_1 + q_2 x_{\alpha_2} q_2 + (\text{other terms}).$$

By exhausting this process after a finite number of steps, we see that $x \sim \sum_i q_i x_{\alpha_i} q_i$ for some projections q_i satisfying $\sum_i q_i = 1$. $\square$

4.2. Corollary. *If $\mathcal{S} \subset M_n$ is a compact set of normals, then $K = C^*\text{-conv } \mathcal{S}$ has no nonnormal C^* -extreme points. Moreover, if $X \subset \mathbb{C}$ is the convex hull of the eigenvalues of the elements of $\mathcal{S}$, then $x \in K$ is C^* -extreme in K if and only if x is normal and the eigenvalues of x are extreme points of X .*

Proof. It is easy to see (e.g., [4]) that $K = C^*\text{-conv}\{\zeta 1 : \zeta \text{ is an extreme point of } X\}$. If $x \in K$ is C^* -extreme in K , then Theorem 4.1 shows that x is a scalar matrix $\zeta 1$, or is reducible and of the form $x = \sum_i q_i (\zeta_i 1) q_i$ for some projections q_i satisfying $\sum_i q_i = 1$. In either case, x is normal and the eigenvalues of x are extreme points of X . Conversely, every normal $x \in M_n$ with eigenvalues in the set of extreme points of X is C^* -extreme in K [4, Theorem 2]. $\square$

In [4, Theorem 3] it was shown that the nilpotent Jordan matrix of order n is an extreme point of $K = C^*\text{-conv}\{\zeta 1_n : \zeta^{n+1} = 1\} \subset M_n$. Therefore, we now have a new set of examples, valid in each matrix algebra M_n , of C^* -convex sets having linear extreme points that are not C^* -extreme. (Previously, the only known example was in M_2 [5].)

The methods of this paper also allow us to recover all of the C^* -extreme points found by Hopenwasser, Moore, and Paulsen of the C^* -convex set in M_2 of matrices with numerical radius not exceeding 1. As is demonstrated by Arveson [1, p. 302], this set is generated by $\begin{pmatrix} 0 & 2 \\ 0 & 0 \end{pmatrix}$. In fact, we have the following general result.

4.3. Theorem. *If $x \in M_2$, then the C^* -extreme points of $C^*\text{-conv}\{x\}$ are all of the matrices unitarily equivalent to x and all of the scalar matrices obtained from the extreme points of the numerical range of x .*

Proof. If $x \in M_2$ is normal, then Corollary 4.2 yields the result. If $x \in M_2$ is nonnormal, then x is irreducible and so elements of the unitary orbit of x are C^* -extreme in $C^*\text{-conv}\{x\}$ (by Theorem 3.1), as are all scalar matrices $\zeta 1$ for which ζ is an extreme point of $W^1(x)$ [4, Theorem 1]. Conversely, suppose that z is C^* -extreme in $C^*\text{-conv}\{x\}$. If z is irreducible, then $z \sim x$; this follows from Theorem 4.1. If z is reducible, then it must be unitarily equivalent to a diagonal matrix, so let us say that

$$z = \begin{pmatrix} \alpha & 0 \\ 0 & \beta \end{pmatrix}.$$

If $\alpha \neq \beta$, then there exist linearly independent unit vectors $\xi, \eta \in \mathbb{C}^2$ such that $(z\xi, \xi) = \alpha$ and $(z\eta, \eta) = \beta$. The matrices

$$t_1 = \sqrt{2}^{-1}[\xi, -\eta] \in M_2, \quad t_2 = \sqrt{2}^{-1}[\xi, \eta] \in M_2$$

are invertible and satisfy $t_1^*t_1 + t_2^*t_2 = 1$ and $t_1^*xt_1 + t_2^*xt_2 = z$. But because z is C^* -extreme in $C^*\text{-conv}\{x\}$, the (reducible) matrix z must be unitarily equivalent to the (irreducible) matrix x , thereby yielding a contradiction. Thus, z must be of the form $\begin{pmatrix} \alpha & 0 \\ 0 & \alpha \end{pmatrix}$. Plainly, in order for the scalar matrix $\alpha 1_2$ to be extreme, let alone C^* -extreme, in $W^2(x)$, it is necessary that α be extreme in $W^1(x)$. $\square$

5. MATRICIAL RANGES OF A HYPONORMAL OPERATOR

In their announcement [8], Percy and Salinas define the $n \times n$ reducing matricial spectrum of $x \in B(H)$ to be the (possibly void) set $R^n(x)$ of $n \times n$ matrices of the form $\rho(x)$, where $\rho : C^*(x) \rightarrow M_n$ is a unital *-homomorphism and $C^*(x)$ is the unital C^* -algebra generated by x . By [8, Theorem 2], $\lambda \in R^1(x)$ if and only if there exists a sequence of unit vectors $\xi_j \in H$ satisfying $\lim_j \|(x - \lambda 1)\xi_j\| = \lim_j \|(x - \lambda 1)^*\xi_j\| = 0$. Therefore, if x is hyponormal, meaning that $x^*x - xx^* \geq 0$, then $R^1(x)$ is precisely the approximate point spectrum $\sigma_l(x)$ of x , and it is $\sigma(x)$ if x is in fact normal. Thus, the well-known result $W^1(x) = \text{conv } \sigma(x)$ for hyponormal x leads to $W^1(x) = \text{conv } R^1(x)$ (by making use of the fact that the extreme points of $\text{conv } \sigma(x)$ must be approximate eigenvalues). An interesting theorem of Stampfli goes even further: if $\lambda = (x\xi, \xi)$ is an extreme point of $W^1(x)$ for some unit vector $\xi \in H$, then $x\xi = \lambda\xi$ and $x^*\xi = \lambda^*\xi$ [10].

The purpose of this section is to show that these numerical range properties of hyponormal operators are present at the level of matricial ranges.

5.1. Theorem. *The following statements hold for every hyponormal operator x acting on a Hilbert space H .*

- (1) $W^n(x) = C^*\text{-conv } R^n(x)$.
- (2) The C^* -extreme points of $W^n(x)$ are elements of $R^n(x)$.
- (3) If $\Lambda = v^*xv$ is a C^* -extreme point of $W^n(x)$ for some isometry $v : \mathbb{C}^n \rightarrow H$, then vv^* is an invariant projection of x and x^* .

Proof. (1) It is slightly more convenient to translate x by a scalar ζ so that the new hyponormal operator $\tilde{x} = x - \zeta 1$ has $0 \in \sigma_l(\tilde{x})$. Because the effect of translation on the matricial spectrum and the matricial range is just translation by the scalar matrix $\zeta 1_n$, we assume without loss of generality that already $0 \in \sigma_l(x)$. Because x is hyponormal, the dilation theorem [11] of Sz.-Nagy and Foiaş asserts that there exists a normal operator z on a Hilbert space H' and a contraction $a : H \rightarrow H'$ such that $x = a^*za$ and $\sigma(z) \subset \sigma_l(x)$. The spectral inclusion as stated is in fact $R^1(z) \subset R^1(x)$, due to the normality of z and the hyponormality of x . If $R^1(z)$ does not contain 0, we can pass to the normal operator $z \oplus 0$ on $H' \oplus \mathbb{C}$, having 0 in its spectrum, so that x is now of the form $a_0^*(z \oplus 0)a_0$, where a_0 is the contraction obtained by the composition $a_0 = pqa$ with q denoting the canonical injection $H' \rightarrow H' \oplus \mathbb{C}$ and p denoting the canonical projection $H' \oplus \mathbb{C} \rightarrow H'$. Hence, again without loss of generality, we will assume that already $0 \in R^1(z)$.

We first show that $W^n(x) \subset W^n(z)$. Because $0 \in (R^1(z) \otimes 1_n)$, it follows that $\lambda^*W^n(z)\lambda \subset W^n(z)$ for every contraction $\lambda \in M_n$, by Remark 7 of [6]. Suppose now that ϕ is a unital completely positive map $C^*(x) \rightarrow M_n$; we will show that $\phi(x) \in W^n(z)$. The argument runs similar to that of Lemma 4.2 in [9]. Consider the completely positive map $\psi : C^*(z) \rightarrow M_n$ defined by $\psi(f) = \phi(a^*fa)$, with $\lambda = \psi(1)$. For the positive contraction λ there exists $b \geq 0$ in M_n such that if p is the projection of $\mathbb{C}^n$ onto the range of λ , then $b\lambda b = p$ (we have $b = \lambda^{-1/2}$ on the range of p). Choose any state θ on $C^*(z)$ and consider the unital completely positive map $\omega : C^*(z) \rightarrow M_n$ defined by $\omega(f) = b\psi(f)b + \theta(f)(1 - p)$. From $\lambda^{1/2}\omega(z)\lambda^{1/2} = \phi(x)$ it follows that

$$\phi(x) \in \lambda^{1/2}W^n(z)\lambda^{1/2} \subset W^n(z),$$

as desired.

Because the sets $R^n(x)$ and $R^n(z)$ consist solely of normal matrices, their C^* -convex hulls are determined by their eigenvalue sets; that is, $R^1(x)$ and $R^1(z)$ determine the C^* -convex hulls of $R^n(x)$ and $R^n(z)$, respectively. Hence,

$$W^n(z) = C^*\text{-conv } R^1(z) \subset C^*\text{-conv } R^1(x) \subset W^n(x) \subset W^n(z).$$

(2) By (1), $W^n(x)$ is the C^* -convex hull of a compact set of normals; hence, if $\Lambda \in W^n(x)$ is a C^* -extreme point of $W^n(x)$, then Corollary 4.2 has that Λ must be normal and its eigenvalues must be extreme in the convex hull of the spectra of $R^n(x)$ (viz., the eigenvalues of Λ are extreme points of $\text{conv } R^1(x) = W^1(x)$). Each eigenvalue of Λ determines a unital $*$ -homomorphism $C^*(x) \rightarrow \mathbb{C}$ and by passing to a direct sum of these homomorphisms, we obtain a unital $*$ -homomorphism $\rho : C^*(x) \rightarrow M_n$ such that $\rho(x)$ is unitarily equivalent to Λ .

(3) By (2), the C^* -extreme point $\Lambda = v^*xv$ is normal and $\sigma(\Lambda)$ is a subset of the extreme points of $W^1(x)$. Thus, whenever ξ is a unit eigenvector of Λ with corresponding eigenvalue λ , then $\lambda = (xv\xi, v\xi)$ is an extreme point of $W^1(x)$. By Stampfli's theorem, $x(v\xi) = \lambda(v\xi)$ and $x^*(v\xi) = \lambda^*(v\xi)$. Furthermore, as Λ is completely determined by its spectral structure, we conclude that $xv = v\Lambda$ and $x^*v = v\Lambda^*$, whence vv^* commutes with x . $\square$

REFERENCES

1. W. B. Arveson, *Subalgebras of C*-algebras. II*, Acta Math. **128** (1972), 271–308.
2. M.-D. Choi, *Completely positive linear maps on complex matrices*, Linear Algebra Appl. **10** (1975), 285–290.
3. D. R. Farenick, *C*-convexity and matricial ranges*, Canad. J. Math. (to appear).
4. ———, *Krein-Milman type problems for compact matricially convex sets*, Linear Algebra Appl. **162–164** (1992), 325–334.
5. A. Hopenwasser, R. L. Moore, and V. I. Paulsen, *C*-extreme points*, Trans. Amer. Math. Soc. **266** (1981), 291–307.
6. R. I. Loebl and V. I. Paulsen, *Some remarks on C*-convexity*, Linear Algebra Appl. **35** (1981), 63–78.
7. K. Morita, *Analytical characterization of displacements in general Poincaré space*, Proc. Imperial Acad. Tokyo **17** (1941), 489–494.
8. C. Pearcy and N. Salinas, *Finite-dimensional representations of separable C*-algebras*, Bull. Amer. Math. Soc. **80** (1974), 970–972.
9. R. R. Smith and J. D. Ward, *The geometric structure of generalized state spaces*, J. Funct. Anal. **40** (1981), 170–184.
10. J. G. Stampfli, *Extreme points of the numerical range of a hyponormal operator*, Michigan Math. J. **13** (1966), 87–89.
11. B. Sz.-Nagy and C. Foiaş, *An application of dilation theory to hyponormal operators*, Acta Sci. Math. (Szeged) **37** (1975), 155–159.

CENTRE DE RECHERCHES MATHÉMATIQUES, UNIVERSITÉ DE MONTRÉAL, C.P. 6128-A, MONTRÉAL, QUÉBEC, CANADA H3C 3J7

Current address, D. R. Farenick: Department of Mathematics and Statistics, University of Regina, Regina, Saskatchewan, Canada S4S 0A2

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO, TORONTO, ONTARIO, CANADA M5S 1A1