LOGARITHMIC CONVEXITY OF PERRON-FROBENIUS EIGENVECTORS OF POSITIVE MATRICES

SIDDHARTHA SAHI

(Communicated by Louis J. Ratliff, Jr.)

ABSTRACT. Let $C(S)$ be the cone of Perron-Frobenius eigenvectors of stochastic matrices that dominate a fixed substochastic matrix S . For each $0 \leq \alpha \leq 1$, it is shown that if u and v are in $C(S)$ then so is w , where $w_j = u_j^\alpha v_j^{1-\alpha}$.

The basic result of Perron-Frobenius theory [S] is that if a matrix has strictly positive entries, then its maximal eigenvalue is unique, positive, occurs with multiplicity 1, and has a (coordinatewise) positive eigenvector.

Subsequent literature on positive matrices contains many results (e.g., [C, F, K]) that deal with convexity properties of the dominant *eigenvalue* as a function of matrix entries. Similar results for the corresponding *eigenvectors* are obtained in [DN, EJN] but *only* for the effects of varying a *single* row of the matrix. Little seems to be known about the behavior of these eigenvectors under a more general perturbation of the matrix.

In this paper we prove a different kind of convexity property for Perron-Frobenius eigenvectors that was motivated by economic considerations in [SY] but that, by virtue of its unexpected and elementary nature, seems to warrant a wider mathematical audience.

For convenience, we formulate the result in terms of stochastic matrices—a positive matrix is called *stochastic* (*substochastic*) if its column sums are equal to (less than) 1.

For a fixed substochastic matrix S , consider the cone $C(S)$ of all (positive) Perron-Frobenius eigenvectors of the various stochastic matrices that (entrywise) dominate S . Thus $C(S) = \{v > 0 \mid \exists \text{ stochastic } A \geq S \text{ such that } Av = v\}$.

Now $C(S)$ need not be a convex subset of $\mathbf{R}^n$. However, we shall show that it has the following remarkable property that may be termed logarithmic, or geometric, convexity.

Theorem. Fix $0 \leq \alpha \leq 1$, and put $\beta = 1 - \alpha$. If $u = (u_1, u_2, \dots, u_n)$ and $v = (v_1, v_2, \dots, v_n)$ are in $C(S)$ then so is $w = (w_1, w_2, \dots, w_n)$ where $w_j = u_j^\alpha v_j^\beta$.

Received by the editors November 13, 1991 and, in revised form, December 13, 1991.

1991 *Mathematics Subject Classification*. Primary 15A48, 15A51.

Key words and phrases. Positive matrix, stochastic matrix, Perron-Frobenius eigenvector, convexity.

The principal difficulty in proving this theorem is the indirect nature of the definition of $C(S)$. The following lemma “eliminates the quantifier” in that definition.

Lemma. *A positive vector v belongs to $C(S)$ if and only if $Sv \leq v$.*

Proof. If v is in $C(S)$, choose $A \geq S$ such that $v = Av$. Then clearly $Sv \leq Av = v$.

Conversely, suppose $v > 0$ with $Sv \leq v$, and put $\delta = v - Sv$. Also, let s_j be the j th column sum of S , and put $\varepsilon_j = 1 - s_j$. Clearly, $0 < \varepsilon_1 v_1 + \cdots + \varepsilon_n v_n = \delta_1 + \cdots + \delta_n = \lambda$, say. Now let A be the matrix whose ij th entry is $a_{ij} = s_{ij} + \frac{1}{\lambda} \delta_i \varepsilon_j$. It is easily checked that A is stochastic, dominates S , and satisfies $Av = v$. $\square$

Proof of Theorem. In view of the lemma, we may assume that $Su \leq u$ and $Sv \leq v$, and we have to show that $Sw \leq w$. Using the Hölder inequality, we get

$$\begin{aligned} (Sw)_i &= \sum_j s_{ij} w_j = \sum_j s_{ij} u_j^\alpha v_j^\beta = \sum_j (s_{ij} u_j)^\alpha (s_{ij} v_j)^\beta \\ &\leq \left(\sum_j s_{ij} u_j \right)^\alpha \left(\sum_j s_{ij} v_j \right)^\beta \leq (u_i)^\alpha (v_i)^\beta = w_i. \quad \square \end{aligned}$$

REFERENCES

- [C] J. Cohen, *Convexity of the dominant eigenvalue of an essentially nonnegative matrix*, Proc. Amer. Math. Soc. **81** (1981), 657–658.
- [DN] E. Deutsch and M. Neumann, *On the first and second order derivatives of the Perron vector*, Linear Algebra Appl. **71** (1985), 57–76.
- [EJN] L. Elsner, C. Johnson, and M. Neumann, *On the effect of the perturbation of a nonnegative matrix on its Perron eigenvector*, Czech. Math. J. **32** (1982), 99–109.
- [F] S. Friedland, *Convex spectral functions*, Linear and Multilinear Algebra **9** (1981), 299–316.
- [K] J. Kingman, *A convexity property of positive matrices*, Quart. J. Math. **81** (1961), 283–284.
- [S] E. Seneta, *Non-negative matrices*, George Allen & Unwin, London, 1973.
- [SY] S. Sahi and S. Yao, *The non-cooperative equilibria of a trading economy with complete markets and consistent prices*, J. Math. Econ. **18** (1989), 325–346.

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY, PRINCETON, NEW JERSEY 08544
E-mail address: sahi@math.princeton.edu