

## RATIOS OF REGULATORS IN EXTENSIONS OF NUMBER FIELDS

ANTONE COSTA AND EDUARDO FRIEDMAN

(Communicated by William W. Adams)

**ABSTRACT.** Let  $L/K$  be an extension of number fields. Then

$$\text{Reg}(L)/\text{Reg}(K) > c_{[L:\mathbf{Q}]}(\log |D_L|)^m,$$

where  $\text{Reg}$  denotes the regulator,  $D_L$  is the absolute discriminant of  $L$ , and  $c_{[L:\mathbf{Q}]} > 0$  depends only on the degree of  $L$ . The nonnegative integer  $m = m(L/K)$  is positive if  $L/K$  does not belong to certain precisely defined infinite families of extensions, analogous to CM fields, along which  $\text{Reg}(L)/\text{Reg}(K)$  is constant. This generalizes some inequalities due to Remak and Silverman, who assumed that  $K$  is the rational field  $\mathbf{Q}$ , and modifies those of Bergé-Martinet, who dealt with a general extension  $L/K$  but used its relative discriminant where we use the absolute one.

### 1. INTRODUCTION

Remak [R1] laid down the principle that a number field ought to have a large regulator if and only if it has a large discriminant. In one direction this follows from work of Landau [L, Sie], who proved that  $\sqrt{|D_L|}(\log |D_L|)^{[L:\mathbf{Q}]-1}$  is an upper bound for  $\text{Reg}(L)$ . To obtain an inequality in the opposite sense, Remak considered the field  $\mathbf{Q}(E_L)$  generated by the units  $E_L$  of  $L$ . The geometry of numbers tells us that  $\mathbf{Q}(E_L)$  can be generated by integral elements (units) whose size at every embedding is bounded in terms of  $\text{Reg}(L)$ . It follows that  $|D_{\mathbf{Q}(E_L)}|$  can be bounded above by a function of  $\text{Reg}(L)$ . Remak then observed that  $\mathbf{Q}(E_L) = L$  unless  $L$  is a CM field (a totally imaginary quadratic extension of a totally real field). Thus he proved [R1]

$$(1.1) \quad \text{Reg}(L) > C_N \log \left( \frac{|D_L|}{N^N} \right),$$

where  $L$  is assumed non-CM,  $N = [L:\mathbf{Q}]$ , and  $C_N > 0$  depends explicitly on  $N$ . In 1984 Silverman [Sil] improved the dependence on  $\log |D_L|$  in (1.1) to

$$\text{Reg}(L) > 2^{-4N^2} \left( \log \left( \frac{|D_L|}{N^{N \log_2(8N)}} \right) \right)^{r_L - \rho},$$

---

Received by the editors September 25, 1991 and, in revised form, February 18, 1992.

1991 *Mathematics Subject Classification*. Primary 11R27, 11R29.

*Key words and phrases*. Regulator, discriminant, unit-weak extensions.

The first author was partially supported by NSA grant MDA90-H-1019.

Research was supported by the Max Planck Institut für Mathematik and the NSF.

where  $|D_L| > N^{N \log_2(8N)}$  is assumed,  $r_L$  is the unit rank of  $L$ , and  $\rho = \max_{F \subsetneq L} \{r_F\}$ .

It follows from (1.1) that given an integer  $N$  and a real number  $\gamma$  there are only finitely many non-CM number fields  $L$  such that  $[L : \mathbf{Q}] \leq N$  and  $\text{Reg}(L) < \gamma$ . CM fields must be excluded since the regulator, being essentially that of a proper subfield, can have the same value for infinitely many CM fields. We can, however, drop all restrictions on the degree  $[L : \mathbf{Q}]$  by using Zimmert's [Z] bound

$$\text{Reg}(L) > (0.04)1.05^{[L : \mathbf{Q}]}$$

In the late 1980s Bergé and Martinet [BM1, BM2] generalized Remak and Silverman's method to the relative case. Given an extension  $L/K$  of number fields their idea was to equate the ratio of regulators  $\text{Reg}(L)/\text{Reg}(K)$  with the covolume of a lattice produced from the units of  $L$ . In their approach the absolute norm  $N(\mathcal{D}_{L/K})$  of the relative discriminant of  $L/K$  appeared naturally and they were able to bound  $\text{Reg}(L)/\text{Reg}(K)$  from below by a power of  $\log(N(\mathcal{E}_{L/K}))$ .

While Bergé and Martinet's results can be used quite effectively [BM3] if  $N(\mathcal{D}_{L/K})$  is large, they are otherwise not so strong. This makes it difficult to obtain inequalities in which  $K$  is allowed to vary, say only fixing  $[L : \mathbf{Q}]$ , as there will be in general infinitely many  $L/K$  with  $N(\mathcal{D}_{L/K}) = 1$ . Our results for totally real fields [CF] suggest that this problem could be overcome by modifying Bergé and Martinet's lattice. We use the lattice associated to the relative units  $E_{L/K}$ . By definition,  $E_{L/K}$  consists of those units of  $L$  whose norm to  $K$  is a root of unity. Since the covolume of  $E_{L/K}$  under the logarithmic embedding is readily related to  $\text{Reg}(L)/\text{Reg}(K)$ , we can apply Remak's geometric method to bound the absolute discriminant of  $\mathbf{Q}(E_{L/K})$  from above in terms of  $\text{Reg}(L)/\text{Reg}(K)$ . It turns out that  $\mathbf{Q}(E_{L/K}) = L$ , except when one of the following three conditions holds:

- (i)  $L = K$ .
- (ii) The field  $L$  is CM (and  $K$  is any subfield of  $L$ ).
- (iii) There is a CM field  $M$  with maximal totally real subfield  $k$  such that  $K$  is a quadratic extension of  $k$ ,  $K \neq M$ , and  $L = MK$ .

We call the extension  $L/K$  *unit-weak* if it satisfies (i), (ii) or (iii) above.

**Theorem.** *Let  $E_{L/K}$  denote, as above, the relative units of an extension  $L/K$  of number fields. Assume that  $|D_L| > 3N^N$ , where  $D_L$  is the discriminant of  $L/\mathbf{Q}$  and  $N = [L : \mathbf{Q}]$ . Then*

$$(1.2) \quad \frac{\text{Reg}(L)}{\text{Reg}(K)} > \frac{C}{N^{2r}} \left( \log \left( \frac{|D_L|}{N^N} \right) \right)^m,$$

where  $r = \text{rank}(E_{L/K}) = r_L - r_K$  is the difference of the unit ranks of  $L$  and  $K$ ,  $\text{Reg}$  is the regulator, and  $C > 0$  is a computable absolute constant. The non-negative integer  $m$  is positive if  $L/K$  is not unit-weak (see the above definition). In general,  $m = m(L/K) = r - \max_{F \subsetneq L} \{\text{rank}(E_{L/K} \cap F)\}$ , where the maximum is taken as  $F$  runs over all proper subfields of  $L$ .

We actually prove the somewhat stronger inequality (3.7) in which  $\text{Reg}(L)/\text{Reg}(K)$  is replaced by the regulator of  $E_{L/K}$ . The exponent  $m$  of  $\log(|D_L|)$  in (1.2) is likely to be best possible. In any case,  $m$  can be computed

by easy linear algebra, without knowledge of any unit, as long as one knows all the subfields of  $L$  (see the end of §3). In contrast, we do not calculate  $C$  here, as we do not obtain a good value. Our proof does yield that one can take  $C = 1$  and  $m = r$ , provided we assume that every proper subfield of  $L$  is actually a subfield of  $K$ .

When  $L/K$  is unit-weak  $m$  vanishes and (1.2) becomes almost useless, however, in this case the ratio of regulators  $\text{Reg}(L)/\text{Reg}(K)$  is essentially that of a proper subextension. Unit-weak extensions can thus be treated inductively and represent no essential complication to the problem of bounding  $\text{Reg}(L)/\text{Reg}(K)$  from below. We treat unit-weak extensions briefly at the end of §§2 and 3.

A consequence of (1.2) is

**Corollary.** *Given an integer  $N$  and a real number  $y$ , there are at most finitely many extensions  $L/K$  such that  $[L:\mathbf{Q}] \leq N$ ,  $\text{Reg}(L)/\text{Reg}(K) < y$ , and  $L/K$  is not unit-weak.*

If  $L$  is totally real even more is true: Given any real number  $y$  there are finitely many pairs of *totally real* fields  $L$  and  $K$ , with  $K \subsetneq L$ , such that  $\text{Reg}(L)/\text{Reg}(K) < y$  [CF]. We do not know yet if this extends to all non-unit-weak  $L/K$ , totally real or not.

## 2. THE FIELD GENERATED BY THE RELATIVE UNITS

Recall that the group of relative units  $E_{L/K}$  of an extension  $L/K$  of number fields is defined by

$$E_{L/K} = \{\alpha \in E_L \mid \text{Norm}_{L/K}(\alpha) \in W_K\},$$

where  $E_L$  denotes the units of  $L$  and  $W_K$  the torsion subgroup of  $E_K$ . The (free) rank of  $E_{L/K}$  is  $r = r_{L/K} = r_L - r_K$ , where  $r_L$  is the rank of  $E_L$ . Let  $\mathcal{S}_L$  denote the set of embeddings of  $L$  into  $\mathbf{C}$ . We embed  $E_L/W_L$  into  $\mathbf{R}^{\mathcal{S}_L}$  by the map  $\mathcal{L} = \mathcal{L}_L: E_L \rightarrow \mathbf{R}^{\mathcal{S}_L}$  defined by

$$(2.1) \quad (\mathcal{L}_L(\alpha))_\sigma = (\mathcal{L}(\alpha))_\sigma = \log |\sigma(\alpha)|, \quad \sigma \in \mathcal{S}_L.$$

We endow  $\mathbf{R}^{\mathcal{S}_L}$  with the Euclidean inner product

$$(2.2) \quad \langle (x_\sigma), (y_\sigma) \rangle = \sum_{\sigma \in \mathcal{S}_L} x_\sigma y_\sigma.$$

Then  $\mathcal{L}_L(E_{L/K})$  is perpendicular to  $\mathcal{L}_L(E_K)$ . A dimension count shows that the  $\mathbf{Q}$ -spans  $\mathbf{Q}\mathcal{L}_L(E_{L/K})$  and  $\mathbf{Q}\mathcal{L}_L(E_K)$  of these two lattices are orthogonal complements of each other inside  $\mathbf{Q}\mathcal{L}(E_L)$ .

Our first goal is to characterize the extensions  $L/K$  for which  $\mathbf{Q}(E_{L/K})$  is a proper subfield of  $L$ . Slightly more generally, we prove

**Proposition 1.** *Let  $L/K$  be an extension of number fields and let  $E_{L/K}$  be its group of relative units. Let  $E$  be a subgroup of finite index in  $E_{L/K}$  and suppose that  $E$  is contained in a proper subfield of  $L$ . Then at least one of (i), (ii), or (iii) below holds:*

- (i)  $L = K$ .
- (ii)  $L$  is CM (and  $K \subset L$  is arbitrary).
- (iii) There is a CM field  $M$  with maximal totally real subfield  $k$  such that  $K$  is a quadratic extension of  $k$ ,  $K \neq M$ , and  $L = MK$ .

Conversely, if (iii), (ii), or (i) holds (with  $L \neq \mathbf{Q}$ ), then  $E_{L/K}$  contains a subgroup  $E$  as above.

*Proof.* The last statement is obvious in cases (i) and (ii). If (iii) holds, let  $H \neq K$ ,  $H \neq M$  be the third field lying strictly between  $k$  and  $L$ . A short computation shows that  $E := E_{H/k} \subset H$  has the same rank as  $E_{L/K}$  and  $E_{H/k} \subset E_{L/K}$ , proving the converse claim.

We now prove the first part of the proposition. Given a subfield  $F \subset L$  and an archimedean place  $\omega$  of  $L$ , let  $e_F(\omega) = e_{L/F}(\omega) = 2$  if  $\omega$  ramifies in  $L/F$ ; otherwise, let  $e_F(\omega) = 1$ . Let  $\infty_F$  denote the set of archimedean places of  $F$ . Then

$$(2.3) \quad r_F + 1 = \frac{1}{[L:F]} \sum_{\omega \in \infty_L} e_F(\omega),$$

because

$$r_F + 1 = \sum_{\nu \in \infty_F} 1 = \sum_{\nu \in \infty_F} \frac{1}{[L:F]} \sum_{\substack{\omega \in \infty_L \\ \omega|\nu}} e_F(\omega) = \frac{1}{[L:F]} \sum_{\omega \in \infty_L} e_F(\omega).$$

Let  $H = \mathbf{Q}(E)$ . Then  $H \subsetneq L$ , by assumption. Since  $E \subset E_H$ , we have  $r_H \geq r_{L/K} = r_L - r_K$ . From this and (2.3) we obtain

$$\frac{1}{[L:H]} \sum_{\omega \in \infty_L} e_H(\omega) + \frac{1}{[L:K]} \sum_{\omega \in \infty_L} e_K(\omega) > \sum_{\omega \in \infty_L} 1.$$

The compositum  $HK \subset L$  contains  $E$  and  $E_K$ . Modulo torsion, these are disjoint (perpendicular!) subgroups of  $E_L/W_L$  of rank  $r_L - r_K$  and  $r_K$ ; hence, the units of  $HK$  have rank  $r_L$ . If  $HK \neq L$ , then  $L$  must be a CM field, in which case the proof is done. We may therefore assume  $HK = L$ . Then we cannot simultaneously have  $e_H(\omega) = 2$  and  $e_K(\omega) = 2$  for  $\omega \in \infty_L$ . Hence,

$$(2.4) \quad \left( \frac{1}{[L:H]} + \frac{1}{[L:K]} \right) \sum_{\omega \in \infty_L} 1 + \max \left( \frac{1}{[L:H]}, \frac{1}{[L:K]} \right) \sum_{\omega \in \infty_L} 1 > \sum_{\omega \in \infty_L} 1.$$

By assumption,  $[L:H] \geq 2$ . Thus, either  $[L:H] = 2$  or  $[L:K] = 2$  (we dismiss the trivial case  $L = K$ ).

We first assume  $[L:K] = 2$ . Let  $\tau$  be the nontrivial element of  $\text{Gal}(L/K) \cong \mathbf{Z}/2\mathbf{Z}$ . For  $\alpha \in E \subset E_{L/K}$ , we have  $\text{Norm}_{L/K}(\alpha) \in W_K$ ; therefore,  $\tau(\alpha) = \eta\alpha^{-1}$ ,  $\eta \in W_K$ . By passing, as we may, to a subgroup of finite index in  $E$ , we can assume  $\tau(\alpha) = \alpha^{-1}$ ; hence,  $\tau$  induces a nontrivial field automorphism of  $H = \mathbf{Q}(E)$ . Let  $H_\tau$  be its fixed field so that  $[H:H_\tau] = 2$ . Since  $H_\tau \subset L_\tau = K$ , we must have either  $H \cap K = H_\tau$  or  $H \cap K = H$ . In the latter case we would have  $E \subset K$ . But then  $E \subset K \cap E_{L/K} = W_K$ . Since  $E$  has finite index in  $E_{L/K}$ , this could only happen if  $L$  is CM. We may thus assume  $H \cap K = H_\tau$ . Then  $E \subset H \cap E_{L/K} = E_{H/H \cap K} \subset E_{L/K}$ . Since  $E$  has finite index in  $E_{L/K}$ ,  $r_{H/H \cap K} = r_{L/K}$ . From this and (2.3) we find

$$\frac{1}{[L:H]} \sum_{\omega \in \infty_L} e_H(\omega) - \frac{1}{[L:H \cap K]} \sum_{\omega \in \infty_L} e_{H \cap K}(\omega) = \sum_{\omega \in \infty_L} 1 - \frac{1}{2} \sum_{\omega \in \infty_L} e_K(\omega).$$

Since  $[L:H \cap K] = 2[L:H]$ , we have

$$(2.5) \quad \frac{1}{[L:H]} \sum_{\omega \in \infty_L} (2e_H(\omega) - e_{H \cap K}(\omega)) = \sum_{\omega \in \infty_L} (2 - e_K(\omega)).$$

Observe that if  $\omega$  ramifies in  $L/K$ , then  $\omega$  ramifies in  $L/H \cap K$  but not in  $L/H$  (since  $L = HK$ ). Thus, if  $e_K(\omega) = 2$ , then  $2e_H(\omega) - e_{H \cap K}(\omega) = 0$ . If  $e_K(\omega) = 1$ , then  $2e_H(\omega) - e_{H \cap K}(\omega) \leq 2$ . It now follows from (2.5) that  $[L : H] = 2$  and that  $e_H(\omega) = 2$  if and only if  $e_K(\omega) = 1$ . Hence  $[L : H] = 2 = [H : H \cap K] = [K : H \cap K]$  and all archimedean places of  $L$  ramify in either  $L/K$  or  $L/H$ , but none in both extensions. It follows that  $L/K$  satisfies condition (iii) in the proposition (let  $k = K \cap H$  and let  $M \neq K$ ,  $M \neq H$ , be the third field lying strictly between  $k$  and  $L$ ). This proves Proposition 1 when  $[L : K] = 2$ .

If  $[L : K] > 2$ , then (2.4) implies  $[L : H] = 2$ . The strategy now is to reverse the roles of  $H$  and  $K$  and thereby reduce the proof to the quadratic case that we just handled. Recall that if  $F$  is any subfield of  $L$ , then the  $\mathcal{Q}$ -spans of  $\mathcal{L}(E_{L/F})$  and  $\mathcal{L}(E_F)$  are orthogonal with respect to the ( $\mathbf{R}$ -valued) inner product (2.2). By construction,  $\mathcal{L}(E) \subset \mathcal{L}(E_H)$ . Since  $E$  has finite index in  $E_{L/K}$ ,  $\mathcal{Q}\mathcal{L}(E) = \mathcal{Q}\mathcal{L}(E_{L/K})$ . Hence

$$(2.6) \quad \mathcal{Q}\mathcal{L}(E_{L/H}) = \mathcal{Q}\mathcal{L}(E_H)^\perp \subset \mathcal{Q}\mathcal{L}(E_{L/K})^\perp = \mathcal{Q}\mathcal{L}(E_K),$$

where  $^\perp$  denotes the orthogonal complement inside  $\mathcal{Q}\mathcal{L}(E_L)$ . Since the kernel  $W_L$  of  $\mathcal{L}$  is finite, (2.6) shows that  $E_{L/H}^n \subset E_K$  for some positive integer  $n$ . Thus  $E' := E_{L/H}^n$  has finite index in  $E_{L/H}$ ,  $[L : H] = 2$ , and  $\mathcal{Q}(E') \subset K$ , a proper subfield of  $L$ ; but this is the quadratic case of the proposition, so the proof is done.

We conclude this section with a brief discussion of the unit-index  $u_{L/K}$  of a unit-weak extension  $L/K$ . We assume first that  $K \neq L$  and that  $L$  is not CM. Let  $k$  and  $M$  be as in (iii) above. Denote by  $K$  and  $H$  the two remaining fields lying strictly between  $k$  and  $L$ . Let  $\tau_H$ ,  $\tau_K$ , and  $\tau_M = \tau_H\tau_K$  be the nontrivial automorphisms of  $L/H$ ,  $L/K$ , and  $L/M$ . Since we assume that  $L$  is not CM, at least one archimedean place of  $k$  ramifies in  $H$ ; hence, at least one archimedean place of  $K$  ramifies in  $L$ . Thus  $W_K = \{\pm 1\}$  and  $-1$  is not a norm in  $L/K$ , whence  $\text{Norm}_{L/K}(E_{L/K}) = \{+1\}$ . Equivalently,  $\tau_K(\alpha) = \alpha^{-1}$  for  $\alpha \in E_{L/K}$ . Hence,  $\text{Norm}_{L/M}(\alpha) = \alpha\tau_H(\tau_K(\alpha)) = \alpha/\tau_H(\alpha)$ . Therefore,  $\text{Norm}_{L/M}(\alpha) = 1$  if and only if  $\alpha \in E_{L/K} \cap H = E_{H/k}$ . In short,  $\text{Norm}_{L/M}$  induces an injection of  $E_{L/K}/E_{H/k}$  into  $W_M = E_{M/k}$ . As  $W_M^2 = \text{Norm}_{L/M}(W_m) \subset \text{Norm}_{L/M}(W_L)$  and  $W_M$  is cyclic, we have  $u_{L/K} := [E_{L/K} : W_L E_{H/k}] = 1$  or  $2$ .

So far we have assumed that  $L$  is not CM. If  $L$  is CM, let  $H$  be its maximal totally real subfield. It is well known that  $[E_L : W_L E_H] = 1$  or  $2$  [R2]. It follows that  $u_{L/K} := [E_{L/K} : W_L E_{H/k}] = 1$  or  $2$ , where  $k = H \cap K$ . Finally, if  $L = K$  we let  $H = k = \mathbf{Q}$  and  $u_{L/K} = 1$ .

We have thus defined, whenever  $L/K$  is unit-weak, a subextension  $H/k$  and a unit-index  $u_{L/K} := [E_{L/K} : W_L E_{H/k}] = 1$  or  $2$ . When  $L$  is CM and  $K = \mathbf{Q}$ ,  $u_{L/\mathbf{Q}}$  is just the usual unit-index of  $L$ . In the next section we relate the regulators of  $E_{L/K}$  and  $E_{H/k}$  using  $u_{L/K}$ . Notice that  $H/k$  itself is not unit-weak unless  $r_{L/K} = 0$ .

### 3. PROOF OF THEOREM

We begin with the definition of the regulator of relative units  $\text{Reg}(E_{L/K})$ . Pick  $\alpha_1, \alpha_2, \dots, \alpha_r$  to be independent generators of  $E_{L/K}/W_L$ , the relative

units modulo torsion. Let  $M$  be the matrix  $M = (\log \|\alpha_i\|_\omega)$ , where  $1 \leq i \leq r$ ,  $\omega$  runs over the set  $\infty_L$  of archimedean places of  $L$ , and  $\|\cdot\|_\omega$  denotes the normalized absolute value at  $\omega$  (so that  $\|\cdot\|_\omega = |\cdot|_\omega^2$  if  $\omega$  is complex, and  $\|\cdot\|_\omega = |\cdot|_\omega$  otherwise). For each place  $\nu \in \infty_K$ , fix a place  $\omega_\nu \in \infty_L$  lying above  $\nu$ . Then  $\text{Reg}(E_{L/K})$  is the absolute value of the determinant of the submatrix of  $M$ , which results when we delete from  $M$  the rows corresponding to the  $\omega_\nu$ 's. In [CF, Theorem 1] we showed, for  $L/K$  of any signature,

$$(3.1) \quad \text{Reg}(E_{L/K}) = \frac{1}{[E_K : W_K \text{Norm}_{L/K}(E_L)]} \frac{\text{Reg}(L)}{\text{Reg}(K)}.$$

We also related [CF, Lemma 2.1]  $\text{Reg}(E_{L/K})$  to the  $r$ -dimensional volume  $V_L(E_{L/K})$  of a fundamental domain for  $\mathcal{L}(E_{L/K})$  (see (2.1)),

$$(3.2) \quad V_L(E_{L/K}) = [L : K]^{(r_1(K)+r_2(K))/2} 2^{(r_2(K)-r_2(L))/2} \text{Reg}(E_{L/K}),$$

where  $(r_1, r_2)$  denotes the number of (real, complex) places. The Euclidean structure (which normalizes volume) is given by  $\|(x_\sigma)\|^2 = \langle (x_\sigma), (x_\sigma) \rangle$ , as in (2.2). For  $\alpha \in E_L$  we write  $\|\alpha\|$  instead of  $\|\mathcal{L}(\alpha)\|$ . Thus,

$$(3.3) \quad \|\alpha\|^2 := \sum_{\sigma \in \mathcal{S}_L} (\log |\sigma(\alpha)|)^2,$$

where  $\mathcal{S}_L$  denotes the set of all embeddings of  $L$  into  $\mathbb{C}$ . We will need the lower bound [F, (3.21)]

$$(3.4) \quad \|\alpha\| > \frac{C'}{\sqrt{N}(\log N)^3},$$

where  $\alpha \in E_L$ ,  $\alpha \notin W_L$ ,  $N = [L : \mathbb{Q}]$ , and  $C' > 0$  is a computable absolute constant (inequality (3.4) follows easily from Dobrowolsky's lower bound for heights [D]).

Let the successive minima of  $\|\cdot\|$  on the lattice  $\mathcal{L}(E_{L/K})$  be attained at  $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_r$ . Thus [GK, pp. 195, 197] the subgroup  $E := \langle \varepsilon_1, \varepsilon_2, \dots, \varepsilon_r \rangle$  of  $E_{L/K}$  generated by the  $\varepsilon_i$  has finite index in  $E_{L/K}$  and

$$(3.5) \quad 0 < \|\varepsilon_1\| \leq \|\varepsilon_2\| \leq \dots \leq \|\varepsilon_r\|,$$

$$(3.6) \quad \prod_{i=1}^r \|\varepsilon_i\| \leq \gamma_r^{r/2} V_L(E_{L/K}),$$

where  $\gamma_r$  denotes Hermite's constant in dimension  $r = r_{L/K}$ .

**Lemma.** *Let  $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_r$  be as above and assume that  $L/K$  is not unit-weak (see §1). Let  $H_0 = \mathbb{Q}$  and  $H_i = H_{i-1}(\varepsilon_i)$ . Then there is an integer  $T$  such that  $H_T \neq L$ ,  $H_{T+1} = L$ ,  $0 \leq T < r$ , and*

$$\frac{1}{[L : \mathbb{Q}]} \log |D_L| \leq \log([L : \mathbb{Q}]) + \frac{1}{\sqrt{3}[L : \mathbb{Q}]} \sum_{i=1}^{T+1} \|\varepsilon_i\| \sqrt{[H_i : H_{i-1}]^2 - 1},$$

where  $D_L$  denotes the absolute discriminant of  $L$  and  $\|\cdot\|$  is given by (3.3).

*Proof.* Proposition 1 implies that there is at least  $T < r$  so that  $L = \mathbb{Q}(\varepsilon_1, \varepsilon_2, \dots, \varepsilon_{T+1})$ . The inequality then follows from [F, (3.3), (3.14) and Lemma 3.5].

**Theorem.** Let  $L/K$  be an extension of number fields and assume that  $D_L > 3N^N$ , where  $D_L$  is the absolute discriminant of  $L$  and  $N = [L : \mathbf{Q}]$ . Then

$$(3.7) \quad \text{Reg}(E_{L/K}) \geq \frac{C}{N^{2r}} \left( \log \left( \frac{|D_L|}{N^N} \right) \right)^m.$$

Here  $\text{Reg}(E_{L/K})$  is the regulator of relative units given by (3.1),  $C > 0$  is a computable absolute constant, and  $r = r_L - r_K = \text{rank}(E_{L/K})$  is the difference of the unit ranks of  $L$  and  $K$ . The nonnegative integer  $m$  is positive if  $L/K$  is not unit-weak (see §1). In general,  $m = m(L/K) = r - \max_{F \subsetneq L} \{\text{rank}(E_{L/K} \cap F)\}$ , where  $F$  runs over all proper subfields of  $L$ .

The slightly simplified version of the theorem given in §1 follows from (3.1) and (3.7).

*Proof.* We first assume that  $L/K$  is not unit-weak. From the Lemma and (3.5) we have

$$(3.8) \quad \frac{1}{N} \log \left( \frac{|D_L|}{N^N} \right) \leq \frac{\|\varepsilon_{T+1}\|}{\sqrt{3N}} \sum_{i=1}^{T+1} \sqrt{[H_i : H_{i-1}]^2 - 1} \leq \|\varepsilon_{T+1}\| \sqrt{\frac{N}{3}},$$

since  $\prod_{i=1}^{T+1} [H_i : H_{i-1}] = N$ . From (3.5), (3.6), and (3.4)

$$(3.9) \quad \|\varepsilon_{T+1}\|^{r-T} \leq \prod_{i=T+1}^r \|\varepsilon_i\| \leq \frac{\gamma_r^{r/2} V_L(E_{L/K})}{(C'/\sqrt{N}(\log N)^3)^T}.$$

If we put this together with (3.2) and (3.8) and use  $\log(|D_L|/N^N) > 0$ , we find

$$(3.10) \quad \begin{aligned} & \frac{1}{N^{2r}} \left( \log \left( \frac{|D_L|}{N^N} \right) \right)^{r-T} \\ & \leq \frac{(([L : K]/2)^{(r_1(K)+r_2(K))/r} 2^{([K : \mathbf{Q}] - r_2(L))/r} \gamma_r / 3N)^{r/2}}{(NC'/\sqrt{3}(\log N)^3)^T} \text{Reg}(E_{L/K}). \end{aligned}$$

If  $[L : K] \geq 3$ , then (2.3) yields

$$r = r_L - r_K = \sum_{\omega \in \infty_L} \left( 1 - \frac{e_K(\omega)}{[L : K]} \right) \geq \sum_{\omega \in \infty_L} \frac{1}{3} \geq \frac{[L : \mathbf{Q}]}{6}.$$

Hence, for  $[L : K] \geq 2$ ,

$$(3.11) \quad \left( \frac{[L : K]}{2} \right)^{(r_1(K)+r_2(K))/r} \leq \left( \frac{[L : K]}{2} \right)^{6/[L : K]} < 3.003.$$

Note that

$$(3.12) \quad [K : \mathbf{Q}] - r_2(L) \leq r_1(L) + r_2(L) - r_1(K) - r_2(K) = r$$

and that, for  $r > 2$ ,  $\gamma_r \leq r/2.1$ . (*Proof.* Use the inequalities quoted in [CF, (2.9)]). We then have in (3.10)

$$(3.13) \quad \left( \left( \frac{[L : K]}{2} \right)^{(r_1(K)+r_2(K))/r} 2^{([K : \mathbf{Q}] - r_2(L))/r} \frac{\gamma_r}{3N} \right)^{r/2} \leq 1,$$

for all  $r > 0$  (do  $e = 1$  or  $2$  separately). Since  $T < r < N$ , (3.10) and (3.13) yield

$$(3.14) \quad \text{Reg}(E_{L/K}) > \frac{C}{N^{2r}} \left( \log \left( \frac{|D_L|}{N^N} \right) \right)^{r-T},$$

with  $C > 0$  a computable absolute constant. To prove (3.7) we must still show that in (3.14) we can replace  $T$  by  $\rho := \max_{F \subsetneq L} \{\text{rank}(E_{L/K} \cap F)\}$ . Since we assume  $D_L > 3N^N$ , it suffices to show  $T \leq \rho$ . By the lemma,  $H_T$  is a proper subfield of  $L$  containing the  $T$  independent relative units  $\varepsilon_1, \varepsilon_2, \dots, \varepsilon_T \in E_{L/K}$ ; hence,  $T \leq \rho$ . Proposition 1 implies that  $m = r - \rho > 0$  which concludes the proof when  $L/K$  is not unit-weak.

If  $L/K$  is unit-weak then  $m = r - \rho = 0$  in (3.7). In this case (3.7) follows from

**Proposition 2.** *Let  $L/K$  be an extension of number fields. Then*

$$(3.15) \quad \text{Reg}(E_{L/K}) \geq \frac{c^r}{(Nr(\log N)^6)^{r/2}}.$$

Here  $\text{Reg}(E_{L/K})$  is the regulator of relative units given by (3.1),  $c > 0$  is a computable absolute constant,  $N = [L : \mathbf{Q}]$ , and  $r = r_L - r_K$  is the difference of the unit ranks of  $L$  and  $K$ . (If  $r = 0$ , (3.15) means the trivial  $1 \geq 1$ .)

*Proof.* From (3.4), (3.6), and (3.2) we obtain

$$\text{Reg}(E_{L/K}) \geq \left( \frac{C'^2}{N\gamma_r(\log N)^6([L : K]/2)^{(r_1(K)+r_2(K))/r} 2^{([K : \mathbf{Q}] - r_2(L))/r}} \right)^{r/2}.$$

Now use (3.11), (3.12), and  $\gamma_r \leq r$  to obtain (3.15), with  $c = C'\sqrt{6.006}$ .

**Corollary.** *Let  $L/K$  (and all notation) be as in the theorem. Suppose further that all proper subfields of  $L$  are actually subfields of  $K$ . Then*

$$(3.16) \quad \text{Reg}(E_{L/K}) \geq \frac{1}{N^{2r}} \left( \log \left( \frac{|D_L|}{N^N} \right) \right)^r.$$

*Proof.* We first dispose of the trivial cases. If  $L/K$  is unit-weak, the hypothesis on  $K$  implies that case (iii) in Proposition 1 cannot hold. If (ii) holds, so  $L$  is CM, then  $K$  must be its maximal totally real subfield. Then  $r = 0$  and (3.16) is trivial. Since case (i) ( $L = K$ ) is equally trivial, we may assume that  $L/K$  is not unit weak. Consider, in the notation of the lemma,  $H_1 = \mathbf{Q}(\varepsilon_1)$ . By assumption, either  $H_1 \subseteq K$  or  $H_1 = L$ . But  $H_1 \subseteq K$  implies  $\varepsilon_1 \in E_{L/K} \cap K$ , which is impossible since  $\varepsilon_1$  is not a root of unity. Thus,  $H_1 = L$  and so  $T = 0$  in the lemma. The corollary now follows from (3.13) and (3.10).

The computation of  $m = m(L/K)$  in the theorem turns out to be elementary. Let  $\mathcal{L}_L$  be the logarithmic embedding (2.1). If  $M \subset \mathcal{L}_L(E_L) \subset \mathbf{R}^{\mathcal{L}_L}$  is a lattice, denote its  $\mathbf{R}$ -span by  $\mathbf{R}M$ . Thus,  $\text{rank}(M) = \dim_{\mathbf{R}}(\mathbf{R}M)$ . If  $F$  is a subfield of  $L$ , observe that

$$\begin{aligned} & \text{rank}(E_{L/K}) + \text{rank}(E_F) - \text{rank}(E_{L/K} \cap F) \\ &= r_L - r_K + r_F - \text{rank}(E_{L/K} \cap F) \\ &= \text{rank}(E_{L/K} E_F) = \dim_{\mathbf{R}}(\mathbf{R}\mathcal{L}_L(E_{L/K} E_F)) \\ &= \dim_{\mathbf{R}}(\mathbf{R}\mathcal{L}_L(E_{L/K}) + \mathbf{R}\mathcal{L}_L(E_F)). \end{aligned}$$



Dirichlet's unit theorem gives an  $\mathbf{R}$ -basis of  $\mathbf{R}\mathcal{L}_L(E_F)$ . It also gives one for  $\mathbf{R}\mathcal{L}_L(E_{L/K})$  as the orthogonal complement of  $\mathbf{R}\mathcal{L}_L(E_K)$  (inside  $\mathbf{R}\mathcal{L}_L(E_L)$ ). It follows that

$$m := \text{rank}(E_{L/K}) - \max_{F \subsetneq L} \{\text{rank}(E_{L/K} \cap F)\}$$

can be calculated by linear algebra from a knowledge of all the subfields of  $L$ , without knowing a single unit. To be precise, one has to know, for each subfield  $F$  of  $L$ , the mapping  $\mathcal{S}_L \rightarrow \mathcal{S}_F$  obtained by restricting the embeddings of  $L$  to embeddings of  $F$ .

We conclude with a comment on  $\text{Reg}(E_{L/K})$  and  $\text{Reg}(L)/\text{Reg}(K)$  for  $L/K$  unit-weak. We defined in §2 a subextension  $H/k$  and a unit index

$$u_{L/K} := [E_{L/K} : W_L E_{H/k}] = 1 \text{ or } 2.$$

On examining the ramification of the archimedean places in  $L/K$  and  $H/k$  one finds, directly from the definition of  $\text{Reg}(E_{L/K})$  as a determinant,

$$(3.17) \quad \text{Reg}(E_{L/K}) = 2^{r_{H/k}} \text{Reg}(E_{H/k}) / u_{L/K}.$$

If we let  $L/K$  range over the infinitely many unit-weak extensions associated to the same  $H/k$ , it is clear from (3.17) that  $\text{Reg}(E_{L/K})$  assumes at most two values. It follows, mainly from (3.1), that  $\text{Reg}(L)/\text{Reg}(K)$  assumes at most  $2^{[H:\mathbf{Q}]}$  values.

## REFERENCES

- [BM1] A.-M. Bergé and J. Martinet, *Sur les minorations géométriques des régulateurs*, Séminaire de Théorie des Nombres de Paris 1987–1988 (C. Goldstein, ed.), Birkhäuser Verlag, Boston, MA, 1990, pp. 23–50.
- [BM2] ———, *Notions relatives de régulateur et de hauteur*, Acta Arith. **54** (1989), 155–170.
- [BM3] ———, *Minorations de hauteurs et petits régulateurs relatifs*, Séminaire de Théorie des Nombres de Bordeaux 1987–1988, Univ. de Bordeaux, 1989.
- [CF] A. Costa and E. Friedman, *Ratios of regulators in totally real extensions of number fields*, J. Number Theory **37** (1991), 288–297.
- [D] E. Dobrowolsky, *On a question of Lehmer and the number of irreducible factors of a polynomial*, Acta Arith. **34** (1979), 391–401.
- [F] E. Friedman, *Analytic formulas for the regulator of a number field*, Invent. Math. **98** (1989), 599–622.
- [GK] P. M. Gruber and C. G. Lekkerkerker, *Geometry of numbers*, North-Holland, Amsterdam, 1987.
- [L] E. Landau, *Abschätzungen von Charaktersummen, Einheiten und Klassenzahlen*, Nachr. Kgl. Ges. Wiss. Göttingen Math.-Phys. Kl. (1918), 79–97; *Collected works*, vol. 7, Thales Verlag, Essen, 1985, pp. 114–132.
- [R1] R. Remak, *Über Größenbeziehungen zwischen Diskriminanten und Regulator eines algebraischen Zahlkörpers*, Compositio Math. **10** (1952), 245–285.
- [R2] ———, *Über algebraische Zahlkörper mit schwachem Einheitsdefekt*, Compositio Math. **12** (1954), 35–80.
- [Sie] C.-L. Siegel, *Abschätzungen von Einheiten*, Nachr. Akad. Wiss. Göttingen Math.-Phys. Kl. **9** (1969), 71–86; *Gesammelte Abhandlungen*, vol. 4, Springer-Verlag, Berlin, 1979, pp. 66–81.

- [Sil] J. Silverman, *An inequality connecting the regulator and discriminant of a number field*, J. Number Theory **19** (1984), 437–442.
- [Z] R. Zimmert, *Ideale kleiner Norm in Idealklassen und eine Regulatorabschätzung*, Invent. Math. **62** (1981), 367–380.

DEPARTMENT OF MATHEMATICS, AMERICAN UNIVERSITY, WASHINGTON, DISTRICT OF COLUMBIA  
20016

*Current address:* Department of Mathematics, George Washington University, Washington, DC  
20052

*E-mail address:* [costa@math.gwu.edu](mailto:costa@math.gwu.edu)

MAX PLANCK INSTITUT FÜR MATHEMATIK, GOTTFRIED CLAREN STRASSE 26, 5300 BONN 3,  
GERMANY