STABLE ABSOLUTELY UBIQUITOUS STRUCTURES

A. A. IVANOV

(Communicated by Andreas R. Blass)

ABSTRACT. It is proved that absolutely ubiquitous structures with trivial algebraic closures are monadically stable.

Let L be a finite language, and let M be an L -structure with countably infinite domain. The age of M , denoted by $\mathcal{F}(M)$, is the set of all isomorphism types of finite substructures of M . A model M is absolutely ubiquitous if M is uniformly locally finite and every countable locally finite L -structure with age $\mathcal{F}(M)$ is isomorphic to M . It is known that an absolutely ubiquitous structure is countably categorical and model-complete. It is not known if an absolutely ubiquitous structure has stable or superstable theory. In [1–3] a classification was given for absolutely ubiquitous structures over relational languages. Such structures are finitely partitioned. This will be proved here in a slightly more general case using completely different techniques.

Notation. $B < M$ if B is a substructure of M ,

$\text{Str}(B)$ is the substructure of M generated by B .

1. LEMMA OF E. A. PALYUTIN

The following lemma is a simple generalization of an old result of E. A. Palyutin on categorical universal theories [4].

Lemma 1. *A countable structure M is absolutely ubiquitous iff*

- (1) *M is uniformly locally finite; and*
- (2) *there is a function $s: \omega \rightarrow \omega$ such that for every n , for every substructure B of M , and for every type $p(y, b_0, \dots, b_{n-1})$ over $\{b_0, \dots, b_{n-1}\} \subseteq B$, if B contains all isomorphism types of $s(n)$ -generated substructures of M then there is $b \in B$ such that $M \models p(b, b_0, \dots, b_{n-1})$.*

Proof. Let M be absolutely ubiquitous. We construct s by induction. Suppose s is defined for all $i < n$. Suppose $R_0, \dots, R_t$ (and $T_0, \dots, T_l$) are complete formulas which determine all complete n -types ($(n+1)$ -types) over $\emptyset$ ($\text{Th}(M)$ is ω -categorical). By model-completeness we may suppose that R_i

Received by the editors March 26, 1992 and, in revised form, August 7, 1992.

1991 *Mathematics Subject Classification.* Primary 03C45, 03C35.

Key words and phrases. Stability, absolutely ubiquitous structures.

are $\forall$ -formulas and T_j are $\exists$ -formulas. We define the set $G_k^{i,j}$ as follows. Let

$$\Phi_{i,j} = \forall x_0, \dots, x_{n-1} (R_i(x_0, \dots, x_{n-1}) \rightarrow \exists y (T_j(y, x_0, \dots, x_{n-1}))).$$

If $M \models \Phi_{i,j}$ let $G_k^{i,j}$ be

$$\begin{aligned} \text{Th}_\forall(M) \cup \{ \text{there are } z_1, \dots, z_m \text{ which generate substructure embedding} \\ \text{all types of } k\text{-generated substructures of } M \} \\ \cup \{ \neg \Phi_{i,j} \}. \end{aligned}$$

If $M \models \neg \Phi_{i,j}$ let $G_k^{i,j}$ be $\forall x (x \neq x)$,

If for all k the set $G_k^{i,j}$ is consistent then there is a countable model N of the theory $\text{Th}_\forall(M)$ with $\mathcal{J}(N) = \mathcal{J}(M)$ and $N \models \neg \Phi_{i,j}$. It is a contradiction. Let

$$\begin{aligned} k(i, j) &= \min(k: G_k^{i,j} \text{ is inconsistent}), \\ s(n) &= \max(k(i, j): 0 \leq i \leq t, 0 \leq j \leq l). \end{aligned}$$

Now if B and p are as in (2), formula R_i determines $\text{tp}(\langle b_0, \dots, b_{n-1} \rangle)$ and formula T_j determines type $p(y, x_0, \dots, x_{n-1})$, then since $G_{s(n)}$ is inconsistent we have

$$B \models \forall x_0, \dots, x_{n-1} (R_i(x_0, \dots, x_{n-1}) \rightarrow \exists y T_j(y, x_0, \dots, x_{n-1})).$$

Since R_i is universal, there is $b \in B$ such that $B \models T_j(b, b_0, \dots, b_{n-1})$, and since T_j is $\exists$ -formula, $M \models T_j(b, b_0, \dots, b_{n-1})$.

If (1) and (2) hold in M then M is countably saturated. If for countable N we have $\mathcal{J}(N) = \mathcal{J}(M)$ then N is isomorphic to a substructure of M . By back-and-forth argument using (2) we can prove M is isomorphic to N .

2. STRUCTURES WITH A FINITE NUMBER OF TERM FUNCTIONS

We suppose here that there are terms $d_1, \dots, d_t$ such that for every term d the corresponding function under essential variables is one of the functions defined by $d_1, \dots, d_t$ in M . For example, a uniformly locally finite unary algebra is such a structure.

Proposition 1. *Let M be an absolutely ubiquitous structure with a finite number of term functions. Then the theory of M is stable.*

Proof. It is sufficient to prove that $\text{Th}(M)$ has not the independence property because by [5] $\text{Th}(M)$ has not the strict order property. For a contradiction suppose that there are a formula $F(\mathbf{x}, \mathbf{y})$ and sequences $\langle \mathbf{a}_i: i \in \omega \rangle$ and $\langle \mathbf{c}_\sigma: \sigma \in 2^\omega \rangle$ such that $\models F(\mathbf{a}_i, \mathbf{c}_\sigma)$ iff $\sigma(i) = 0$. For ease of notation suppose that $\mathbf{a}_i$ and $\mathbf{c}_\sigma$ are singletons. Assume terms $d_1, \dots, d_t$ depend on at most m variables. Let B be a finite substructure of M containing parameters of F and all types of $s(2m+2)$ -generated substructures of M . We may suppose that $\mathbf{a}_1, \dots, \mathbf{a}_{2m+2}$ is a sequence of order indiscernibles over B , and by Lemma 1 suppose

$$\{ \mathbf{c}_\sigma: \sigma \in 2^{2m+2} \} \subseteq \text{Str}(B \cup \{ \mathbf{a}_1, \dots, \mathbf{a}_{2m+2} \})$$

(of course, we neglect parameters). Define $\sigma \in 2^{2m+2}$ by $\sigma(i) = 0$ iff i is odd. We know that for some $\mathbf{a}_{j_1}, \dots, \mathbf{a}_{j_k}$ ($k \leq m$) and $\mathbf{b} \subseteq B$ and term d_i , $\mathbf{c}_\sigma = d_i(\mathbf{b}, \mathbf{a}_{j_1}, \dots, \mathbf{a}_{j_k})$. There are $\mathbf{a}_{2n-1}$ and $\mathbf{a}_{2n}$ ($n \leq m+1$) not among

$\mathbf{a}_{j_1}, \dots, \mathbf{a}_{j_k}$. By indiscernibility, $\models F(\mathbf{a}_{2n-1}, \mathbf{c}_\sigma)$ iff $\models F(\mathbf{a}_{2n}, \mathbf{c}_\sigma)$, which is a contradiction.

3. STRUCTURES WITH TRIVIAL CLOSURE

A structure M has trivial closure if $\text{acl}(X \cup Y) = \text{acl}(X) \cup \text{acl}(Y)$ holds in M . For example, absolutely ubiquitous structures with unary terms only have trivial closure. Indeed, by Lemma 1 there is some type of finite substructures of M such that for every B of this type $\text{acl}(X \cup Y)$ is in the substructure generated by $X \cup Y \cup B$. Since every term is unary, we have $\text{acl}(X \cup Y) \subseteq \text{acl}(X) \cup \text{acl}(Y) \cup B$. We can choose B with $B \cap \text{acl}(X \cup Y) \subseteq \text{acl}(\emptyset)$. So, $\text{acl}(X \cup Y) = \text{acl}(X) \cup \text{acl}(Y)$.

By [1, 6] absolutely ubiquitous structures in a relational language are monadically stable. By the description of countably categorical monadically stable structures [6], every absolutely ubiquitous monadically stable structure has trivial closure. It is not hard to find an absolutely ubiquitous structure with a finite number of term functions which does not have trivial closure. So, the situation of Proposition 1 is quite different than that of [1–3]. But in the case of trivial closure we have the following

Proposition 2. *Absolutely ubiquitous structures with trivial closure are monadically stable.*

Proof. Let us prove that the relation $x \in \text{acl}(y)$ on nonalgebraic elements is an equivalence relation. If $a \in \text{acl}(b)$ and $b \notin \text{acl}(a)$ then there is some type of finite substructures of our model M such that for every B of this type some realization of $\text{tp}(b|a)$ is in the substructure generated by $B \cup \{a\}$. By triviality of closure it is in $\text{acl}(B)$ and so $a \in \text{acl}(B)$. But we can choose $\text{acl}(B)$ so that $\text{acl}(B) \cap \{a\} \subseteq \text{acl}(\emptyset)$. We have now $a \in \text{acl}(\emptyset)$.

Let us prove that Shelah's R -rank of M is one. Let $\mathbf{a}_i$, $i \in \omega$, be an infinite sequence, and suppose the set of nonalgebraic complete formulas $F(x, \mathbf{a}_i)$ is n -inconsistent. Choose a finite $B < M$ containing all types of $s(\text{ln}(\mathbf{a}_i))$ -generated substructures of M . By Lemma 1 $\text{Str}(B \cup \{\mathbf{a}_i : i \in \omega\})$ contains realizations of all formulas $F(x, \mathbf{a}_i)$. Of course, the corresponding realization of formula $F(x, \mathbf{a}_i)$ is in $\text{Str}(B \cup \mathbf{a}_i)$ and by triviality of closure it is in $\text{acl}(B)$. This contradicts n -inconsistency of $\{F(x, \mathbf{a}_i) : i \in \omega\}$.

Since ω -categorical superstable theory is ω -stable, the Morley rank of M is one. There is $E_0 \in FE(\emptyset)$ with strongly minimal classes. Let $G = \{\text{acl}(a) - \text{acl}(\emptyset) : a \text{ is nonalgebraic}\}$. For $X_1, X_2 \in G$ let $\langle X_1, X_2 \rangle \in E_1$ iff X_1, X_2 meet the same E_0 -classes. Hence E_1 is a finite equivalence relation.

If a and b are in the same E_0 -class and c is a finite sequence disjoint to $(\text{acl}(a) \cup \text{acl}(b)) - \text{acl}(\emptyset)$ then $\text{tp}(a|c) = \text{tp}(b|c)$. Indeed otherwise for some formula $H(x, c)$, $M \models H(a, c) \& \neg H(b, c)$. For the corresponding strongly minimal type $p(x, \mathbf{d})$ (consisting of the E_0 -class with a and b) we have that, for example, $p(x, \mathbf{d}) \& H(x, c)$ is algebraic. Since acl is trivial and $x \in \text{acl}(y)$ is an equivalence relation, $a \in \text{acl}(\emptyset)$. This is a contradiction.

We now have a 0-definable equivalence relation E on M with finite classes such that M/E is finitely partitioned: E is $\text{acl}(x) = \text{acl}(y)$, E_1 -classes on $M/E - \text{acl}(\emptyset)/E$ are infinite, E_1 is invariant under $\text{Aut}(M)$, and for every such E_1 -class Q the pointwise stabilizer of $M/E - Q$ in $\text{Aut}(M)$ induces

$\text{Sym } Q$). By [6] M is monadically stable (M is cellular in the notation of Schmerl [2]).

Proposition 2 yields an easy proof of the main result of [1]. It is enough to note that in a relational language $\text{acl}(a) = \text{acl}(\emptyset) \cup \{a\}$. But this is an easy corollary of Lemma 1.

REFERENCES

1. I. M. Hodkinson and H. D. Macpherson, *Relational structures induced by their finite induced substructures*, J. Symbolic Logic **53** (1988), 222–230.
2. J. Schmerl, *Coinductive ω -categorical theories*, J. Symbolic Logic **55** (1990), 1130–1137.
3. A. H. Lachlan, *Complete theories with only universal and existential axioms*, J. Symbolic Logic **52** (1987), 698–711.
4. E. A. Palyutin, *Modeli so schetno kategorichnymi universalnymi teoriami*, Algebra i Logika **10** (1971), 23–32.
5. H. D. Macpherson, *Absolutely ubiquitous structures and ω -categorical groups*, Quart. J. Math. Oxford (2) **39** (1988), 483–500.
6. A. H. Lachlan, *ω -categorical tree-decomposable structures*, preprint, 1990.

MATHEMATICAL INSTITUTE, UNIVERSITY OF WROCLAW, PL. GRUNWALDZKI 2/4, 50-384 WROCLAW, POLAND

E-mail address: logicuwr@plwruw11.bitnet