PURELY INSEPARABLE EXTENSIONS OF $\mathbf{k}[X, Y]$

D. DAIGLE

(Communicated by Eric Friedlander)

ABSTRACT. Let $\mathbf{k}$ be a field of characteristic $p > 0$ and R a polynomial ring in two variables over $\mathbf{k}$. Define *weak variable* of R to mean an element u of R such that $u - \lambda$ is irreducible for each $\lambda \in \mathbf{k}$ and such that $R^{p^n} \subseteq \mathbf{k}[u, v]$ for some $v \in R$ and some integer $n \geq 0$. Given a weak variable u of R , consider all $v \in R$ such that $R^{p^n} \subseteq \mathbf{k}[u, v]$ for some n ; if one of these v is “absolutely smaller” than u (roughly, $\deg_X v < \deg_X u$ for all coordinate systems (X, Y) of R), we call it an R -companion of u . The main result gives a connection between the structure of a purely inseparable extension $R \supset A$, where A is a polynomial ring in two variables, and whether or not there exists a companion for each u in a suitable set of weak variables of R .

Throughout this paper, $\mathbf{k}$ is a field of characteristic $p > 0$ and R is a polynomial ring in two variables over $\mathbf{k}$ (which is abbreviated $R = \mathbf{k}^{[2]}$).

A *variable* of R is an element u of R such that $R = \mathbf{k}[u, v]$ for some $v \in R$; the set of variables of R is denoted $\text{Var}(R)$. A *coordinate system* of R is an ordered pair $(u, v) \in R \times R$ such that $R = \mathbf{k}[u, v]$; the set of coordinate systems of R is denoted $\Gamma(R)$.

One can safely assert that little is known about the subalgebras $A = \mathbf{k}^{[2]}$ of R which satisfy $R^{p^n} \subseteq A$ for n large enough—the set of all such A ’s is denoted $\Sigma(R)$. For instance, the question whether every $u \in R$ which satisfies $R/uR = \mathbf{k}^{[1]}$ is a variable of such a subalgebra A is still unanswered, after at least twelve years of existence (this is a weak version of the conjecture discussed in [7]). Questions like this one would be easier if we understood how the variables of these subalgebras look, as elements of R .

We define *weak variable* of R to mean an element u of R which is a variable of some $A \in \Sigma(R)$ and which is such that $u - \lambda$ is irreducible in R for every $\lambda \in \mathbf{k}$. A weak variable is *proper* if it is not a variable of R . Given a weak variable u of R , we discuss the following two questions:

(1) Can we describe the set $\Sigma_u(R) = \{A \in \Sigma(R) | u \in \text{Var}(A)\}$? The first section shows, in particular, that if $u \notin \mathbf{k}[R^p]$ (which is always the case if $\mathbf{k}$ is perfect) then that collection of rings is totally ordered by inclusion and, consequently, has a unique maximal element. The second section is concerned with:

Received by the editors June 8, 1992.

1991 *Mathematics Subject Classification*. Primary 13F20.

The author was supported by a grant from NSERC Canada.

(2) Consider all $v \in R$ such that $\mathbf{k}[u, v] \in \Sigma(R)$; are some of these v better than the others? More precisely, we are interested in the possible existence of a v which satisfies the following two conditions: First, $\mathbf{k}[u, v]$ is a maximal member of the collection of rings $\Sigma_u(R)$ considered in the first question. Second,

$$\deg_X v < \deg_X u \quad \text{for all } (X, Y) \in \Gamma(R).$$

We call such a v an *R-companion* of u ; it is easily seen to be a weak variable of R and, if $u \notin \mathbf{k}[R^p]$, to be almost unique (if v and v' are *R-companions* of u then $v' = av + b$ for some $a, b \in \mathbf{k}$, $a \neq 0$).

It is clear that no variable of R can have an *R-companion* but, as far as we can see, no obvious reason prevents the proper weak variables of R from having *R-companions*. The main result of the second section, Theorem (2.5), asserts the equivalence of the following two conditions on $A \in \Sigma(R)$:

(*) Every weak variable u of R such that A is a maximal member of $\Sigma_u(R)$ has an *R-companion* in A .

(**) Some variable of A belongs to R^p .

Note that, by [5] and [3], condition (**) is equivalent to the seemingly weaker:

(***) There exists $B \in \Sigma(R)$ such that $B^p \subset A \subset B$ (strictly).

Finally, we point out the following consequence of (2.5):

Corollary. *If $\mathbf{k}$ is perfect then the following are equivalent:*

- (1) *Every proper weak variable of R has an *R-companion*.*
- (2) *Every $A \in \Sigma(R)$ other than R itself has a variable in R^p .*

Conventions. If A is a ring (i.e., a commutative ring with a unity) then A^* is its set of units; if A is an integral domain then $\text{qt } A$ is its field of fractions. If $A \subseteq B$ are integral domains then the degree $[\text{qt } B : \text{qt } A]$ of the field extension $\text{qt } B / \text{qt } A$ is abbreviated $[B : A]$.

If $n > 0$ is an integer, $A = \mathbf{k}^{[n]}$ means that A is a polynomial ring in n variables over $\mathbf{k}$; by a *subalgebra* of $A = \mathbf{k}^{[n]}$ we mean a subring that contains $\mathbf{k} = \{0\} \cup A^*$.

The symbol $\subset$ is reserved for *strict* inclusion of sets.

1. STRUCTURE OF $\Sigma_u(R)$

1.1. Recall from the introduction that $\Sigma(R)$ denotes the collection of subalgebras $A = \mathbf{k}^{[2]}$ of R such that $A \supseteq R^{p^n}$ for some integer $n \geq 0$. We note that if $A, B \in \Sigma(R)$ then

- (i) $\text{qt } A \subseteq \text{qt } B \Leftrightarrow A \subseteq B$;
- (ii) if $A \subseteq B$ then $[B : A] = p^k$ for some integer $k \geq 0$ and $[B : A] = 1 \Leftrightarrow A = B$.

For each $u \in R$ we define $\Sigma_u(R) = \{A \in \Sigma(R) \mid u \text{ is a variable of } A\}$. A *weak variable* of R is an element u of R such that $\Sigma_u(R) \neq \emptyset$ and such that $u - \lambda$ is irreducible in R for every $\lambda \in \mathbf{k}$. It is easy to see that if $u \in R$ satisfies $\Sigma_u(R) \neq \emptyset$ then $u = aw^{p^k} + b$ for some weak variable w of R , some $a \in \mathbf{k}^*$,

$b \in \mathbf{k}$, and some integer $k \geq 0$. A weak variable u of R is *proper* if it is not a variable of R .

Remark. Suppose $\mathbf{k}$ is perfect. Then a weak variable of R is simply an irreducible element $u \in R$ such that $\Sigma_u(R) \neq \emptyset$, and if u is any element of R which satisfies $\Sigma_u(R) \neq \emptyset$ then $u = w^{p^k}$ for some weak variable w of R and some integer $k \geq 0$.

Before examining the structure of $\Sigma_u(R)$ we gather some elementary observations into a lemma. These probably exist in the literature, but we could not find appropriate references.

1.2. Lemma. (1) Let T be a set of indeterminates over $\mathbf{k}$, $\mathbf{k}_0$ a subfield of $\mathbf{k}$, and S a subset of $\mathbf{k}_0[T]$. Then $\mathbf{k}_0[T] \cap \mathbf{k}[S] = \mathbf{k}_0[S]$.

(2) If A is a subalgebra of R then $R^p \cap \mathbf{k}[A^p] = A^p$.

(3) Let $A = \mathbf{k}[u, v] = \mathbf{k}^{[2]}$ be a subalgebra of R such that $v \notin \mathbf{k}[R^p]$. Then $R^p \cap \mathbf{k}[u^p, v] = A^p$.

(4) Let $A = \mathbf{k}^{[2]}$ and $B = \mathbf{k}^{[2]}$ be subalgebras of R . If $\exists u, v, s, t$ such that $A = \mathbf{k}[u, v]$, $B = \mathbf{k}[s, t]$, and $\mathbf{k}[u^p, v] = \mathbf{k}[s^p, t] \not\subseteq \mathbf{k}[R^p]$ then $A = B$.

Proof. (1) Let $\mathcal{M}$ be the set of monomials, i.e., the set of finite products of elements of $\{1\} \cup T$. Then $\mathcal{M}$ is a basis for the $\mathbf{k}$ -vector space $\mathbf{k}[T]$ and also for the $\mathbf{k}_0$ -vector space $\mathbf{k}_0[T]$.

Note that if $\bar{S} = \mathbf{k}_0[S]$ then $\mathbf{k}_0[\bar{S}] = \mathbf{k}_0[S]$ and $\mathbf{k}[\bar{S}] = \mathbf{k}[S]$; in particular, we may assume that $\mathbf{k}[S] = \text{Span}_{\mathbf{k}} S$.

Clearly, $\mathbf{k}_0[T] \cap \mathbf{k}[S] \supseteq \mathbf{k}_0[S]$. To prove the other inclusion, consider $\xi \in \mathbf{k}_0[T] \cap \mathbf{k}[S]$ and note that $\xi = \sum_{j=1}^n \xi_j s_j$ for some $\xi_1, \dots, \xi_n \in \mathbf{k}$ and $s_1, \dots, s_n \in S$. Since $s_1, \dots, s_n \in \mathbf{k}_0[T] = \text{Span}_{\mathbf{k}_0} \mathcal{M}$, we may select a finite subset $\mathcal{B} = \{M_1, \dots, M_m\}$ of $\mathcal{M}$ such that $s_1, \dots, s_n \in \text{Span}_{\mathbf{k}_0} \mathcal{B}$. Note that $\xi = \sum_{j=1}^n \xi_j s_j \in \text{Span}_{\mathbf{k}} \mathcal{B}$ and $\text{Span}_{\mathbf{k}} \mathcal{B} \cap \mathbf{k}_0[T] = \text{Span}_{\mathbf{k}_0} \mathcal{B}$ imply $\xi \in \text{Span}_{\mathbf{k}_0} \mathcal{B}$. Now consider the $\mathbf{k}_0$ -linear map $\sigma_0: \mathbf{k}_0^n \rightarrow \text{Span}_{\mathbf{k}_0} \mathcal{B}$ which maps the standard basis of $\mathbf{k}_0^n$ on $s_1, \dots, s_n$, and its unique extension to a $\mathbf{k}$ -linear map $\sigma: \mathbf{k}^n \rightarrow \text{Span}_{\mathbf{k}} \mathcal{B}$. The equality $\sigma_0(X_1, \dots, X_n) = \xi$ can be viewed as a system of m linear equations in the unknowns $X_1, \dots, X_n$; this system is consistent by elementary linear algebra and the fact that $\sigma(\xi_1, \dots, \xi_n) = \sum_{j=1}^n \xi_j s_j = \xi$. Hence for some $\xi'_1, \dots, \xi'_n \in \mathbf{k}_0$ we have

$$\xi = \sigma_0(\xi'_1, \dots, \xi'_n) = \sum_{j=1}^n \xi'_j s_j \in \mathbf{k}_0[S].$$

(2) Write $R = \mathbf{k}[X, Y]$, and use the first part with $\mathbf{k}_0 = \mathbf{k}^p$, $T = \{X^p, Y^p\}$, and $S = A^p$:

$$R^p \cap \mathbf{k}[A^p] = \mathbf{k}_0[T] \cap \mathbf{k}[S] = \mathbf{k}_0[S] = \mathbf{k}^p[A^p] = A^p.$$

(3) $R^p \cap \mathbf{k}[u^p, v] \supseteq A^p$ is trivial. Let $\xi \in R^p \cap \mathbf{k}[u^p, v]$, and write $\xi = F(u^p, v)$, where $F \in \mathbf{k}[T_1, T_2]$ (T_1, T_2 indeterminates over $\mathbf{k}$). Write $R = \mathbf{k}[X_1, X_2]$; then for some $i \in \{1, 2\}$ the derivation $\partial/\partial X_i: R \rightarrow R$ satisfies $\partial v/\partial X_i \neq 0$. Since $\xi \in R^p$,

$$0 = \frac{\partial \xi}{\partial X_i} = F_{T_1}(u^p, v) \frac{\partial u^p}{\partial X_i} + F_{T_2}(u^p, v) \frac{\partial v}{\partial X_i} = F_{T_2}(u^p, v) \frac{\partial v}{\partial X_i},$$

whence $F_{T_2}(u^p, v) = 0$, i.e., $F \in \mathbf{k}[T_1, T_2^p]$. Thus, $\xi \in \mathbf{k}[u^p, v^p] = \mathbf{k}[A^p]$, and the conclusion follows from (2).

(4) Since $\mathbf{k}[u^p, v] \not\subseteq \mathbf{k}[R^p]$, we have $v \notin \mathbf{k}[R^p]$, so $R^p \cap \mathbf{k}[u^p, v] = A^p$ by (3). It follows that $s \in A$, since $s^p \in R^p \cap \mathbf{k}[u^p, v]$ is clear. Hence $B \subseteq A$ and we are done by symmetry. $\square$

We now turn to the set $\Sigma_u(R)$. The first two results are easy and useful.

1.3. Lemma. *Let $u \in R$ be such that $\Sigma_u(R) \neq \emptyset$, and let $A, B \in \Sigma_u(R)$ be such that $A \subseteq B$. Then $A = \mathbf{k}[B^{p^s}, u]$ for some integer $s \geq 0$, where $\mathbf{k}[B^{p^s}, u]$ denotes the subalgebra of R generated by the set $B^{p^s} \cup \{u\}$.*

Proof. Let v and w be such that $A = \mathbf{k}[u, v]$ and $B = \mathbf{k}[u, w]$, and let $K = \mathbf{k}(u, v)$ and $L = \mathbf{k}(u, w)$. Since $w \notin B^p \supseteq A^p$ and $w^{p^n} \in R^{p^n} \subseteq A$ for n large enough, we may consider $s \geq 0$ such that $w^{p^s} \in A \setminus A^p$. Now $w^{p^s} \in K \setminus K^p$ as well (for $K^p \cap A = A^p$), so $[L : K] = [K[w] : K] = p^s = [L : \mathbf{k}(u, w^{p^s})]$. Since $\mathbf{k}(u, w^{p^s}) \subseteq K \subseteq L$, we conclude that $K = \mathbf{k}(u, w^{p^s})$, which means that the rings A and $\mathbf{k}[B^{p^s}, u] = \mathbf{k}[u, w^{p^s}]$ have the same field of fractions. Since these two rings belong to $\Sigma(R)$, they must be equal. $\square$

1.4. Proposition. *If u is a weak variable of R such that $u \notin \mathbf{k}[R^p]$ then $\Sigma_u(R)$ is totally ordered by inclusion. In particular, $\Sigma_u(R)$ has a unique maximal element.*

Proof. Let $A, B \in \Sigma_u(R)$. For $i = 0, 1, 2, \dots$, define $A_i = \mathbf{k}[A^{p^i}, u]$ and $B_i = \mathbf{k}[B^{p^i}, u]$; note that $A_i, B_i \in \Sigma_u(R)$ and that $A_i \not\subseteq \mathbf{k}[R^p]$ and $B_i \not\subseteq \mathbf{k}[R^p]$. If i is large enough then $A_i \subseteq B$, so, by (1.3), $A_i = B_s$ for some $s \geq 0$. Hence, $E \neq \emptyset$, where E is the set of pairs (i, j) of nonnegative integers such that $A_i = B_j$. If $(i, j) \in E$, $i > 0$, and $j > 0$ then 1.2(4) implies that $(i-1, j-1) \in E$. Thus E contains an element (i, j) such that $i = 0$ or $j = 0$, which means that $A \subseteq B$ or $B \subseteq A$, as desired. $\square$

When $u \in \mathbf{k}[R^p]$, the structure of $\Sigma_u(R)$ can be more complicated. To describe it, we will need some definitions.

1.5. If $u \in R$ then $\text{Max } \Sigma_u(R)$ denotes the set of maximal members of $\Sigma_u(R)$.

1.6. Let $\vec{\Sigma}(R)$ be the directed graph whose set of vertices is $\Sigma(R)$ and where, given $A, B \in \Sigma(R)$, we draw an edge $A \rightarrow B$ if and only if there exist $x, y \in B$ such that $B = \mathbf{k}[x, y]$ and $A = \mathbf{k}[x, y^p]$. Note that there is at most one edge from A to B and no edge from A to itself. It is not known whether $\vec{\Sigma}(R)$ is connected, but it is clear that it contains lots of cycles. If $u \in R$ is such that $\Sigma_u(R) \neq \emptyset$ then let $\vec{\Sigma}_u(R)$ be the subgraph of $\vec{\Sigma}(R)$ generated by $\Sigma_u(R)$. Observe that (1.3) implies the following:

- (i) If $A, B \in \Sigma_u(R)$ then there is an edge $A \rightarrow B$ if and only if there exists v such that $B = \mathbf{k}[u, v]$ and $A = \mathbf{k}[u, v^p]$.
- (ii) If $A_1, A_2, B \in \Sigma_u(R)$ and $A_1 \rightarrow B \leftarrow A_2$ then $A_1 = A_2$.
- (iii) $\vec{\Sigma}_u(R)$ is connected.

Hence $\vec{\Sigma}_u(R)$ is a tree, by (ii) and (iii). An element A of $\Sigma_u(R)$ is a *branch point* (of $\vec{\Sigma}_u(R)$) if there exist $B_1, B_2 \in \Sigma_u(R)$ such that $B_1 \neq B_2$ and $B_1 \leftarrow A \rightarrow B_2$. Clearly, $\Sigma_u(R)$ is totally ordered by inclusion $\Leftrightarrow \text{Max } \Sigma_u(R)$ has only one element $\Leftrightarrow \vec{\Sigma}_u(R)$ has no branch point.

1.7. Example. Suppose that $\mathbf{k}$ is not perfect, choose $\theta \in \mathbf{k} \setminus \mathbf{k}^p$ and $(X, Y) \in \Gamma(R)$, and let $u = X^{p^2} + \theta$. Then $\Sigma_u(R) \neq \emptyset$ and u is an irreducible element of R but not a weak variable of R . Consider the $\mathbf{k}$ -vector space $\mathbf{k}[X]$ and its subspace $\mathbf{k}[X^p]$; let $E \subset \mathbf{k}[X]$ be a set of representative elements for the cosets of $\mathbf{k}[X^p]$ in $\mathbf{k}[X]$, except that E contains no representative element for the coset $\mathbf{k}[X^p]$ itself. Consider the following rings:

- $A_n = \mathbf{k}[X^{p^2}, Y^{p^n}]$, for $n \geq 0$;
- $A_{n,f} = \mathbf{k}[X^{p^2}, Y^{p^{n-1}} + f^p]$, for $n \geq 1$ and $f \in E$;
- $A_{n,f,g} = \mathbf{k}[X^{p^2}, Y^{p^{n-2}} + f + g^p]$, for $n \geq 2$ and $f, g \in E$.

Now all these rings belong to $\Sigma_u(R) = \Sigma_{X^{p^2}}(R)$ and are distinct. If $n \geq 2$ then we have $A_n \rightarrow A_{n,f} \rightarrow A_{n,f,g} \in \text{Max } \Sigma_u(R)$ for all $f, g \in E$, so that A_n and $A_{n,f}$ are branch points of $\tilde{\Sigma}_u(R)$. We conclude that, for each integer $n \geq 1$, each one of the sets $\{A \in \text{Max } \Sigma_u(R) \mid [R:A] = p^n\}$ and $\{A \in \Sigma_u(R) \mid [R:A] = p^n \text{ and } A \text{ is a branch point}\}$ has cardinality equal to $|\mathbf{k}|$.

The above example shows that $\tilde{\Sigma}_u(R)$ can have lots of branch points when u is not a weak variable of R . In contrast, we will prove:

1.8. Proposition. *If u is a weak variable of R then $\tilde{\Sigma}_u(R)$ has at most one branch point.*

1.9. Example. Suppose that $\mathbf{k}$ is not perfect, choose $\theta \in \mathbf{k} \setminus \mathbf{k}^p$ and $(X, Y) \in \Gamma(R)$, and let $n > 0$ be an integer. Then $u = X^{p^n} + \theta Y^{p^n}$ is a weak variable of R . It is not difficult to see that $\mathbf{k}[X^{p^n}, Y^{p^n}]$ is a branch point (the only one) of $\tilde{\Sigma}_u(R)$ and that if we define the sets

$$E_r = \{A \in \text{Max } \Sigma_u(R) \mid [R:A] = p^r\}, \quad r = 0, 1, 2, \dots,$$

then $|E_r| = |\mathbf{k}|$ when $n \leq r \leq 2n - 1$ and E_r is empty otherwise.

Some notation and observations will be needed:

1.10. Let $\text{Der}(\mathbf{k}, \mathbf{k})$ be the set of all derivations $\mathbf{k} \rightarrow \mathbf{k}$. Recall that an element ρ of $\mathbf{k}$ is in $\mathbf{k}^p$ if and only if $D\rho = 0$ for all $D \in \text{Der}(\mathbf{k}, \mathbf{k})$ (see for instance [6, p. 183]).

Suppose that $(X, Y) \in \Gamma(R)$ has been chosen. Then each $D \in \text{Der}(\mathbf{k}, \mathbf{k})$ determines a derivation $\tilde{D}: R \rightarrow R$ defined by $\sum a_{ij} X^i Y^j \mapsto \sum D(a_{ij}) X^i Y^j$ (where $a_{ij} \in \mathbf{k}$). Given a weak variable u of R such that $u \in \mathbf{k}[R^p]$, we clearly have

$$\tilde{D}u \notin \mathbf{k} \quad \text{for some } D \in \text{Der}(\mathbf{k}, \mathbf{k}).$$

Proof of 1.8. Let u be a weak variable of R , and suppose that $\tilde{\Sigma}_u(R)$ has at least one branch point. Then u is proper and, by 1.4, $u \in \mathbf{k}[R^p]$. Choose $(X, Y) \in \Gamma(R)$. We will show that if $A \in \Sigma_u(R)$ is a branch point of $\tilde{\Sigma}_u(R)$ then

$$(1) \quad \{\mathbf{k}[u, \tilde{D}u] \mid D \in \text{Der}(\mathbf{k}, \mathbf{k})\} \cap \Sigma(R) = \{A\},$$

where $\tilde{D}$ is defined as in 1.10. Observe that the proofs of the facts labeled 2.6, 2.7, and 2.8, in the next section, are independent of this proposition. So we may apply those facts and conclude that $\deg_X u > 0$ and that u is almost monic in X . Replacing if necessary u by a constant multiple of it, we may therefore assume that u is monic in X , so that $\deg_X \tilde{D}u < \deg_X u$ for all D .

Suppose that A is a branch point of $\tilde{\Sigma}_u(R)$. Then there exist distinct elements $\mathbf{k}[u, v]$ and $\mathbf{k}[u, w]$ of $\Sigma_u(R)$ such that $\mathbf{k}[u, v^p] = A = \mathbf{k}[u, w^p]$, and consequently

$$(2) \quad w^p = av^p + P(u),$$

for some $a \in \mathbf{k}^*$ and $P \in \mathbf{k}[T]$ (where T is an indeterminate). This implies that

$$(3) \quad 0 = Dav^p + P^{(D)}(u) + P'(u)\tilde{D}u \quad \text{for all } D \in \text{Der}(\mathbf{k}, \mathbf{k}).$$

We claim that $P' \neq 0$. Indeed, if $P' = 0$ then (3) and the fact that u and v are algebraically independent over $\mathbf{k}$ imply that $Da = 0$ and $P^{(D)} = 0$ for all D . Thus $a \in \mathbf{k}^p$, $P \in (\mathbf{k}[T])^p$, and (2) yields $\mathbf{k}[u, v] = \mathbf{k}[u, w]$, a contradiction.

Let $D \in \text{Der}(\mathbf{k}, \mathbf{k})$, and let us determine the ring $\mathbf{k}[u, \tilde{D}u]$. Note that

$$(4) \quad \tilde{D}u = -\frac{Dav^p + P^{(D)}(u)}{P'(u)},$$

so in particular $\tilde{D}u \in \mathbf{k}(u, v^p)$; since $\tilde{D}u \in R$, it is integral over $\mathbf{k}[u, v^p]$ and consequently

$$(5) \quad \tilde{D}u \in \mathbf{k}[u, v^p].$$

Case $Da = 0$. Then (4) and (5) say that $\tilde{D}u \in \mathbf{k}(u) \cap \mathbf{k}[u, v^p] = \mathbf{k}[u]$; together with $\deg_X \tilde{D}u < \deg_X u$, this gives $\tilde{D}u \in \mathbf{k}$. In particular, $\mathbf{k}[u, \tilde{D}u] = \mathbf{k}[u] \notin \Sigma(R)$ in this case.

Observe that 1.10 and the above paragraph imply that some $D \in \text{Der}(\mathbf{k}, \mathbf{k})$ satisfies $Da \neq 0$.

Case $Da \neq 0$. By (3) and (5), $P'(u)$ divides $Dav^p + P^{(D)}(u)$ in the ring $\mathbf{k}[u, v^p]$. Hence $P'(u) \in \mathbf{k}^*$ and, by (4), $\mathbf{k}[u, \tilde{D}u] = \mathbf{k}[u, v^p] = A$.

This establishes (1) and the proof of the proposition is complete. $\square$

2. COMPANIONS OF WEAK VARIABLES

2.1. Definition. Let u be a weak variable of R . An R -cogenerator of u is an element v of R such that $\mathbf{k}[u, v] \in \text{Max } \Sigma_u(R)$; an R -companion of u is an R -cogenerator v which satisfies $\deg_X v < \deg_X u$ for all $(X, Y) \in \Gamma(R)$.

If u is a weak variable of R then, clearly, u has R -cogenerators; moreover, we note:

2.2. Every R -cogenerator of u is a weak variable of R .

2.3. Definition. If $A \in \Sigma(R)$ then $W(A, R)$ denotes the set of weak variables u of R which satisfy $\text{Max } \Sigma_u(R) \ni A$.

For later use, we record:

2.4. If $A, B \in \Sigma(R)$ satisfy $A \subseteq B$ then $W(A, R) \subseteq W(B, R)$.

We can now state the main result of this section:

2.5. Theorem. Given $A \in \Sigma(R)$, the following are equivalent:

- (1) Every member of $W(A, R)$ has an R -companion which belongs to A .
- (2) Some variable of A belongs to R^p .

Remark. If $\mathbf{k}$ is perfect then the first condition of the theorem is equivalent to:

(1') *Every member of $W(A, R)$ has an R -companion.*

The implication (1) $\Rightarrow$ (2) is trivial and we dispose of it right away. Suppose that condition (1) holds. The reader will verify that, to prove that (2) holds, it suffices to show that $W(A, R)$ is a *proper* subset of $\text{Var}(A)$. In particular, we may assume that $W(A, R) \neq \emptyset$. Choose $(X, Y) \in \Gamma(R)$, and then select an element u of $W(A, R)$ which minimizes $\deg_X u$. By (1), u has an R -companion v which belongs to A ; since $\deg_X v < \deg_X u$, we have $v \notin W(A, R)$. But $v \in A$ implies that $\mathbf{k}[u, v] = A$, so $v \in \text{Var}(A)$ and we are done.

The rest of this paper is devoted to the proof that (2) $\Rightarrow$ (1). The heart of the proof is 2.11, but some preliminaries are necessary.

Recall that an irreducible element u of R is said to *have one place at infinity* if only one valuation ring $\mathcal{O}$ of $\text{qt}(R/uR)$ over $\mathbf{k}$ does not contain R/uR ; in that case $\mathcal{O}$ is called the place of u at infinity. Moreover, if the residue field of $\mathcal{O}$ coincides with $\mathbf{k}$ (resp., is purely inseparable over $\mathbf{k}$) then u is said to have one *rational* (resp., *purely inseparable*) place at infinity. We note:

2.6. *If u is a weak variable of R then u has one purely inseparable place at infinity.*

Proof. Choose an element $A = \mathbf{k}[u, v]$ of $\Sigma_u(R)$, and let $p^n = [R: A]$. Then $R^{p^n} \subseteq A \subseteq R$, and consequently $(R/uR)^{p^n} \subseteq A/uA \subseteq R/uR$. Since $A/uA = \mathbf{k}^{[1]}$ has one place $\mathcal{O}_0$ at infinity and $\text{qt}(R/uR)$ is purely inseparable over $\text{qt}(A/uA)$, it follows that R/uR has one place $(\mathcal{O}, \kappa)$ at infinity (where κ stands for the residue field of $\mathcal{O}$). Now $\mathcal{O}^{p^n} \subseteq \mathcal{O}_0$ is clear, so κ^{p^n} is contained in the residue field of $\mathcal{O}_0$, which is $\mathbf{k}$. Hence κ is purely inseparable over $\mathbf{k}$, as claimed. $\square$

Remarks. (1) One can see that $[\kappa: \mathbf{k}]$ divides $[R: A]$ for all $A \in \Sigma_u(R)$ (notation as in the proof), but this will not be needed.

(2) If $\mathbf{k}$ is perfect and u is a weak variable of R then u has one *rational* place at infinity, by 2.6. Moreover, one can see that the integral closure of R/uR in its field of fractions is $\mathbf{k}^{[1]}$ (u is a so-called “polynomial curve”).

Curves with one rational place at infinity have several interesting properties (see [1, 2, 4, 8], among other references), and some of those properties are in fact shared by all curves with one (not necessarily rational) place at infinity. In particular, we note that the result (11.19) of [1] holds in general:

2.7. *Let F be a field and $A = F^{[2]}$, and suppose that $u \in A$ has one place at infinity. Let $\gamma = (X, Y) \in \Gamma(A)$ and $(m, n) = \text{bideg}_\gamma u$. Then every $(i, j) \in \text{Supp}_\gamma u$ satisfies $ni + mj \leq mn$ (we say that the support of u has triangular shape). In particular, if $n > 0$ then u is almost monic in Y ; i.e., if we regard u as a polynomial in Y with coefficients in $F[X]$ then the leading coefficient of u is in F^* .*

Here, $\text{bideg}_\gamma u = (\deg_X u, \deg_Y u)$ and $\text{Supp}_\gamma u = \{(i, j) | a_{ij} \neq 0\}$ where $u = \sum a_{ij} X^i Y^j$ and $a_{ij} \in F$. Although we do not know a reference for 2.7 we will only prove the special case which interests us, because in this situation we can easily reduce to the rational case: If $u \in R$ satisfies $\Sigma_u(R) \neq \emptyset$, then $\Sigma_u(\bar{R}) \neq \emptyset$ as well, where $\bar{R} = \bar{\mathbf{k}} \otimes R$ and $\bar{\mathbf{k}}$ is the perfect closure of $\mathbf{k}$.

Consequently, $u = w^{p^k}$ for some weak variable w of $\bar{R}$ and some integer $k \geq 0$, and $w \in \bar{R}$ has one rational place at infinity. Thus 2.7 holds for u , since it holds for w .

Observe that if u is a weak variable of R which satisfies $\deg_X u = 1$ for some $\gamma = (X, Y) \in \Gamma(R)$ then u is a variable of R (by 2.7, $u = aX + f(Y)$ for some $a \in \mathbf{k}^*$ and $f(Y) \in \mathbf{k}[Y]$). If $\deg_X u = 0$ then (notation as above) $u = w^{p^k}$ where $w \in \bar{R} = \bar{\mathbf{k}}[X, Y]$ has one rational place at infinity and must satisfy $\text{bideg}_\gamma w = (0, 1)$, so $u = aY^{p^k} + b$ for some $a \in \mathbf{k}^*$, $b \in \mathbf{k}$, and $k \geq 0$; since $u - b$ is irreducible, $k = 0$ and u is a variable in this case too. Hence:

2.8. *If u is a proper weak variable of R then $\deg_X u > 1$ for all $(X, Y) \in \Gamma(R)$.*

The following device will be needed in the proof of 2.11.

2.9. Suppose that $u, v \in R$ have one place at infinity and that u satisfies $\deg_X u > 0$ for all $(X, Y) \in \Gamma(R)$ (e.g., u is a proper weak variable of R). For each $\gamma = (X, Y) \in \Gamma(R)$ we define the positive rational number

$$M_\gamma(v/u, R) = \max \left(\frac{\deg_X v}{\deg_X u}, \frac{\deg_Y v}{\deg_Y u} \right).$$

We also define $M(v/u, R) = \sup_{\gamma \in \Gamma(R)} M_\gamma(v/u, R)$.

A pair (a, b) of nonnegative integers is *nonprincipal* if either $ab = 0$ or both $a \nmid b$ and $b \nmid a$ hold.

2.10. **Lemma.** *Let $u, v \in R$ be as in 2.9. If $\gamma \in \Gamma(R)$ is such that $\text{bideg}_\gamma u$ is nonprincipal then $M_\gamma(v/u, R) = M(v/u, R)$.*

Proof. Write $\gamma = (X, Y)$, $(m, n) = \text{bideg}_\gamma u$, $(r, s) = \text{bideg}_\gamma v$, and $M = M_\gamma(v/u, R) = \max(r/m, s/n)$. It suffices to show that $\deg_X v(f, g) / \deg_X u(f, g) \leq M$ whenever $(f, g) \in \Gamma(R)$. Let $\alpha = \deg_X f$ and $\beta = \deg_X g$.

If $(\alpha, \beta) = (1, 0)$ then $\deg_X v(f, g) / \deg_X u(f, g) = r/m \leq M$. If $(\alpha, \beta) = (0, 1)$ then $\deg_X v(f, g) / \deg_X u(f, g) = s/n \leq M$.

Now suppose (α, β) is neither $(1, 0)$ nor $(0, 1)$. By basic properties of coordinate systems of R we have $\alpha\beta \neq 0$ and either $\alpha|\beta$ or $\beta|\alpha$. Since, by 2.7, $\text{Supp}_\gamma u$ and $\text{Supp}_\gamma v$ have triangular shapes, $\deg_X u(f, g) \leq \max(m\alpha, n\beta)$ and $\deg_X v(f, g) \leq \max(r\alpha, s\beta)$. Moreover, the fact that (α, β) is principal and that (m, n) is nonprincipal implies that $m\alpha \neq n\beta$ and consequently that $\deg_X u(f, g) = \max(m\alpha, n\beta)$. Hence

$$\deg_X v(f, g) \leq \max(r\alpha, s\beta) \leq \max(Mm\alpha, Mn\beta) = M \deg_X u(f, g),$$

as asserted. $\square$

2.11. **Proposition.** *If $A = \mathbf{k}[u, v] \in \Sigma(R)$, $u \in W(A, R)$, and $v \notin W(A, R)$ then v is an R -companion of u .*

Proof. Note that $A \neq R$ (since $W(A, R) \neq \text{Var}(A)$), so u is a proper weak variable of R . Since v is clearly an R -cogenerator of u , 2.2 and the assumption $v \notin W(A, R)$ imply that $A \notin \text{Max } \Sigma_v(R)$. Thus 1.3 implies that $A = \mathbf{k}[v, w^p]$ for some $w \in R$ and consequently

$$(1) \quad au = w^p + P(v),$$

for some $a \in \mathbf{k}^*$ and some polynomial $P \in \mathbf{k}[T]$. Observe that P and a are completely determined by w , and assume that w has been chosen so as to minimize $d = \deg_T P$.

Clearly, if $d = 0$ then $u - \lambda$ is reducible for some $\lambda \in \mathbf{k}$, and if $d = 1$ then $A = \mathbf{k}[u, w^p]$, so $A \notin \text{Max}_{\Sigma_u}(R)$. Hence the fact that $u \in W(A, R)$ implies that $d > 1$.

Claim 1. If $p \mid d$ then v is an R -companion of u .

To see this, suppose $p \mid d$ and observe that the leading coefficient of P is not in $\mathbf{k}^p$ (otherwise, we could replace w by $w + bv^{d/p}$, for suitable $b \in \mathbf{k}^*$, and obtain a P of lower degree, contradicting the minimality of d). Now let $(X, Y) \in \Gamma(R)$, and given $F \in \mathbf{k}[X, Y]$ let $\text{lc}_X F$ denote the leading coefficient of F , when F is regarded as a polynomial in X with coefficients in $\mathbf{k}[Y]$. It suffices to show that $\deg_X u > \deg_X v$. This is trivial if $\deg_X v = 0$, because $X^{p^n} \in \mathbf{k}[u, v]$ for some n . Suppose that $\deg_X v \neq 0$; since $\text{lc}_X P(v) \notin \mathbf{k}[Y]^p$ is clear, we have $\text{lc}_X P(v) \neq -\text{lc}_X(w^p) \in \mathbf{k}[Y]^p$ and consequently

$$\deg_X u = \max(\deg_X(w^p), \deg_X P(v)) \geq d \deg_X v > \deg_X v,$$

which proves Claim 1.

From now on we assume that $p \nmid d$. The proof goes by induction on n , where $[R : A] = p^n$.

If $n = 1$ then write $\gamma = (v, w) \in \Gamma(R)$ and observe that $\text{bideg}_\gamma u = (d, p)$ is nonprincipal. So, by 2.10, $M(v/u, R) = M_\gamma(v/u, R) = \max(1/d, 0/p) \leq 1/2$ and v is an R -companion of u .

Suppose $n > 1$, and let $\gamma = (X, Y) \in \Gamma(R)$. Again, we have to show that $\deg_X u > \deg_X v$. Observe that the derivative P' has degree $d - 1 > 0$ (for $p \nmid d$) and that (1) implies that

$$au_X = P'(v)v_X \quad \text{and} \quad au_Y = P'(v)v_Y.$$

In particular, $u_X = 0 \Leftrightarrow v_X = 0$ and $u_Y = 0 \Leftrightarrow v_Y = 0$.

If $u_X \neq 0$ then $\deg_X u > \deg_X u_X \geq \deg_X v$ and we are done. So we may assume that $u_X = 0$ (thus $v_X = 0$).

Suppose $u_Y \neq 0$ (and $v_Y \neq 0$). Since $[\mathbf{k}[v, w] : A] = p < p^n = [R : A]$, we have $\mathbf{k}[v, w] \neq R$, which implies the nullity of the jacobian: $0 = v_X w_Y - v_Y w_X = -v_Y w_X$. Whence $u, v, w \in B$, where we define $B = \mathbf{k}[X^p, Y]$. Now $A \in \Sigma(B)$, $u \in W(A, B)$ (by 2.4), and $v \notin W(A, B)$ (because $A \subset \mathbf{k}[v, w] \in \Sigma_v(B)$), so by the inductive hypothesis v is a B -companion of u . Since $(X^p, Y) \in \Gamma(B)$, it follows that $\deg_{(X^p)} u > \deg_{(X^p)} v$ and hence that $\deg_X u > \deg_X v$, as desired.

The last case to consider is $u_Y = 0$, i.e., $u, v \in \mathbf{k}[X^p, Y^p] = \mathbf{k}[R^p]$. Let us proceed by contradiction, and assume that $\deg_X u \leq \deg_X v$. Observe that $1 < \deg_X u \leq \deg_X v$ by 2.8, so u and v are almost monic in X by 2.7 and 2.6. Multiplying if necessary u, v , and the coefficients of P by elements of $\mathbf{k}^*$, we may therefore assume that

$$(2) \quad v \text{ is monic in } X \text{ and } u = w^p + P(v).$$

In view of 1.10, it is enough to prove

Claim 2. $\tilde{D}(v) \in \mathbf{k}$, for all $D \in \text{Der}(\mathbf{k}, \mathbf{k})$.

Let $D : \mathbf{k} \rightarrow \mathbf{k}$ be a derivation such that $\tilde{D}v \neq 0$. Let $n = \deg_X v$ and $r = \deg_X(\tilde{D}v)$; we have $r < n$, since v is monic in X . Let $P^{(D)} \in \mathbf{k}[T]$

denote the polynomial $\sum D(b_i)T^i$, where $P = \sum b_i T^i$ and $b_i \in \mathbf{k}$. Then, by (2),

$$(3) \quad \tilde{D}u = P^{(D)}(v) + P'(v)\tilde{D}v.$$

We claim that $D(b_d)=0$ and $r=0$. Indeed, if $D(b_d) \neq 0$ then $\deg_X(P^{(D)}(v)) = dn > (d-1)n+r = \deg_X(P'(v)\tilde{D}v)$, so the right-hand side of (3) has X -degree dn ; this is absurd, since $dn > n \geq \deg_X u \geq \deg_X \tilde{D}u$. Hence $D(b_d) = 0$. If $r > 0$ then $\deg_X(P'(v)\tilde{D}v) = (d-1)n+r > (d-1)n \geq \deg_X P^{(D)}(v)$, so the right-hand side of (3) has X -degree $(d-1)n+r$; this is impossible because $(d-1)n+r > n \geq \deg_X \tilde{D}u$. Hence $r = 0$.

So $\deg_X P^{(D)}(v) \leq (d-1)n = \deg_X(P'(v)\tilde{D}v)$, and consequently the coefficient of $X^{(d-1)n}$ in the right-hand side of (3) is $D(b_{d-1}) + db_d\tilde{D}v$. This means that

$$D(b_{d-1}) + db_d\tilde{D}v = (\text{coefficient of } X^{(d-1)n} \text{ in } \tilde{D}u) = \tilde{D}(u_{(d-1)n}),$$

where $u = \sum u_i X^i$ and $u_i \in \mathbf{k}[Y^p]$. Since u is almost monic in X and $(d-1)n \geq n \geq \deg_X u$, we have $u_{(d-1)n} \in \mathbf{k}$, so $D(b_{d-1}) + db_d\tilde{D}v = \tilde{D}(u_{(d-1)n}) \in \mathbf{k}$. We conclude that $\tilde{D}v \in \mathbf{k}$, which completes the proof of Claim 2 and also that of the proposition. $\square$

2.12. Definition. Let $A \in \Sigma(R)$ and $\gamma = (X, Y) \in R \times R$ be such that $A \subseteq \mathbf{k}[X, Y]$. An (A, R, γ) -sequence is a sequence $u^* = (u_0, \dots, u_k)$ of elements of $W(A, R)$ such that $k \geq 0$ and such that the following condition holds if $k > 0$:

$$A = \mathbf{k}[u_{i-1}, u_i] \text{ and } \deg_X u_{i-1} > \deg_X u_i \text{ for } 1 \leq i \leq k.$$

An (A, R, γ) -sequence $(u_0, \dots, u_k)$ is *maximal* if no $u_{k+1} \in W(A, R)$ is such that $(u_0, \dots, u_{k+1})$ is an (A, R, γ) -sequence. Clearly, every (A, R, γ) -sequence $(u_0, \dots, u_k)$ can be extended to a maximal one, $(u_0, \dots, u_k, \dots, u_l)$.

2.13. Lemma. Let $A, B \in \Sigma(R)$ be such that $A \subset B$, let $\gamma \in \Gamma(B) \cup \Gamma(R)$, and let $(u_0, \dots, u_k)$ be an (A, R, γ) -sequence. If some $u_{k+1} \in A$ is both a B -companion and an R -companion of u_k then u_1 is both a B -companion and an R -companion of u_0 .

Proof. We may assume that $k > 0$. We have $A = \mathbf{k}[u_{k-1}, u_k]$, and $u_{k-1} \in W(A, R) \subseteq W(A, B)$ implies that u_k is both a B -cogenerator and an R -cogenerator of u_{k-1} . Clearly, it suffices to prove:

Claim. u_k is both a B -companion and an R -companion of u_{k-1} .

Consider $u_{k+1} \in A$, as in the statement. Since $\Sigma_{u_k}(R) \ni A \supseteq \mathbf{k}[u_k, u_{k+1}] \in \text{Max } \Sigma_{u_k}(R)$, we see that $\mathbf{k}[u_k, u_{k+1}] = A = \mathbf{k}[u_{k-1}, u_k]$ and consequently

$$u_{k-1} = \lambda u_{k+1} + P(u_k),$$

for some $\lambda \in \mathbf{k}^*$ and some $P \in \mathbf{k}[T]$ (where T is an indeterminate). Let $d = \deg_T P$; if we write $\gamma = (X_1, Y_1)$ then

$$\deg_{X_1} u_{k-1} > \deg_{X_1} u_k > \deg_{X_1} u_{k+1},$$

and these inequalities imply that $d > 1$. Consequently, for each $(X, Y) \in \Gamma(B) \cup \Gamma(R)$ we have

$$\deg_X u_{k-1} = \deg_X(\lambda u_{k+1} + P(u_k)) = d \deg_X u_k > \deg_X u_k,$$

which means that u_k is both a B -companion and an R -companion of u_{k-1} , as claimed. $\square$

2.14. Corollary. *Let $(X, Y) \in \Gamma(R)$ and $A = \mathbf{k}[X, Y^p]$. Then every element of $W(A, R)$ has an R -companion which belongs to A .*

Proof. Let $u_0 \in W(A, R)$; note that (u_0) is an (A, R, γ) -sequence with $\gamma = (X, Y)$. Let us extend this sequence to a maximal (A, R, γ) -sequence $u^* = (u_0, \dots, u_k)$. Since $u_k \in W(A, R)$, 2.8 implies that $\deg_X u_k > 1$. By well-known properties of coordinate systems (of A), there exists $u_{k+1} \in A$ such that $A = \mathbf{k}[u_k, u_{k+1}]$ and $\deg_X u_{k+1} < \deg_X u_k$. By maximality of u^* , $u_{k+1} \notin W(A, R)$; hence 2.11 implies that u_{k+1} is an R -companion of u_k . Applying 2.13 with $B = R$, we conclude that u_1 is an R -companion of u_0 . $\square$

2.15. Corollary. *Let $A, B \in \Sigma(R)$ be such that $A \subset B$ and such that every element of $W(A, B)$ has a B -companion in A . Then every element of $W(A, R)$ has an R -companion in A ; more precisely, if $u \in W(A, R)$ then every B -companion of u in A is also an R -companion of u .*

Proof. Let $u \in W(A, R)$; since $W(A, R) \subseteq W(A, B)$, u has a B -companion $v \in A$. We show that v is also an R -companion of u .

In view of 2.11, we may assume that $v \in W(A, R)$. Choose $\gamma \in \Gamma(B)$; then (u, v) is an (A, R, γ) -sequence and can be extended to a maximal one, $u^* = (u_0, \dots, u_k)$, with $k \geq 1$, $u_0 = u$, and $u_1 = v$. Since $u_k \in W(A, R) \subseteq W(A, B)$, u_k has a B -companion $u_{k+1} \in A$. Now $u_{k+1} \notin W(A, R)$ by maximality of u^* , so u_{k+1} is also an R -companion of u_k by 2.11. By 2.13, we conclude that $u_1 = v$ is an R -companion of $u_0 = u$. $\square$

We can now finish the proof of the main theorem:

Proof of 2.5. Suppose (2) holds, i.e., $A = \mathbf{k}[X, Y^p]$ for some $X, Y \in R$. Let $B = \mathbf{k}[X, Y] \in \Sigma(R)$. By 2.14, every element of $W(A, B)$ has a B -companion in A ; thus 2.15 implies that every element of $W(A, R)$ has an R -companion in A . $\square$

ACKNOWLEDGMENT

The author thanks R. Ganong for many substantial discussions about weak variables and related questions.

REFERENCES

1. S. Abhyankar, *Lectures on expansion techniques in algebraic geometry* (Notes by B. Singh), Tata Inst. Fund. Res. Lectures on Math. and Phys., vol. 57, Tata Inst. Fund. Res., Bombay, 1977.
2. S. Abhyankar and B. Singh, *Embeddings of certain curves in the affine plane*, Amer. J. Math. **100** (1978), 99–175.
3. D. Daigle, *Plane Frobenius sandwiches of degree p* , Proc. Amer. Math. Soc. **117** (1993), 885–889.
4. R. Ganong, *On plane curves with one place at infinity*, J. Reine Angew. Math. **307/308** (1979), 173–193.
5. —, *Plane Frobenius sandwiches*, Proc. Amer. Math. Soc. **84** (1982), 474–478.
6. N. Jacobson, *Lectures in abstract algebra*, Vol. 3, Graduate Texts in Math., vol. 32, Springer-Verlag, New York, 1964.

7. T. T. Moh, *On the classification problem of embedded lines in characteristic p* , Algebraic Geometry and Commutative Algebra in Honor of Masayoshi NAGATA, Kinokuniya, 1987, pp. 267–279.
8. K. P. Russell, *Hamburger-Noether expansions and approximate roots of polynomials*, Manuscripta Math. **31** (1980), 25–95.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OTTAWA, OTTAWA, CANADA K1N 6N5
E-mail address: `daniel@mathstat.uottawa.ca`