GENERALIZATIONS OF DEODHAR'S α -LOCALIZATION FUNCTOR

BEN COX

(Communicated by Roe Goodman)

ABSTRACT. In this paper we generalize the result of Deodhar (see *Invent. Math.* **57** (1980), 101–118) on α -localization functors. Namely, we show that localization with respect to a larger family of left denominator sets “intertwines” with tensoring by finite-dimensional representations. In the language of the author’s previous work, localization with respect to such a left denominator set produces a new example of an $\mathfrak{F}$ -functor and an $\mathfrak{F}$ -category.

1. INTRODUCTION

Let $\mathfrak{g}$ be a semisimple finite-dimensional Lie algebra with triangular decomposition $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$. In [D], Deodhar discovered that Enright’s completion functor (see [E1, E2]) is actually a subfunctor of an α -localization functor. This latter functor is obtained by noncommutative localization with respect to the left denominator set $S_\alpha = \{y_\alpha^n | n \in \mathbb{N}\}$ where y_α is a nilpotent element in $\mathfrak{n}_-$. Localization with respect to this set “intertwines” with tensoring by finite-dimensional $\mathfrak{g}$ -modules. Below we find new examples of noncommutative localization which also intertwine with tensoring by finite-dimensional representations.

2. PRELIMINARIES AND NOTATION

2.1. Recall that an *additive category* $\mathfrak{A}$ is a category satisfying the following three axioms:

- (i) $\mathfrak{A}$ has a zero object;
- (ii) any two objects in $\mathfrak{A}$ have a product; and
- (iii) for all objects $A, B \in \text{Ob } \mathfrak{A}$ the set of morphisms $\text{Hom}_{\mathfrak{A}}(A, B)$ forms an abelian group such that the composition

$$\text{Hom}_{\mathfrak{A}}(A, B) \times \text{Hom}_{\mathfrak{A}}(B, C) \rightarrow \text{Hom}_{\mathfrak{A}}(A, C)$$

is bilinear. One also has the following proposition.

Proposition [HS, Chapter 2]. *Let $\mathfrak{A}$ and $\mathfrak{B}$ be two additive categories and $F: \mathfrak{A} \rightarrow \mathfrak{B}$ a functor. Then the following are equivalent:*

- (i) *F preserves sums (of two objects).*
- (ii) *F preserves products (of two objects).*

Received by the editors April 30, 1992 and, in revised form, October 28, 1992.
 1991 *Mathematics Subject Classification*. Primary 17B35; Secondary 16P50.

(iii) For each $A, A' \in \text{Ob } \mathfrak{A}$ one has that

$$F: \text{Hom}_{\mathfrak{A}}(A, A') \rightarrow \text{Hom}_{\mathfrak{B}}(FA, FA')$$

is a group homomorphism.

A functor satisfying the above equivalent conditions is called an *additive functor*.

For a vector space V over a field k , let $T^n(V)$ denote the n -fold tensor product of V with itself, and let $T^0 = k$. Then $T(V) := \bigoplus_{n=0}^{\infty} T^n(V)$ is the *tensor algebra* of V . Elements in $T^n(V)$ are called *homogeneous of degree n* . If $\mathfrak{g}$ is a Lie algebra, let $U(\mathfrak{g})$ denote the universal enveloping algebra of $\mathfrak{g}$.

3. $\mathfrak{F}$ -CATEGORIES AND $\mathfrak{F}$ -FUNCTORS

3.1. For any Lie algebra $\mathfrak{g}$ (possibly infinite dimensional) defined over a field k of characteristic zero, let $M_{\mathfrak{g}}$ denote the category of all $\mathfrak{g}$ -modules. Throughout we will assume that $\mathfrak{F}$ denotes an additive subcategory of $M_{\mathfrak{g}}$ satisfying the following two conditions:

- (1) $\mathfrak{F}$ is closed under tensoring; i.e., if $E_j, F_j \in \mathfrak{F}$ and $f_j \in \text{Hom}_{\mathfrak{F}}(E_j, F_j)$ for $j = 1, 2$, then $E_1 \otimes E_2, F_1 \otimes F_2 \in \text{Ob } \mathfrak{F}$ and $f_1 \otimes f_2 \in \text{Hom}_{\mathfrak{F}}(E_1 \otimes E_2, F_1 \otimes F_2)$.
- (2) $\mathfrak{g} \in \text{Ob } \mathfrak{F}$ as a $\mathfrak{g}$ -module under the adjoint action.

For $F \in \text{Ob } \mathfrak{F}$, $\mathfrak{g}$ -modules A and B , and $h \in \text{Hom}_{\mathfrak{g}}(A, B)$, let T_F denote the tensor product functor on $M_{\mathfrak{g}}$ given by $A \mapsto F \otimes A$ and $h \mapsto 1_F \otimes h$. If $\mathfrak{n} \subset \mathfrak{g}$ is a Lie subalgebra and $\mathfrak{C}$ is a subcategory of $M_{\mathfrak{n}}$, we shall use the symbol T_F to denote the tensor product functor on $\mathfrak{C}$ when no confusion is likely to arise. We call the category $\mathfrak{C}$ an *$\mathfrak{F}$ -category* if it is additive and T_F carries $\mathfrak{C}$ into itself for all $F \in \text{Ob } \mathfrak{F}$.

Now let $\mathfrak{a}$ and $\mathfrak{b}$ be two Lie subalgebras of $\mathfrak{g}$, and let $\mathfrak{A}$ (resp. $\mathfrak{B}$) be an additive subcategory of $M_{\mathfrak{a}}$ (resp. $M_{\mathfrak{b}}$). Suppose further that both $\mathfrak{A}$ and $\mathfrak{B}$ are $\mathfrak{F}$ -categories and τ is a functor from $\mathfrak{A}$ to $\mathfrak{B}$. We call τ an *intertwining functor* (or *$\mathfrak{F}$ -intertwining functor* when more precision is necessary) if τ is additive and there exists a natural equivalence, for each $F \in \text{Ob } \mathfrak{F}$, $i_F: T_F \circ \tau \rightarrow \tau \circ T_F$.

Suppose τ is an intertwining functor, and let $\mathcal{I} = \{i_F | F \in \text{Ob } \mathfrak{F}\}$ denote the family of natural equivalences above. Recall that a natural equivalence $i_E: T_E \circ \tau \rightarrow \tau \circ T_E$ is a rule that assigns to each object A of $\mathfrak{A}$ an isomorphism $i_E(A): T_E \circ \tau(A) \rightarrow \tau \circ T_E(A)$ such that for every homomorphism $f: A \rightarrow B$ in $\mathfrak{A}$ one has $i_E(B) \circ ((T_E \circ \tau)(f)) = (\tau \circ T_E)(f) \circ i_E(A)$. For convenience we set $i_{E,A} = i_E(A)$ for all $A \in \text{Ob } \mathfrak{A}$, $E \in \text{Ob } \mathfrak{F}$.

Suppose now that for every $E, F \in \text{Ob } \mathfrak{F}$ and $h \in \text{Hom}_{\mathfrak{F}}(E, F)$ one has $h \otimes 1_A \in \text{Hom}_{\mathfrak{A}}(E \otimes A, F \otimes A)$ for $A \in \text{Ob } \mathfrak{A}$. Assume the category $\mathfrak{B}$ has this same property. Then we say that $\mathcal{I}$ is *natural in the $\mathfrak{F}$ -variable* (or *natural in $\mathfrak{F}$*) if the following diagram is commutative for all $E, F \in \text{Ob } \mathfrak{F}$, $A \in \text{Ob } \mathfrak{A}$, and $f \in \text{Hom}_{\mathfrak{F}}(E, F)$:

$$(1) \quad \begin{array}{ccc} E \otimes \tau A & \xrightarrow{i_{E,A}} & \tau(E \otimes A) \\ f \otimes 1_{\tau A} \downarrow & & \downarrow \tau(f \otimes 1_A) \\ F \otimes \tau A & \xrightarrow{i_{F,A}} & \tau(F \otimes A) \end{array}$$

We call $\mathcal{S}$ *distributive* if the following diagram is commutative for $E, F \in \text{Ob } \mathfrak{F}$ and $A \in \text{Ob } \mathfrak{A}$:

$$(2) \quad \begin{array}{ccc} (E \oplus F) \otimes \tau A & \xrightarrow{i_{E \oplus F, A}} & \tau((E \oplus F) \otimes A) \\ \downarrow & & \downarrow \\ (E \otimes \tau A) \oplus (F \otimes \tau A) & \xrightarrow{i_{E, A} \oplus i_{F, A}} & \tau(E \otimes A) \oplus \tau(F \otimes A) \end{array}$$

The left map (2) expresses the bilinearity of $\otimes$, and the right map expresses this bilinearity combined with additivity of τ .

We say that $\mathcal{S}$ is *associative* if the following diagram is commutative for all $E, F \in \text{Ob } \mathfrak{F}$ and $A \in \text{Ob } \mathfrak{A}$:

$$(3) \quad \begin{array}{ccc} E \otimes F \otimes \tau A & \xrightarrow{1 \otimes i_{F, A}} & E \otimes \tau(F \otimes A) \\ & \searrow i_{E \otimes F, A} & \downarrow i_{E, F \otimes A} \\ & & \tau(E \otimes F \otimes A) \end{array}$$

3.2 Lemma [C]. Suppose $\mathfrak{A}$ and $\mathfrak{B}$ are $\mathfrak{F}$ -categories and $\tau: \mathfrak{A} \rightarrow \mathfrak{B}$ is an intertwining functor. If the family $\mathcal{S} = \{i_{F, A} | F \in \text{Ob } \mathfrak{F}, A \in \text{Ob } \mathfrak{A}\}$ is natural in the $\mathfrak{F}$ -variable then $\mathcal{S}$ is distributive.

Suppose τ is an intertwining functor with the family of natural equivalences $\mathcal{S} = \{i_F | F \in \text{Ob } \mathfrak{F}\}$. We call the pair $(\tau, \mathcal{S})$ an $\mathfrak{F}$ -functor whenever $\mathcal{S}$ is both distributive and associative. When $\mathcal{S}$ is understood to be fixed we call τ an $\mathfrak{F}$ -functor. (See [C] and the references listed there for several examples of $\mathfrak{F}$ -functors in the representation theory of Lie algebras.)

4. DEODHAR'S α -LOCALIZATION FUNCTOR AND NEW EXAMPLES OF $\mathfrak{F}$ -FUNCTORS

4.1. Let $\mathfrak{g}$ be a Kac-Moody algebra, and let $\mathfrak{F}$ be the category of integrable $\mathfrak{g}$ -modules. In this section we review Deodhar's α -localization functor D_α for α a real root of $\mathfrak{g}$. Our main result, Theorem 4.18, is a substantial generalization of Deodhar's results on localization and will provide us with new examples of $\mathfrak{F}$ -functors. Let us begin by reviewing some noncommutative ring theory (our general reference will be by [GW]).

4.2. All rings in this article are assumed to have an identity. If X is a multiplicative subset then a *left ring of fractions for R with respect to X* is a ring homomorphism $\phi: R \rightarrow S$ such that

- (a) $\phi(x)$ is a unit in S for all $x \in X$,
- (b) each element of S can be written in the form $\phi(x)^{-1}\phi(r)$ for some $x \in X$ and $r \in R$, and
- (c) $\ker(\phi) = \{r \in R | xr = 0 \text{ for some } x \in X\}$.

A multiplicative set X in a ring R that satisfies the following two conditions is called a *left denominator set*:

- (Da) $Xr \cap Rx \neq \emptyset$ for all $r \in R$ and $x \in X$;
- (Db) if $r \in R$ and $x \in X$ are such that $xr = 0$ then there exists $x' \in X$ such that $rx' = 0$.

A well-known result due to Goldie is

4.3. Theorem [GW, Proposition 9.7]. *Let X be a multiplicative set in a ring R . Then there exists a left ring of fractions for R with respect to X if and only if X is a left denominator set.*

Moreover, if a left ring of fractions exists then it is unique up to isomorphism (see [GW, Corollary 9.5]). If X is a left denominator set then we let $X^{-1}R$ denote its unique ring of fractions. This (as is well known) can be constructed as follows: Define on $X \times R$ an equivalence relation $\sim$ where $(x, r) \sim (x', r')$ if there exists $s \in R$ and $y \in X$ such that $yr = sr'$ and $yx = sx'$. The set of equivalence classes of $X \times R$ has an obvious ring structure which gives us $X^{-1}R$. We let $x \setminus r$ denote the equivalence class of (x, r) .

4.4. Let X be a left denominator set, and let A be a left R -module. A *module of fractions for A with respect to X* is an R -module homomorphism $f: A \rightarrow B$ where B is a left $X^{-1}R$ -module such that

- (a) every element of B can be written in the form $x^{-1}f(a)$ for some $x \in X$ and $a \in A$, and
- (b) $\ker f = \{a \in A \mid ax = 0 \text{ for some } x \in X\}$.

Another basic fact is

Theorem [GW, Corollary 9.11 and Theorem 9.13]. *If X is a left denominator set in a ring R then there exists a unique (up to isomorphism) module of fractions for any left R -module A with respect to X .*

4.5. When X is a left denominator set and A is a left R -module then we let $X^{-1}A$ denote its unique module of fractions with respect to X . In addition there is the following useful construction of $X^{-1}A$.

4.6. Theorem [GW, Proposition 9.14]. *For X a left denominator set in R and A a left R -module we have*

$$X^{-1}R \otimes_R A \cong X^{-1}A$$

where the map is given by $s \otimes a \mapsto sa$ for $a \in A$ and $s \in X^{-1}R$.

One has the following universal mapping property of localizations.

Proposition [GW, Proposition 9.10]. *Let X be a left denominator set in a ring R , let A be a left R -module, and suppose C is a left $X^{-1}R$ -module with $g: A \rightarrow C$ an R -module homomorphism. If $f: A \rightarrow X^{-1}A$ is the module of fractions for A with respect to X , then there exists a unique $X^{-1}R$ -module homomorphism $h: X^{-1}A \rightarrow C$ such that $g = h \circ f$.*

4.7. Next we define Deodhar's α -localization functor and record the well-known fact that it is just "localization" with respect to the multiplicative set $S_\alpha = \{y_\alpha^n \mid n \in \mathbb{N}\}$ where $y_\alpha \in \mathfrak{g}_{-\alpha}$ is nonzero and α is a positive root.

Let $\mathcal{A}$ denote the category of $\mathfrak{h}$ -semisimple $U(\mathfrak{n}_-)$ -torsionfree $\mathfrak{g}$ -modules. If $A \in \text{Ob } \mathcal{A}$, let $A' = \{y_\alpha^{-n} \mid n \in \mathbb{N}\} \times A$, and define an equivalence relation on A' by $(y_\alpha^{-n}, a) \sim (y_\alpha^{-m}, a')$ if and only if $y_\alpha^n a' = y_\alpha^m a$. Set $D_\alpha(A) = A' / \sim$. $D_\alpha(A)$ is given a $\mathfrak{g}$ -module structure as follows.

4.8. Lemma [D, Lemma 2.1]. *If $z \in \mathfrak{g}$ and $0 \leq r \in \mathbb{Z}$ then there exists $0 \leq s \in \mathbb{Z}$ such that $y_\alpha^s z = u y_\alpha^r$ for some $u \in U(\mathfrak{g})$.*

Remark. Deodhar's proof is based on a straightforward $\mathfrak{sl}(2, \mathbb{C})$ computation, and from the proof it is easy to see that this lemma is also true for $z \in U(\mathfrak{g})$.

so that S_α is a left Ore set, i.e., it satisfies (Da). It then follows from a result of Goldie (see [GW, Proposition 9.9]) that S_α is a left denominator set.

Now if $(y_\alpha^{-r}, a) \in D_\alpha(A)$ and $z \in \mathfrak{g}$ then $z(y_\alpha^{-r}, a)$ is defined to be (y_α^{-s}, ua) where u is in the lemma above.

4.9. Proposition [D, Proposition 2.2]. *Under the action of $\mathfrak{g}$ on D_α given above, $D_\alpha(A)$ becomes a well-defined $\mathfrak{g}$ -module.*

Remark. By definition D_α is Deodhar's α -localization functor.

Next for $A \in \mathcal{A}$, $A \hookrightarrow D_\alpha(A)$ as a $\mathfrak{g}$ -module so that by the universal mapping property of the module of fractions there exists a unique $S_\alpha^{-1}U(\mathfrak{g})$ -module map $S_\alpha^{-1}A \rightarrow D_\alpha A$. In fact this map is given by $y_\alpha^{-n}a \mapsto (y_\alpha^{-n}, a)$. This map is clearly a bijection. Hence $S_\alpha^{-1}|_{\mathcal{A}} \cong D_\alpha$.

4.10. We now turn to our generalization of these results of Deodhar. Let $X_0 \subset \mathfrak{g} \oplus \mathbb{C} = U(\mathfrak{g})_1$, and let $X \subset U(\mathfrak{g})$ be a left denominator set generated by 1 and X_0 with the following property: for $x \in X_0$ and E any finite-dimensional $\mathfrak{g}$ -module

(1) if λ is an eigenvalue of the action of x on E then $x + \lambda \cdot 1 \in X$.

We shall suppose throughout this section that X satisfies (1).

For convenience we set $R = U(\mathfrak{g})$ and $S = X^{-1}R$.

4.11. Lemma. *Let $x \in X$. Then x acts by an isomorphism on $S \otimes E$.*

Proof. It is sufficient to check the lemma for $x \in X_0$. Introduce an x -stable filtration $0 = E_0 \subset \cdots \subset E_d$ on E with x acting by $\lambda_i \cdot 1$ on E_i/E_{i-1} . Then suppose x acts by isomorphism on $S \otimes E_{i-1}$. Thus for $\{e_k\}$ a basis of E with $e_{i_j} \in E_i$

$$\begin{aligned} x(s \otimes e_{i_j}) &\equiv xs \otimes e_{i_j} + s \otimes \lambda_i e_{i_j} \mod S \otimes E_{i-1} \\ &\equiv (x + \lambda_i)s \otimes e_{i_j} \mod S \otimes E_{i-1}. \end{aligned}$$

This proves the lemma.

S is an R -bimodule. We shall need other bimodules and so introduce the notation $E \otimes \mathbb{C}$ to denote the R -bimodule with left action given on E and trivial right action. Similarly, let E^σ be the right R -module defined by $r(x) \cdot e = \sigma(x) \cdot e$, $e \in E$, $x \in R$, and σ is the involutive antiautomorphism of $U(\mathfrak{g})$ equal to -1 on $\mathfrak{g}$. Then $\mathbb{C} \otimes E^\sigma$ is an R -bimodule with trivial left action and right action given on E^σ .

4.12. Lemma. *As S -bimodules we have an isomorphism $S \otimes (E \otimes \mathbb{C}) \cong S \otimes (\mathbb{C} \otimes E^\sigma)$.*

Proof. Consider the map $\underline{\phi}: (R \otimes (E \otimes \mathbb{C})) \rightarrow R \otimes (\mathbb{C} \otimes E^\sigma)$ given by $x \otimes e \otimes 1 \mapsto (1 \otimes 1 \otimes e) \cdot \sigma(x)$. An easy induction argument using the filtration on $U(\mathfrak{g})$ shows that $\underline{\phi}$ is surjective even at the filtered left; i.e., $\underline{\phi}: R_i \otimes (E \otimes \mathbb{C}) \rightarrow R_i \otimes (\mathbb{C} \otimes E^\sigma)$ is surjective.

To prove injectivity suppose $0 = \sum_i (1 \otimes 1 \otimes e_i) \cdot \sigma(x_i)$ where the e_i are as in the proof of Lemma 4.11 and let d be the maximal integer such that some $x_j \in R_d$ but $x_j \notin R_{d-1}$. Then $0 \equiv \sum_i \sigma(x_i) \otimes 1 \otimes e_i \mod R_{d-1} \otimes (\mathbb{C} \otimes E^\sigma)$. However, the e_i are linearly independent, so each $\sigma(x_i) \in R_{d-1}$. We clearly have a contradiction. This proves injectivity; thus $\underline{\phi}$ is an isomorphism.

Now we extend ϕ to a map $\phi: S \otimes (E \otimes \mathbb{C}) \rightarrow S \otimes (\mathbb{C} \otimes E^\sigma)$. We must define $\phi(x \setminus r \otimes e \otimes 1)$. Recall that x acts on the right of $S \otimes (\mathbb{C} \otimes E^\sigma)$ by an isomorphism $i(x)$. Set

$$\phi(x \setminus r \otimes e \otimes 1) = (1 \otimes 1 \otimes e)i(x)^{-1}a(r) = (1 \otimes 1 \otimes e)a(x \setminus r)$$

where $a(-)$ denotes the right action of R on $S \otimes (\mathbb{C} \otimes E^\sigma)$. To see that ϕ is well defined suppose $x \setminus r = x' \setminus r'$, i.e., there exists $y \in X$ and $s \in R$ such that $yr = sr'$ and $yx = sx'$. Then we have $a(yr) = a(sr')$ and $a(yx) = a(sx')$; thus, ϕ is well defined.

Next we check that ϕ is an R -bimodule isomorphism. Suppose $x \in \mathfrak{g}$, and let $l(x)$ ($r(x)$) denote the left (resp. right) action. Then for $y \in X$, $z \in R$, $e \in E$ we have

$$\begin{aligned} l(x)\phi(y \setminus z \otimes e \otimes 1) &= l(x)(1 \otimes 1 \otimes e)r(y)^{-1}r(z) = (x \otimes 1 \otimes e)r(y)^{-1}r(z) \\ &= (x \otimes 1 \otimes e)r(y)^{-1}r(z) + (1 \otimes 1 \otimes \sigma(x) \cdot e)r(y)^{-1}r(z) \\ &\quad + (1 \otimes 1 \otimes x \cdot e)r(y)^{-1}r(z) \\ &= (1 \otimes 1 \otimes e)r(x)r(y)^{-1}r(z) + (1 \otimes 1 \otimes x \cdot e)r(y)^{-1}r(z) \\ &= \phi(x(y \setminus z) \otimes e \otimes 1) + \phi(y \setminus z \otimes x \cdot e \otimes 1). \end{aligned}$$

Now for the right action we have

$$\begin{aligned} r(x)\phi(y \setminus z \otimes e \otimes 1) &= (1 \otimes 1 \otimes e)r(y)^{-1}r(z)r(x) \\ &= \phi(y \setminus zx \otimes e \otimes 1) = \phi(y \setminus z \otimes e \otimes 1)r(x). \end{aligned}$$

Finally we check that ϕ is an S -bimodule map. Suppose $x \in X$; then from above $l(x) \circ \phi = \phi \circ l(x)$. By Lemma 4.11 we can invert these left actions. Multiplying out we have $\phi \circ l(x)^{-1} = l(x)^{-1} \circ \phi$; so ϕ intertwines with the left action of S . Similarly for the right action. This proves the lemma.

4.13. Lemma. *Let F be any left R -module. As left S -modules we have the isomorphism*

$$S \otimes_R (E \otimes F) \cong S \otimes (\mathbb{C} \otimes E^\sigma) \otimes_R F.$$

Proof. The left tensor product above is the quotient of $S \otimes E \otimes F$ by the relations determined by a $s \cdot x \otimes e \otimes f = s \otimes x \cdot (e \otimes f)$ where $s \in S$, $e \in E$, $f \in F$, and $x \in \mathfrak{g}$. However, this is precisely the same set of relations as $(s \otimes e')x \otimes f = s \otimes e' \otimes x \cdot f$ for $e' \in E^\sigma$. This proves the lemma.

In the next lemma we introduce another multiplicative subset of $U(\mathfrak{g})$. We will see later (4.18) that this gives us another example of an $\mathfrak{F}$ -functor. First we define an $\mathfrak{h}$ -graded multiplicative subset of $U(\mathfrak{g})$ to be a multiplicative subset Y of $U(\mathfrak{g})$ such that $Y = p \bigcup_{\eta \in \mathfrak{h}^*} Y_\eta$ and $Y_\eta \cdot Y_\beta \subset Y_{\eta+\beta}$ where $Y_\eta := \{y \in Y \mid [h, y] = \eta(h)y \text{ for all } h \in \mathfrak{h}^*\}$.

4.14. Lemma. *Let Y be an $\mathfrak{h}$ -graded multiplicative subset of $U(\mathfrak{n}_-)$, W an $\mathfrak{h}$ -semisimple Y -divisible $\mathfrak{g}$ -module, and E a finite-dimensional $\mathfrak{g}$ -module. Then $E \otimes W$ is an $\mathfrak{h}$ -semisimple Y -divisible $\mathfrak{g}$ -module.*

Proof. The fact that $E \otimes W$ is $\mathfrak{h}$ -semisimple is obvious. Thus we need only show that $x(E \otimes W) = E \otimes W$ for all $x \in Y$. We may assume x is not an element in $Y_0 = \{y \in Y \mid h.y = 0 \text{ for all } h \in \mathfrak{h}\}$. The result will follow if we can show that, for arbitrary $e \in E_\gamma$ and $w \in W_\lambda$, $e \otimes w$ is in $x(E \otimes W)$. For $\lambda \in \mathfrak{h}^*$

let E_λ denote the λ th weight space of E . Since E is finite dimensional, the set $\Lambda(E) = \{\lambda \in \mathfrak{h}^* | E_\lambda \neq 0\}$ is finite; hence there exists a minimal $\mu \in \Lambda(E)$. If $\gamma \in \Lambda(E)$ is minimal then $xe = 0$ for $e \in E_\gamma$. Now as W is Y -divisible there exists for any $w \in W$ an $w' \in W$ such that $xw' = w$ and $x(e \otimes w') = e \otimes w$. Thus we may assume as an induction hypothesis that γ is not minimal, and the previous statement is true for all $e \in E_\mu$, $w \in W$ with $\mu < \gamma$. Now if $e' \in E_\gamma$ and $w \in W$ then

$$x(e' \otimes w') = e' \otimes w + \sum_i e'_i \otimes w'_i$$

where $e'_i \in E_{\mu_i}$ with $\mu_i < \gamma$ and $xw' = w$. By induction $e'_i \otimes w'_i \in x(E \otimes W)$; thus $e' \otimes w \in x(E \otimes W)$. This completes the proof of the lemma.

4.15. Lemma. *Let Y be an $\mathfrak{h}$ -graded multiplicative subset of $U(\mathfrak{n}_-)$, W an $\mathfrak{h}$ -semisimple Y -torsionfree $\mathfrak{g}$ -module, and E a finite-dimensional $\mathfrak{g}$ -module. Then $E \otimes W$ is an $\mathfrak{h}$ -semisimple Y -torsionfree $\mathfrak{g}$ -module.*

Proof. Let $\sum_i e_{\mu_i} \otimes w_{\lambda_i} \neq 0$ with $e_{\mu_i} \in E_{\mu_i}$, $w_{\lambda_i} \in W_{\lambda_i}$ linearly independent and $x(\sum_i e_{\mu_i} \otimes w_{\lambda_i}) = 0$ for some $x \in Y_\eta$ where $\eta \neq 0$. Lexicographically order the set $\Lambda(E) \times \Lambda(W)$ (notation as in the previous lemma), and choose (μ_j, λ_j) maximal with respect to this ordering and such that $e_{\mu_i} \otimes w_{\lambda_i} \neq 0$. Then $0 = \sum_i e_{\mu_i} \otimes xw_{\lambda_i} + \sum_i e_{\mu'_i} \otimes w_{\lambda'_i}$ where $\mu'_k < \mu_j$ for all k . But then $e_{\mu_j} \otimes xw_{\lambda_j} = 0$; thus $xw_{\lambda_j} = 0$. Since W is Y -torsionfree, $x = 0$. Hence $E \otimes W$ is Y -torsionfree.

4.16. Lemma. *If E and W are $\mathfrak{g}$ -modules with E finite dimensional then*

$$\text{Hom}(E, W) \cong E^* \otimes W.$$

4.17. Theorem (Mackey Isomorphism Theorem). *Suppose V and E are $\mathfrak{g}$ -modules with E finite dimensional and V $\mathfrak{h}$ -semisimple. Let $Y \subseteq U(\mathfrak{n}_-)$ be an $\mathfrak{h}$ -graded left denominator set. Then we have a natural isomorphism*

$$Y^{-1}(V \otimes E) \cong (Y^{-1}V) \otimes E$$

of $\mathfrak{g}$ -modules.

Proof. For the proof of this theorem set $R = U(\mathfrak{n}_-)$, and let W be an $\mathfrak{h}$ -semisimple $Y^{-1}R$ -module. Now consider the following sequence of isomorphisms:

$$\begin{aligned} & \text{Hom}_{Y^{-1}R}(Y^{-1}(E \otimes V), W) \\ & \cong \text{Hom}_R(E \otimes V, W) \cong \text{Hom}_R(V, \text{Hom}(E, W)) \\ & \cong \text{Hom}_{Y^{-1}R}(Y^{-1}V, \text{Hom}(E, W)) \cong \text{Hom}_{Y^{-1}R}(E \otimes Y^{-1}V, W). \end{aligned}$$

The first isomorphism is just given by Proposition 4.6. The second and fourth isomorphisms are derived from the adjoint associativity property of Hom . Now an R -module is a $Y^{-1}R$ -module if and only if it is Y -divisible and Y -torsionfree (see [GW, Proposition 9.12]). Thus Lemmas 4.14 through 4.16 imply that $\text{Hom}(E, W)$ is a Y^{-1} -module so that we can apply Proposition 4.6 to obtain the third isomorphism. We now proceed as in [Kn, Proposition 5.14] and let $W = Y^{-1}(E \otimes V)$. The identity in $\text{Hom}_{Y^{-1}R}(Y^{-1}(E \otimes V), Y^{-1}(E \otimes V))$ induces a map in $\text{Hom}_{Y^{-1}R}(E \otimes Y^{-1}V, Y^{-1}(E \otimes Y^{-1}V))$ which we will denote by φ . Similarly the identity morphism in $\text{Hom}_{Y^{-1}R}(E \otimes Y^{-1}V, E \otimes Y^{-1}V)$ induces a map $\psi \in \text{Hom}_{Y^{-1}R}(Y^{-1}(E \otimes V), E \otimes Y^{-1}V)$. One can check as in

[Kn, Proposition 5.14] that $\psi \circ \varphi = 1$ and $\varphi \circ \psi = 1$. This completes the proof of the theorem.

We now consider localization as a functor on the category $M_{\mathfrak{g}}$ of all $\mathfrak{g}$ -modules. Let $\mathfrak{F}$ denote the subcategory of all finite-dimensional $\mathfrak{g}$ -modules. Set $\tau_X M = X^{-1}M$ and $\tau_Y M = Y^{-1}M$ where X satisfies 4.10.1, and $Y \subseteq U(\mathfrak{n}_-)$ is an $\mathfrak{h}$ -graded left denominator set. The main result of the article is

4.18. Theorem. τ_X and τ_Y are intertwining $\mathfrak{F}$ -functors that are natural in $\mathfrak{F}$.

Proof. Theorem 4.17 proves the theorem for Y . For X we let $E \in \text{Ob } \mathfrak{F}$, $M \in M_{\mathfrak{g}}$. Then

$$\begin{aligned} \tau_X(E \otimes M) &= S \otimes_R (E \otimes M) \cong S \otimes (\mathbb{C} \otimes E^\sigma) \otimes_R M \quad (\text{by Lemma 4.13}) \\ &\cong S \otimes (E \otimes \mathbb{C}) \otimes_R M \cong E \otimes \tau_X M \quad (\text{by } \phi^{-1} \text{ and Lemma 4.12}). \end{aligned}$$

The first isomorphism is induced by the identity map, so we shall identify these spaces. For the second isomorphism we have ϕ^{-1} where ϕ is given by $\phi = \phi_E$ where $\phi_E(x \setminus r \otimes e \otimes 1) = (1 \otimes 1 \otimes e)a(x)^{-1}a(r)$.

Now suppose $\gamma \in \text{Hom}_{\mathfrak{g}}(E, F)$; $E, F \in \text{Ob } \mathfrak{F}$. Since $1 \otimes \gamma$ intertwines the (right) action of $\mathfrak{g}$ from $S \otimes E^\sigma$ to $S \otimes F^\sigma$, $1 \otimes \gamma$ intertwines that of $a(x)^{-1}$ as well. This implies that the family of equivalences $\{\phi_E | E \in \text{Ob } \mathfrak{F}\}$ is natural in $\mathfrak{F}$. This proves the theorem.

Remark. For the case $X = S_\alpha$ the result above is due to Deodhar (see [D, Theorem 3.1]) where again the proof is based on an $\text{sl}(2, \mathbb{C})$ calculation.

4.19. Corollary. τ_X and τ_Y are $\mathfrak{F}$ -functors.

Proof. By Theorem 4.18 and Lemma 3.2 we need only show that $\tau = \tau_X$ and $\tau = \tau_Y$ are associative. First consider $\tau = \tau_X$. Suppose $E, F \in \text{Ob } \mathfrak{F}$, $M \in \text{Ob } M_{\mathfrak{g}}$. Since the maps involved are all left S -module maps, we need only verify the identity on a set of S -generators for the space. Clearly $\mathbb{C} \otimes \mathbb{C} \otimes E^\sigma \otimes F^\sigma \otimes_R M$ is a set of generators; so $\mathbb{C} \otimes E \otimes F \otimes \mathbb{C} \otimes_R M$ is a set of S -generators for $S \otimes E \otimes F \otimes \mathbb{C} \otimes_R M$. But $\phi_{E \otimes F}$ and also $\phi_E \circ (1 \otimes \phi_F)$ essentially equal the identity map on these generators;

$$\begin{aligned} \phi_{E \otimes F}(1 \otimes e \otimes f \otimes 1 \otimes m) &= 1 \otimes 1 \otimes e \otimes f \otimes m \\ &= \phi_E \circ (1 \otimes \phi_F)(1 \otimes e \otimes f \otimes 1 \otimes m). \end{aligned}$$

If $\tau = \tau_Y$, one uses the identification $Y^{-1}R \otimes_R A \cong Y^{-1}A$ of Theorem 4.6 and observes that the isomorphism of 4.17 is given by $1 \otimes v \otimes e \mapsto 1 \otimes v \otimes e$ on a set of S generators of $Y^{-1}(V \otimes E)$. Now one checks as before that the appropriate identity is satisfied on the set $\mathbb{C} \otimes E \otimes F \otimes A$ of S -generators of $\tau_Y(E \otimes F \otimes A)$.

4.20. Examples. Here we describe some multiplicative sets X other than Deodhar's S_α , for which the hypothesis of 4.18 is satisfied. Set $\mathbb{Z}_{\geq 0} = \{k \in \mathbb{Z} | k \geq 0\}$.

(a) If A is a $\mathbb{C}$ -algebra and $x \in A$ then let $d_x: A \rightarrow A$ denote the $\mathbb{C}$ -algebra map given by $d_x(y) = xy - yx$ for all $y \in A$. For this first example we need

Theorem [BR, Satz 2.2]. *Let A be a prime Noetherian $\mathbb{C}$ -algebra. Suppose $x \in A$ is an element such that d_x is locally nilpotent on A . If x is not*

nilpotent, then x is a nonzero divisor in A and the localization $S^{-1}A$ exists where $S = \{x^n | n \in \mathbb{Z}_{\geq 0}\}$.

First let $\mathfrak{g} = \mathfrak{sl}(3, \mathbb{C})$, $\mathfrak{h}$ a Cartan subalgebra of $\mathfrak{g}$, $R \subset \mathfrak{h}^*$ the set of nonzero roots, $R^+ \subset R$ a set of positive roots, and $B = \{\alpha, \beta\} \subset R^+$ a basis for R . Fix a Chevalley basis $\{x_\gamma, h_\alpha, h_\beta | \gamma \in R\}$ of $\mathfrak{g}$ where x_γ denotes the element in this basis with weight $\gamma \in R$. Let $n_\pm = \bigoplus_{\alpha \in R_\pm} \mathbb{C}x_\alpha$. For $\gamma \in R_+$ set $y_\gamma = x_{-\gamma}$.

Fix $\varepsilon \in \mathbb{C} \setminus \{0, 1\}$, and define $S_\varepsilon = \{(y_\alpha y_\beta - \varepsilon[y_\alpha, y_\beta])^n | n \in \mathbb{Z}_{\geq 0}\}$. By the Theorem above S_ε is a multiplicative left denominator subset of $U(\mathfrak{g})$. It is also clear that S_ε is $\mathfrak{h}$ -graded. Consequently S_ε satisfies the hypotheses of 4.18. Our goal is to show that localization with respect to S_ε is not the same as localization with respect to the denominator set $S_\gamma^\mu := \{(\mu y_\gamma)^n | n \in \mathbb{Z}_{\geq 0}\}$ for any $\gamma \in R$ and $\mu \in \mathbb{C} \setminus \{0\}$.

We first consider the case that $\gamma = \alpha$. Suppose now that localization with respect to S_α^μ is the same as localization with respect to S_ε . Then y_α is invertible in $S_\varepsilon^{-1}U(\mathfrak{g})$; thus, $(s \setminus m)y_\alpha = 1$ for some $s \in S_\varepsilon$ and $m \in U(\mathfrak{g})$. Consequently there exists $a \in U(\mathfrak{g})$ and $b \in S_\varepsilon$ with $as = b \in S_\varepsilon$ and $amy_\alpha = b$. This implies that $my_\alpha = s = (y_\alpha y_\beta - \varepsilon[y_\alpha, y_\beta])^n$ for some $n \geq 0$ ($a \neq 0$ since otherwise $0 = as = b \in S_\varepsilon$).

Define an ordering of the Chevalley basis by

$$x_\alpha < x_\beta < x_{\alpha+\beta} < h_\alpha < h_\beta < y_{\alpha+\beta} < y_\beta < y_\alpha.$$

Then the Poincaré-Birkhoff-Witt Theorem tells us that $U(\mathfrak{g})$ has a basis of monomials $\bar{x}\bar{h}\bar{y}$ where

$$\bar{x} = x_\alpha^{n_1} x_\beta^{n_2} x_{\alpha+\beta}^{n_3}, \quad \bar{h} = h_\alpha^{n_4} h_\beta^{n_5}, \quad \bar{y} = y_{\alpha+\beta}^{n_6} y_\beta^{n_7} y_\alpha^{n_8},$$

and $n_i \in \mathbb{Z}_{\geq 0}$. Consequently $m = \sum u_{\bar{n}} \bar{x}\bar{h}\bar{y}$ where $u_{\bar{n}} \in \mathbb{C}$ and $\bar{n} = (n_1, \dots, n_8) \in \mathbb{Z}_{\geq 0}^8$.

Thus

$$my_\alpha = \sum u_{\bar{n}} \bar{x}\bar{h}\bar{y}y_\alpha = (y_\alpha y_\beta - \varepsilon[y_\alpha, y_\beta])^n \in U(n_-)n_-.$$

By the Poincaré-Birkhoff-Witt Theorem we have that $m \in U(n_-)n_-$. A straightforward calculation shows that

$$my_\alpha = \sum u_{\bar{n}} \bar{x}\bar{h}\bar{y}y_\alpha = (1 - \varepsilon)^n y_{\alpha+\beta}^n + p(y_\alpha, y_\beta, y_{\alpha+\beta})y_\alpha$$

for some polynomial p . Thus

$$(m - p(y_\alpha, y_\beta, y_{\alpha+\beta}))y_\alpha = (1 - \varepsilon)^n y_{\alpha+\beta}^n.$$

If $\varepsilon \neq 1$, this is impossible by the Poincaré-Birkhoff-Witt Theorem. This proves that localization with respect to S_α^μ is not the same as localization with respect to S_ε if $\varepsilon \neq 1$. A very similar argument (but with a different ordering on the basis) shows that localization with respect to S_γ^μ where $\gamma \in \{\pm\alpha, \pm\beta, \pm(\alpha+\beta)\}$ is not the same as localization with respect to S_ε if $\varepsilon \neq 0, 1$.

(b) Let $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{C})$ with $\{x, h, y\}$ a Chevalley basis of $\mathfrak{g}$ such that $[h, x] = 2x$, $[x, y] = h$, and $[h, y] = -2y$. Let $X_0 = \{h - n1 | n \in \mathbb{Z}\}$ and $X = \{(h - n1)^k | n, k \in \mathbb{Z}, k \geq 0\}$ be subsets of $U(\mathfrak{g})$. This is Example 1.8 in [BR] of an Ore subset of $U(\mathfrak{g})$. Now it is straightforward to check that X satisfies condition 4.10(1). Thus we only need to see that localization with respect to X is not the same as localization with respect to $S_\alpha = \{x^n | n \in \mathbb{Z}_{\geq 0}\}$ or

$S_{-\alpha} = \{y^n | n \in \mathbb{Z}_{\geq 0}\}$. We will prove the case $S_{\alpha} = \{x^n | n \in \mathbb{Z}_{\geq 0}\}$ and leave the other case to the reader. As in the previous example we assume the contrapositive so that there exists $m \in U(\mathfrak{sl}(2, \mathbb{C}))$, $n_1, \dots, n_t \in \mathbb{Z}$, and $k_1, \dots, k_t \in \mathbb{Z}_{\geq 0}$ such that

$$my = (h - n_1)^{k_1} \cdots (h - n_t)^{k_t}.$$

Using the Poincaré-Birkhoff-Witt Theorem we see that this is impossible. Thus, localization with respect to X is not the same as localization with x or y .

ACKNOWLEDGMENT

This paper is a portion of the author's thesis, and he would like to thank his advisor Professor Thomas J. Enright for his generous help and many useful suggestions. He would also like to thank the referee for pointing out some errors in an earlier version of this paper.

REFERENCES

- [BR] W. Borho and R. Rentschler, *Oresche Teilmengen in Einhüllenden Algebren*, Math. Ann. **217** (1975), 201–210.
- [C] B. Cox, *$\mathfrak{F}$ -categories and $\mathfrak{F}$ -functors in the representation theory of Lie algebras*, preprint.
- [D] V. Deodhar, *On a construction of representations and a problem of Enright*, Invent. Math. **57** (1980), 101–118.
- [E1] T. J. Enright, *On the fundamental series of a real semisimple Lie algebra and their irreducibility, resolution and multiplicity formulae*, Ann. of Math. (2) **110** (1979), 1–82.
- [E2] ———, *Representations of complex semisimple Lie groups*, Springer-Verlag, New York, 1981.
- [GW] K. Goodearl and R. Warfield, *An introduction to noncommutative Noetherian rings*, Cambridge Univ. Press, Cambridge and New York, 1989.
- [HS] P. Hilton and U. Stambach, *A course in homological algebra*, Springer-Verlag, New York, 1971.
- [Kn] A. Knapp, *Lie groups, Lie algebras, and cohomology*, Princeton Univ. Press, Princeton, NJ, 1988.

DEPARTMENT OF MATHEMATICAL SCIENCES, UNIVERSITY OF MONTANA, MISSOULA, MONTANA 59812-1032

E-mail address: ma-blc@selway.umt.edu