GROWTH PROPERTY FOR THE MINIMAL SURFACE EQUATION IN UNBOUNDED DOMAINS

JENN-FANG HWANG

(Communicated by Peter Li)

ABSTRACT. Here we prove that if u satisfies the minimal surface equation in an unbounded domain Ω which is properly contained in a half plane, then the growth rate of u is of the same order as the shape of Ω and $u|_{\partial\Omega}$.

1. INTRODUCTION

The purpose of this paper is to improve a Phragmén-Lindelöf Theorem for the minimal surface equation in $\mathbb{R}^2$. Hwang has proved that if u satisfies the minimal surface equation in an unbounded domain Ω , which is properly contained in a half plane, the growth property of u depends on Ω and $u|_{\partial\Omega}$ only, without requiring any restriction for u [3]. In this respect, the Phragmén-Lindelöf Theorem for the minimal surface equation is better than that of the Laplace equation. We remark that if u satisfies the Laplace equation in an unbounded domain Ω , the growth property of u cannot be determined completely by the shape of Ω and $u|_{\partial\Omega}$ alone [7].

But the estimate in [3] is not good enough. For example, let $\Omega = \{-\cosh y < x < \cosh y | y > 0\}$ and $\operatorname{div} Tu = 0$ in Ω , $u|_{\partial\Omega} = \sqrt{(\cosh y)^2 - x^2}|_{\partial\Omega}$. Then, by Example 3.4 of [3], we know that $u = O(ye^y)$ as $y \rightarrow \infty$, but the growth rate of the solution $\sqrt{(\cosh y)^2 - x^2}$ (catenoid) is $O(e^y)$.

The purpose of this paper is to improve the estimate of [3]. We will prove that the growth rate of u is of the same order as the shape of Ω and $u|_{\partial\Omega}$ (Theorems 2.12 and 2.13). In fact, let $\Omega = \{-\cosh y < x < \cosh y\}$. We will prove that a catenoid is the maximum solution among the solutions with vanishing boundary value (Corollary 2.3).

2. PHRAGMÉN-LINDELÖF THEOREMS FOR $\mathbb{R}^2$

Throughout the paper, Ω will be a connected domain (bounded or unbounded) in $\mathbb{R}^2$ and, for any function $u \in C^2(\Omega)$, Tu will denote the vector $Du/\sqrt{1 + |Du|^2}$, where Du is the gradient vector of u and the minimal surface

Received by the editors September 22, 1992.

1991 *Mathematics Subject Classification*. Primary 35B05; Secondary 35J10, 49Q05, 53A10.

©1994 American Mathematical Society
0002-9939/94 \$1.00 + \$.25 per page

operator $\mathfrak{M}$ is given by

$$\mathfrak{M}u = (1 + |Du|^2)\Delta u - D_i u D_j u D_{ij} u = (1 + |Du|^2)^{3/2} \operatorname{div} Tu,$$

where Δu is the Laplacian of u .

We will use functions of the form

$$F(x, y) = (G(x, y))^{1/2} g(y) + h(y)$$

as comparison functions, and we compute $\mathfrak{M}F$ in the following lemma.

Lemma 2.1. *Let*

$$F(x, y) = (G(x, y))^{1/2} g(y) + h(y),$$

where $F, G: \mathbb{R}^2 \rightarrow \mathbb{R}^1$, $g, h: \mathbb{R}^1 \rightarrow \mathbb{R}^1$, $F, G, g, h \in C^2$, and $G > 0$. Then $G^{3/2}\mathfrak{M}F = I + II + III$, where

$$\begin{aligned} I &= g^3 \left(\frac{1}{4} G_x^2 \left(\frac{1}{2} G_{yy} + G \frac{g''}{g} - 3G \left(\frac{g'}{g} \right)^2 \right) \right. \\ &\quad \left. + \frac{1}{2} G_{xx} \left(\frac{1}{4} G_y^2 + G_y G \frac{g'}{g} + G^2 \left(\frac{g'}{g} \right)^2 \right) - \frac{1}{2} G_x G_{xy} \left(\frac{1}{2} G_y + G \frac{g'}{g} \right) \right), \\ II &= g \left(-\frac{1}{4} G_y^2 + \frac{1}{2} G G_{yy} + G G_y \frac{g'}{g} + G^2 \frac{g''}{g} - \frac{1}{4} G_x^2 + \frac{1}{2} G_{xx} G \right), \\ III &= G^{1/2} h'' \left(G + \frac{1}{4} G_x^2 g^2 \right) \\ &\quad + G^{1/2} h' \left(-\frac{1}{2} G_x G_{xy} g^2 - G_x^2 g g' + \frac{1}{2} G_y G_{xx} g^2 + G G_{xx} g g' \right) \\ &\quad + h'^2 \left(-\frac{1}{4} G_x^2 g + \frac{1}{2} G G_{xx} g \right). \end{aligned}$$

Proof. By simple computation we have

$$\begin{aligned} (1 + F_x^2) F_{yy} &= \left(1 + \frac{1}{4} G^{-1} G_x^2 g^2 \right) \\ &\quad \times \left(-\frac{1}{4} G^{-3/2} G_y^2 g + \frac{1}{2} G^{-1/2} G_{yy} g + G^{-1/2} G_y g' + G^{1/2} g'' + h'' \right), \\ -2F_x F_y F_{xy} &= -2 \left(\frac{1}{2} G^{-1/2} G_x g \right) \left(\frac{1}{2} G^{-1/2} G_y g + G^{1/2} g' + h' \right) \\ &\quad \times \left(-\frac{1}{4} G^{-3/2} G_x G_y g + \frac{1}{2} G^{-1/2} G_{xy} g + \frac{1}{2} G^{-1/2} G_x g' \right), \\ (1 + F_y^2) F_{xx} &= \left(1 + \left(\frac{1}{2} G^{-1/2} G_y g + G^{1/2} g' + h' \right)^2 \right) \\ &\quad \times \left(-\frac{1}{4} G^{-3/2} G_x^2 g + \frac{1}{2} G^{-1/2} G_{xx} g \right). \end{aligned}$$

Hence

$$\begin{aligned}
G^{5/2}\mathfrak{M}F &= (G + \frac{1}{4}G_x^2g^2)(-\frac{1}{4}G_y^2g + \frac{1}{2}GG_{yy}g + GG_yg' + G^2g'' + G^{3/2}h'') \\
&\quad - 2(\frac{1}{2}G_xg)(\frac{1}{2}G_yg + Gg' + G^{1/2}h')(-\frac{1}{4}G_xG_yg + \frac{1}{2}GG_{xy}g + \frac{1}{2}GG_xg') \\
&\quad + (G + (\frac{1}{2}G_yg + Gg' + h'G^{1/2})^2)(-\frac{1}{4}G_x^2g + \frac{1}{2}GG_{xx}g) \\
&= \frac{1}{4}G_x^2g^2(-\frac{1}{4}G_y^2g + \frac{1}{2}GG_{yy}g + GG_yg' + G^2g'') \\
&\quad - 2(\frac{1}{2}G_xg)(\frac{1}{2}G_yg + Gg')(-\frac{1}{4}G_xG_yg + \frac{1}{2}GG_{xy}g + \frac{1}{2}GG_xg') \\
&\quad + (\frac{1}{2}G_yg + Gg')^2(-\frac{1}{4}G_x^2g + \frac{1}{2}GG_{xx}g) \\
&\quad + G(-\frac{1}{4}G_y^2g + \frac{1}{2}GG_{yy}g + GG_yg' + G^2g'' - \frac{1}{4}G_x^2g + \frac{1}{2}G_{xx}gG) \\
&\quad + G^{3/2}h''(G + \frac{1}{4}G_x^2g^2) \\
&\quad + G^{1/2}h'(-G_xg)(-\frac{1}{4}G_xG_yg + \frac{1}{2}GG_{xy}g + \frac{1}{2}GG_xg') \\
&\quad + 2h'G^{1/2}(\frac{1}{2}G_yg + Gg')(-\frac{1}{4}G_x^2g + \frac{1}{2}GG_{xx}g) \\
&\quad + h'^2G(-\frac{1}{4}G_x^2g + \frac{1}{2}GG_{xx}g) \\
&= g^3(\frac{1}{4}G_x^2(\frac{1}{2}GG_{yy} + G^2\frac{g''}{g} - 3G^2(\frac{g'}{g})^2) \\
&\quad + \frac{1}{2}GG_{xx}(\frac{1}{2}G_y + G\frac{g'}{g})^2 - \frac{1}{2}GG_xG_{xy}(\frac{1}{2}G_y + G\frac{g'}{g})) \\
&\quad + Gg(-\frac{1}{4}G_y^2 + \frac{1}{2}GG_{yy} + GG_y\frac{g'}{g} + G^2\frac{g''}{g} - \frac{1}{4}G_x^2 + \frac{1}{2}G_{xx}G) \\
&\quad + G(G^{1/2}h''(G + \frac{1}{4}G_x^2g^2) \\
&\quad + G^{1/2}h'(-\frac{1}{2}G_xG_{xy}g^2 - G_x^2gg' + \frac{1}{2}G_yG_{xx}g^2 + GG_{xx}gg') \\
&\quad + h'^2(-\frac{1}{4}G_x^2g + \frac{1}{2}GG_{xx}g)) \\
&= G(I + II + III).
\end{aligned}$$

The lemma follows.

Now we treat the Phragmén-Lindelöf Theorem for comparison functions $H(x, y)$ with faster growth. Since, in a half plane, the bound of the solutions with vanishing boundary value does not even exist, the domain must be properly contained in a half plane.

Lemma 2.2. Let $H(x, y) = a\sqrt{f^2(y) - x^2}$ and $\Omega \subset \{-f_1(y) < x < f_1(y), y > 0\}$, where $f, f_1 : [0, \infty) \rightarrow [0, \infty)$, $f \in C^2$, $f_1 \in C^0$, $f \geq f_1 > 0$ for $y > 0$, $f' \equiv \frac{df}{dy} > 0$, and a is a positive constant. Let $u \in C^2(\Omega) \cap C^0(\bar{\Omega})$, and suppose that

$$(i) \quad \operatorname{div} TH - \operatorname{div} Tu \leq 0 \text{ in } \Omega,$$

$$(ii) \quad (u - H)|_{\partial\Omega} \leq 0,$$

$$(iii) \quad \liminf_{y \rightarrow \infty} f_1(y)/f'^2 = 0.$$

Then $u \leq H$ in Ω .

Remark. It is easy to see that if $\lim_{y \rightarrow \infty} f(y)/f'^2 = 0$, then the rate of growth of f must be faster than y^2 as $y \rightarrow \infty$.

Proof of Lemma 2.2. If $\{(x, y) \in \Omega | u(x, y) - H(x, y) > 0\}$ is nonempty, there exists $\varepsilon > 0$ such that $\Omega' = \{(x, y) \in \Omega | u(x, y) - H(x, y) > \varepsilon\}$ is non-empty and $\partial\Omega' \cap \Omega$ is smooth (Sard's Theorem). Since $(u - H)|_{\partial\Omega} \leq 0$, we have $\partial\Omega' \subset \Omega$ and $\partial\Omega' = \{(x, y) \in \Omega | u - H = \varepsilon\}$.

For every $y_0 > 0$, let $\Omega_{y_0} = \Omega' \cap \{y < y_0\}$ and $\Gamma_{y_0} = \partial\Omega_{y_0} \cap \{y = y_0\}$. By the divergence theorem, we have

$$\begin{aligned} & \int_{\partial\Omega_{y_0}} \tan^{-1}(u - H - \varepsilon)(Tu - TH) \cdot \nu d\sigma \\ &= \int \int_{\Omega_{y_0}} \frac{(Du - DH)}{1 + (u - H - \varepsilon)^2} (Tu - TH) dx \\ & \quad + \int \int_{\Omega_{y_0}} (u - H - \varepsilon)(\operatorname{div} Tu - \operatorname{div} TH) dx, \end{aligned}$$

where ν is the unit outer normal of $\partial\Omega_{y_0}$. Noticing that $\partial\Omega_{y_0} \setminus \Gamma_{y_0} \subset \partial\Omega'$, we have $u - H = \varepsilon$ on $\partial\Omega_{y_0} \setminus \Gamma_{y_0}$. Since $\tan^{-1}(u - H - \varepsilon)(\operatorname{div} Tu - \operatorname{div} TH) \geq 0$ in Ω_{y_0} and $Tu \cdot \nu \leq 1$,

$$\begin{aligned} & \int \int_{\Omega_{y_0}} \frac{(Du - DH)}{1 + (u - H - \varepsilon)^2} (Tu - TH) dx \\ & \leq \int_{\Gamma_{y_0}} \tan^{-1}(u - H - \varepsilon)(Tu - TH) \cdot \nu d\sigma \\ & \leq \int_{\Gamma_{y_0}} \tan^{-1}(u - H - \varepsilon)(1 - TH \cdot \nu) d\sigma. \end{aligned}$$

Since $\nu = \langle 0, 1 \rangle$ on Γ_{y_0} ,

$$H = a\sqrt{f^2 - x^2}, \quad H_y = a\frac{ff'}{\sqrt{f^2 - x^2}}, \quad H_x = -\frac{ax}{\sqrt{f^2 - x^2}},$$

$$\begin{aligned} TH \cdot \nu|_{\Gamma_{y_0}} &= \frac{H_y}{\sqrt{1 + H_x^2 + H_y^2}} = \frac{aff' / \sqrt{f^2 - x^2}}{\sqrt{1 + a^2x^2/(f^2 - x^2) + a^2f^2f'^2/(f^2 - x^2)}} \\ &= \frac{aff'}{\sqrt{f^2 - x^2 + a^2x^2 + a^2f^2f'^2}}, \end{aligned}$$

$$\begin{aligned} (1 - TH \cdot \nu)|_{\Gamma_{y_0}} &= \frac{\sqrt{f^2 - x^2 + a^2x^2 + a^2f^2f'^2} - aff'}{\sqrt{f^2 - x^2 + a^2x^2 + a^2f^2f'^2}} \\ &= \frac{\sqrt{f^2 - x^2 + a^2x^2 + a^2f^2f'^2} + aff'}{\sqrt{f^2 - x^2 + a^2x^2 + a^2f^2f'^2} + aff'} \\ &= \frac{f^2 - x^2 + a^2x^2}{\sqrt{f^2 - x^2 + a^2x^2 + a^2f^2f'^2}(\sqrt{f^2 - x^2 + a^2x^2 + a^2f^2f'^2} + aff')} \\ &\leq \frac{(1 + a^2)f^2}{a^2f^2f'^2} = \frac{1 + a^2}{a^2f'^2}. \end{aligned}$$

Now

$$\begin{aligned} & \int \int_{\Omega_{y_0}} \frac{(Du - DH)}{1 + (u - H - \varepsilon)^2} (Tu - TH) dx \\ & \leq \int_{\Gamma_{y_0}} \tan^{-1}(u - H - \varepsilon)(1 - TH \cdot \nu) d\sigma \\ & \leq \frac{\pi}{2} \int_{\Gamma_{y_0}} \frac{1 + a^2}{a^2 f'^2} d\sigma \leq \pi \cdot \frac{(1 + a^2)f_1}{a^2 f'^2}(y_0). \end{aligned}$$

Let $y_0 \rightarrow \infty$. We have

$$0 \leq \int \int_{\Omega'} \frac{(Du - DH)}{1 + (u - H - \varepsilon)^2} (Tu - TH) dx \leq 0.$$

Since $(Du - DH) \cdot (Tu - TH) \geq 0$ and the equality holds when $Du = DH$, we have $Du - DH \equiv 0$ in Ω' . Then $u \equiv H + \varepsilon$ in Ω' , and, by definition, Ω' must be empty. This is impossible, and we conclude that $u(x) \leq H(x)$ for all x in Ω .

Remark. The following well-known fact is used to prove Lemma 2.2: $Du/\sqrt{1 + |Du|^2}$ has norm less than 1. It is a very important idea for the capillary surface equation (cf. [1, Theorem 5.1]).

Now we obtain the result: catenoid is the maximum solution among those surfaces on $\Omega = \{-\cosh y < x < \cosh y\}$ satisfying the minimal surface equation and with vanishing boundary value.

Corollary 2.3. Let $\Omega = \{-\cosh y < x < \cosh y\}$, and let $u \in C^2(\Omega) \cap C^0(\bar{\Omega})$. Suppose that

- (i) $\operatorname{div} Tu \geq 0$ in Ω ,
- (ii) $u|_{\partial\Omega} \leq 0$.

Then $u \leq \sqrt{(\cosh y)^2 - x^2}$.

Proof. The corollary can be proved by the fact that

$$\operatorname{div} Tu - \operatorname{div} T\sqrt{(\cosh y)^2 - x^2} \geq 0, \quad \lim_{y \rightarrow \pm\infty} \frac{\cosh y}{[(\cosh y)']^2} = 0,$$

and a similar argument as in the proof of Lemma 2.2.

By Lemma 2.2, it is sufficient to show that $\operatorname{div} T(a\sqrt{f^2 - x^2}) \leq 0$ holds when $\liminf_{y \rightarrow \infty} (f_1/f'^2) = 0$. Let $f \in C^2$, $f \geq 0$, and $f' > 0$. Define $p(f) = (1/(\log f)')' = (f'^2 - ff'')/f'^2 = 1 - ff''/f'^2$; some properties of $p(f)$ will be discussed later. By setting $F = a\sqrt{f^2 - x^2}$, $G = f^2 - x^2$, and $g = a$ in Lemma 2.1 where a is a positive constant, we get

$$\begin{aligned} G^{3/2}\mathfrak{M}F &= a^3(x^2 f'^2 + x^2 f f'' - f^2 f'^2) + a(ff''(f^2 - x^2) - x^2 f'^2 - f^2) \\ &= f'^2 a^3(-f^2 + (2 - p(f))x^2) + f'^2 a(f^2(1 - p(f)) - (2 - p(f))x^2) - af^2 \\ &= -(a - a^3)(2 - p(f))x^2 f'^2 + f^2 f'^2 a((1 - p(f)) - a^2) - af^2. \end{aligned}$$

Lemma 2.4. Let f , $p(f)$, F , G , and a be defined as above, and let $\Omega \subset \{(x, y) | -f_1(y) < x < f_1(y), y > 0\}$, where $f_1 > 0$ is given. Suppose $p(f) \geq p_0$, where p_0 is a constant. Then

- (i) if $1 \geq p_0 \geq 0$, $a = \sqrt{1 - p_0}$, and $f \geq f_1$, or

(ii) if $p_0 < 0$, $a > \sqrt{1-p_0}$, a is a positive constant, and $f^2 \geq (a^2 - 1)(2 - p_0)f_1^2/(a^2 - (1 - p_0))$,

we have $G^{3/2}\mathfrak{M}F \leq 0$ in Ω .

Proof. (i) If $2 \geq p(f) \geq p_0$, it is easy to see that

$$G^{3/2}\mathfrak{M}F \leq 0.$$

If $p(f) \geq 2$, we have

$$G^{3/2}\mathfrak{M}F \leq (p(f) - 2)\sqrt{1-p_0}p_0x^2f'^2 + f^2f'^2\sqrt{1-p_0}(-p(f) + p_0) \leq 0.$$

(ii) By assumption, it is easy to have the following:

$$\begin{aligned} G^{3/2}\mathfrak{M}F &= f'^2a^3(-f^2 + (2 - p(f))x^2) \\ &\quad + f'^2a(f^2(1 - p(f)) - (2 - p(f))x^2) - af^2 \\ &= af'^2((a^2 - 1)(2 - p(f))x^2 - f^2(a^2 - (1 - p(f)))) - af^2 \leq 0. \end{aligned}$$

Hence it is easy to derive the following theorem.

Theorem 2.5. Let $f, f_1, p(f)$, and Ω be defined as in Lemma 2.4. Let $\liminf_{y \rightarrow \infty} (f_1/f'^2) = 0$, $u \in C^2(\Omega) \cap C^0(\Omega)$, and $\operatorname{div} Tu \geq 0$ in Ω . Then

- (i) if $p(f) \geq p_0 \geq 0$ where p_0 is a constant, $f \geq f_1$, $1 \geq p_0 \geq 0$, and $u|_{\partial\Omega} \leq \sqrt{1-p_0}\sqrt{f^2-x^2}|_{\partial\Omega}$, then we have $u \leq \sqrt{1-p_0}\sqrt{f^2-x^2}$ in Ω ,
(ii) if $p(f) \geq p_0$, where p_0 is a negative constant, and

$$f^2 \geq \frac{(a^2 - 1)(2 - p_0)}{a^2 - (1 - p_0)} f_1^2, \quad u|_{\partial\Omega} \leq a\sqrt{f^2 - x^2}|_{\partial\Omega},$$

where a is a positive constant satisfying $a^2 - 1 + p_0 > 0$, then we have $u \leq a\sqrt{f^2 - x^2}$ in Ω .

We will show later that the above theorem still remains valid without the condition $\liminf_{y \rightarrow \infty} (f_1/f'^2) = 0$. We first investigate the properties of $p(f)$. Let $f \in C^2$, $f \geq 0$, $f' > 0$, a be a positive constant, and $\alpha \neq 0$ be a constant. Since $p(af^\alpha) = (1/(\log(af^\alpha)))' = \frac{1}{\alpha}(1/(\log f))'$, we have $p(af^\alpha) = \frac{1}{\alpha}p(f)$. Then we prove the following lemma.

Lemma 2.6. Let $f, f_1 \in C^2$, and $f, f', f_1, f'_1 > 0$. Then

- (i) if $p(f), p(f_1) \geq p_0 > 0$, where $0 < p_0 \leq 1$, is a constant, then $p(ff_1) \geq p_0/2 > 0$;
(ii) if $p(f) \geq 0$, then $p(e^{ay}f) \geq 0$ for any positive constant a ;
(iii) if $p(f) \geq p_0$, where p_0 is a negative constant, then $p(e^{ay}f) \geq p_0$ for any positive constant a .

Proof. (i) Let $h = (\log f)' = f'/f > 0$ and let $h_1 = (\log f_1)' = f'_1/f_1 > 0$. Since $p(f) = (1/h)' = -h'/h^2 \geq p_0 > 0$ and $p(f_1) = (1/h_1)' = -h'_1/h_1^2 \geq p_0 > 0$, we have $p(ff_1) = (1/(h+h_1))' = -(h'+h'_1)/(h+h_1)^2 \geq -(h'+h'_1)/2(h^2+h_1^2) \geq p_0/2$.

(ii) Since $p(f) = -h'/h^2 \geq 0$, we have $p(e^{ay}f) = (1/((\log e^{ay})' + (\log f)'))' = (1/(a+h))' = -h'/(a+h)^2 \geq 0$.

(iii) Since $p(f) = -h'/h^2$ and $p(e^{ay}f) = -h'/(a+h)^2$ have the same sign and $|p(e^{ay}f)| \leq |p(f)|$, the result follows immediately.

Remark 2.7. (i) If $f = y^m$, where m is a positive constant, then $p(f) = \frac{1}{m}$.

(ii) $p(e^y) = 0$.

(iii) If we set $f = e^{y^\alpha}$, where $\alpha > 1$ is a constant, then $p(f) = (1/(y^\alpha)')' = (y^{1-\alpha}/\alpha)' = (1-\alpha)y^{-\alpha}/\alpha$. This implies that $p(f) \rightarrow 0^-$ as $y \rightarrow \infty$, so, for sufficiently large y , we have $p(f) > -\varepsilon$ for some small positive number ε .

Similarly, if f increases faster than the exponential function, then we can assume $p(f) \geq -\varepsilon$ for some small positive constant ε essentially.

Lemma 2.8. Let $f \in C^1$, $f, f' > 0$, and $\limsup_{y \rightarrow \infty} (f(y)/y^2) = +\infty$. Then $\liminf_{y \rightarrow \infty} (f/f'^2) = 0$.

Proof. Suppose not; then there exist positive constants y_0 and C such that, for every $y > y_0$, we have $f(y)/f'^2(y) \geq C$, so $f = O(y^2)$. Contradiction arises and the lemma follows.

We are now ready to remove the condition $\liminf_{y \rightarrow \infty} (f/f'^2) = 0$ in Theorem 2.5.

We will start with a theorem.

Theorem 2.9. Let $f \in C^0[0, \infty) \cap C^2(0, \infty)$, $f \geq 0$, $f' > 0$, $\Omega \subset \{(x, y) | -f(y) < x < f(y), y > 0\}$, $u \in C^2(\Omega) \cap C^0(\bar{\Omega})$, $\operatorname{div} Tu \geq 0$, and $p(f) \geq p_0 \geq 0$, where p_0 is a constant. Then

(i) if $p_0 = 0$ and $(u - \sqrt{f^2 - x^2})|_{\partial\Omega} \leq 0$, we have $u \leq \sqrt{f^2 - x^2}$ in Ω ,

(ii) if $1 \geq p_0 > 0$ and $(u - \sqrt{1 - q_0} \sqrt{f^2 - x^2})|_{\partial\Omega} \leq 0$, where $q_0 = \min(1/8, p_0/2)$, we have $u \leq \sqrt{1 - q_0} \sqrt{f^2 - x^2}$ in Ω .

Proof. (i) Let $h = e^{ay} f$, where a is a positive constant. Then by Lemma 2.6 $p(h) \geq 0$. So $\mathfrak{M}(\sqrt{h^2 - x^2}) \leq 0$ in Ω . Since $f' > 0$, $\limsup_{y \rightarrow \infty} (h/y^2) = +\infty$. By Lemma 2.8, $\liminf_{y \rightarrow \infty} (h/h'^2) = 0$. Since $u|_{\partial\Omega} \leq \sqrt{f^2 - x^2}|_{\partial\Omega} \leq \sqrt{h^2 - x^2}|_{\partial\Omega}$, by Theorem 2.5, $u \leq \sqrt{h^2 - x^2} = \sqrt{(e^{ay} f)^2 - x^2}$ in Ω . Setting $a \rightarrow 0$, the result follows.

(ii) Let $h = (1 + by)^4 f$, where b is a positive constant. Since $p((1 + by)^4) = \frac{1}{4}$, by Lemma 2.6, $p(h) \geq \min(1/4, p_0)/2 = q_0 > 0$. By Lemma 2.4, $\mathfrak{M}(\sqrt{1 - q_0} \sqrt{h^2 - x^2}) \leq 0$ in Ω . Since $f' > 0$, $\limsup_{y \rightarrow \infty} (h/y^2) = +\infty$. So $\liminf_{y \rightarrow \infty} (h/h'^2) = 0$. Since

$$u|_{\partial\Omega} \leq \sqrt{1 - q_0} \sqrt{f^2 - x^2}|_{\partial\Omega} \leq \sqrt{1 - q_0} \sqrt{h^2 - x^2}|_{\partial\Omega},$$

by Theorem 2.5, $u \leq \sqrt{1 - q_0} \sqrt{h^2 - x^2} = \sqrt{1 - q_0} \sqrt{(1 + by)^8 f^2 - x^2}$ in Ω ; the result then follows by letting $b \rightarrow 0$.

Theorem 2.10. Let f, Ω , and u be as in Theorem 2.9, $\lim_{y \rightarrow \infty} f(y) = +\infty$, and $p(f) \geq p_0 > 0$, where $1 \geq p_0 > 0$ is a constant. Suppose $u|_{\partial\Omega} \leq \sqrt{1 - p_0} \sqrt{f^2 - x^2}$. Then $u \leq \sqrt{1 - p_0} \sqrt{f^2 - x^2}$ in Ω .

Proof. Case 1: $0 < p_0 < 1$. Let $q_0 = \min(1/8, p_0/2)$. Then

$$u|_{\partial\Omega} \leq \sqrt{1 - p_0} \sqrt{f^2 - x^2}|_{\partial\Omega} \leq \sqrt{1 - q_0} \sqrt{f^2 - x^2}|_{\partial\Omega}.$$

By Theorem 2.9, $u \leq \sqrt{1 - q_0} \sqrt{f^2 - x^2}$ in Ω . Let

$$F = \sqrt{f^2 - x^2} \sqrt{1 - bh(y)^{-\alpha}},$$

where α is some constant in $(0, 1)$ to be determined later, ε is a constant such that $0 < \varepsilon < p_0$ and $b = p_0 - \varepsilon$, and $h(y) = f(y_0 + y)$, where y_0 is a positive constant. Since $f' > 0$ and $\lim_{y \rightarrow \infty} f = +\infty$, we can choose $y_0 > 0$ such that $h \geq 1$ for every $y > 0$. Let

$$\begin{aligned} G &= f^2 - x^2, & g &= (1 - bh^{-\alpha})^{1/2}, \\ g' &= \frac{1}{2}(1 - bh^{-\alpha})^{-1/2}(-b)(-\alpha)h^{-\alpha-1}h' \\ &= \frac{1}{2}(1 - bh^{-\alpha})^{-1/2}b\alpha h^{-\alpha-1}h', \\ g'' &= -\frac{1}{4}(1 - bh^{-\alpha})^{-3/2}(-b)(-\alpha)h^{-\alpha-1}h'b\alpha h^{-\alpha}h' \\ &\quad + \frac{b\alpha}{2}(1 - bh^{-\alpha})^{-1/2}((-\alpha - 1)h^{-\alpha-2}h'^2 + h^{-\alpha-1}h''). \end{aligned}$$

But $p(h) \geq p_0 > 0$, so $h'^2 \geq hh''$. Hence $g'' \leq 0$. By Lemma 2.1,

$$\begin{aligned} G^{3/2}\mathfrak{M}F &= g \left(-f^2 f'^2 + (f^2 - x^2)(f'^2 + f f'') - x^2 - (f^2 - x^2) \right. \\ &\quad \left. + (f^2 - x^2)2f f' \frac{g'}{g} + G^2 \frac{g''}{g} \right) \\ &\quad + g^3 \left(x^2(f'^2 + f f'') - f^2 f'^2 \right. \\ &\quad \left. + x^2 \left(G \frac{g''}{g} - 3G \left(\frac{g'}{g} \right)^2 \right) - \left(G_y G \frac{g'}{g} + G^2 \left(\frac{g'}{g} \right)^2 \right) \right) \\ &\leq g(-f^2 f'^2 + (f^2 - x^2)(f'^2 + f f'') \\ &\quad - x^2 - (f^2 - x^2) + 2f f' \frac{g'}{g}(f^2 - x^2)) \\ &\quad + g^3(x^2(f'^2 + f f'') - f^2 f'^2) \\ &\leq (-g + g^3)(2 - p(f))x^2 f'^2 + f^2 f'^2 g((1 - p(f)) - g^2) \\ &\quad - g f^2 + 2f f' g'(f^2 - x^2) \\ &\leq -(g - g^3)(2 - p(f))x^2 f'^2 - f^2 f'^2 g \max(p(f) - 2, 0) \\ &\quad + f^2 f'^2 g((1 - \min(2, p(f)) - g^2) - g f^2 + 2f f' g'(f^2 - x^2)). \end{aligned}$$

Since $1 \geq g \geq \sqrt{1 - b}$,

$$\begin{aligned} G^{3/2}\mathfrak{M}F &\leq f^2 f'^2 g(1 - \min(2, p(f)) - g^2) + 2f f' g' f^2 \\ &\leq f^2 f'^2 g(-\min(2, p(f)) + bh^{-\alpha}) + 2f^3 f' \frac{1}{2}(1 - bh^{-\alpha})^{-1/2}b\alpha h^{-\alpha-1}h' \\ &\leq f^2 f'^2 \sqrt{1 - b}(-\min(2, p(f)) + bh^{-\alpha}) + b\alpha h^{-\alpha-1}h' \frac{1}{\sqrt{1 - b}} f^3 f' \\ &\leq f^2 f'^2 \sqrt{1 - b}(-\min(2, p(f)) + b)h^{-\alpha} + b\alpha h^{-\alpha-1}h' \frac{1}{\sqrt{1 - b}} f^3 f' \\ &\leq h^{-\alpha} f^2 f'^2 \left(-\sqrt{1 - b}\varepsilon + \alpha b h^{-1} h' \frac{1}{\sqrt{1 - b}} \frac{f}{f'} \right). \end{aligned}$$

Since $(f/f')' = p(f) > 0$, f/f' is monotone increasing. But $h(y) = f(y_0 + y)$, y_0 is a positive constant; therefore $\frac{h'}{h} \frac{f}{f'} \leq 1$. So when $0 < \alpha < (1-b)\varepsilon/b$, we have $G^{3/2}\mathfrak{M}F \leq 0$. Since

$$\lim_{y \rightarrow \infty} f(y) = +\infty, \quad \lim_{y \rightarrow \infty} \sqrt{1 - bh(y)^{-\alpha}} \rightarrow 1,$$

we have $F(x, y) \geq \sqrt{1 - q_0} \sqrt{f^2 - x^2}$ for sufficiently large y . Since $u \leq \sqrt{1 - q_0} \sqrt{f^2 - x^2}$ in Ω and $u|_{\partial\Omega} \leq F|_{\partial\Omega}$ by hypothesis, we have $u \leq F$ in Ω . So $u \leq \sqrt{1 - bh^{-\alpha}} \sqrt{f^2 - x^2}$ in Ω . Letting $\alpha \rightarrow 0$, $u \leq \sqrt{1 - b} \sqrt{f^2 - x^2} = \sqrt{1 - (p_0 - \varepsilon)} \sqrt{f^2 - x^2}$ in Ω . Letting $\varepsilon \rightarrow 0$, we get $u \leq \sqrt{1 - p_0} \sqrt{f^2 - x^2}$ in Ω .

Case 2: $p_0 = 1$. Then for any constant p_1 , $0 < p_1 < 1$, we have $p(f) \geq 1 \geq p_1$. Since $u|_{\partial\Omega} \leq \sqrt{1 - 1} \sqrt{f^2 - x^2} \leq \sqrt{1 - p_1} \sqrt{f^2 - x^2}$, by Case 1, we have $u \leq \sqrt{1 - p_1} \sqrt{f^2 - x^2}$ in Ω . Letting $p_1 \rightarrow 1$, the result follows.

Since the case $\lim f < +\infty$ is not very important, we omit that case.

The case for a negative constant p_0 is studied in the following theorem.

Theorem 2.11. Let $f \in C^0[0, \infty) \cap C^2(0, \infty)$, $f \geq 0$, $f' > 0$, $f_1 \in C^0[0, \infty)$, $f_1 > 0$, $\Omega \subset \{(x, y) | -f_1(y) < x < f_1(y), y > 0\}$, $u \in C^2(\Omega) \cap C^0(\bar{\Omega})$, $\operatorname{div} Tu \geq 0$ in Ω , and $p(f) \geq p_0$, where p_0 is a negative constant. Then if $f^2 \geq (a^2 - 1)(2 - p_0)f_1^2/(a^2 - (1 - p_0))$ and $u|_{\partial\Omega} \leq a\sqrt{f^2 - x^2}|_{\partial\Omega}$, where a is a positive constant satisfying $a^2 - 1 + p_0 > 0$, we have $u \leq a\sqrt{f^2 - x^2}$ in Ω .

Proof. Let $f_2 = e^{\alpha y} f$, where α is a positive constant. Then $f_2^2 \geq f^2 \geq (a^2 - 1)(2 - p_0)f_1^2/(a^2 - (1 - p_0))$. By Lemma 2.6, $p(f_2) \geq p_0$. It is easy to see that $\limsup_{y \rightarrow \infty} (f_2/y^2) = +\infty$, by Lemma 2.8, and $\liminf_{y \rightarrow \infty} (f_2/f_2'^2) = 0$. Since $u|_{\partial\Omega} \leq a\sqrt{f^2 - x^2}|_{\partial\Omega} \leq a\sqrt{f_2^2 - x^2}|_{\partial\Omega}$, by Theorem 2.5, $u \leq a\sqrt{f_2^2 - x^2} = a\sqrt{e^{2\alpha y} f^2 - x^2}$ in Ω . The result then follows by letting $\alpha \rightarrow 0$.

Let $\Omega \subset \{-f_1(y) < x < f_1(y) | y > 0\}$ and $u|_{\partial\Omega} \leq af_1$, where a is a positive constant. The growing rate of u is stated in the following theorem.

Theorem 2.12. Let $f_1 \in C^0[0, \infty) \cap C^2(0, \infty)$, $f_1, f_1' > 0$, $\lim f_1 = \infty$, and $p(f_1) \geq p_0 \geq 0$, where $1 \geq p_0 \geq 0$ is a constant. Moreover, let $\Omega \subset \{(x, y) | -f_1(y) < x < f_1(y), y > 0\}$ and $f = af_1$, where $a \geq 0$ is a constant. Then if $\operatorname{div} Tu \geq 0$ in Ω and $u|_{\partial\Omega} \leq f$, we have $u \leq \sqrt{(a^2 + 1 - p_0)f_1^2 + (1 - p_0)x^2}$ in Ω .

Proof. *Case 1:* $1 > p_0 \geq 0$. Since

$$u|_{\partial\Omega} \leq f = af_1 \leq \sqrt{1 - p_0} \sqrt{\frac{a^2 f_1^2}{1 - p_0} + f_1^2 - x^2}|_{\partial\Omega}$$

and $p(\sqrt{\frac{a^2}{1 - p_0} + 1} f_1) = p(f) \geq p_0$,

$$\begin{aligned} u &\leq \sqrt{1 - p_0} \sqrt{\frac{a^2}{1 - p_0} f_1^2 + f_1^2 - x^2} \\ &= \sqrt{(a^2 + (1 - p_0))f_1^2 - (1 - p_0)x^2} \quad \text{in } \Omega \end{aligned}$$

by Theorem 2.10.

Case 2: $p_0 = 1$. For every constant b where $1 \geq b \geq 0$, by Case 1, we have $u \leq \sqrt{(a^2 + (1-b))f_1^2 - (1-b)x^2}$ in Ω . The result then follows by letting $b \rightarrow 0$.

Now we assume that $p(f_1) \geq p_0$ with p_0 being a negative constant, and set $f^2 = (a^2 - 1)(2 - p_0)f_1^2 / (a^2 - (1 - p_0))$, where $a > \sqrt{1 - p_0}$ is a constant. If $u|_{\partial\Omega} \leq a\sqrt{f^2 - x^2}|_{\partial\Omega} \leq a\sqrt{(a^2 - 1)(2 - p_0)/(a^2 - (1 - p_0)) - 1}f_1$, by Theorem 2.11, we have $u \leq a\sqrt{f^2 - x^2}$ in Ω . Now we want to compute the minimum of $a\sqrt{(a^2 - 1)(2 - p_0)/(a^2 - (1 - p_0)) - 1}$. For convenience, let $b = a^2$ and $q_1 = 1 - p_0 > 1$, and we need to compute the minimum of

$$b \left(\frac{(b-1)(1+q_1)}{b-q_1} - 1 \right) = b \frac{q_1 b - 1}{b - q_1}.$$

Take logarithms and differentiate the above function; we have $1/(b+q_1/(q_1 b - 1)) = 1/(b - q_1)$. From this, we get $b = q_1 \pm \sqrt{q_1^2 - 1}$. Since $b = a^2 > q_1$, we have $b = q_1 + \sqrt{q_1^2 - 1}$. So the minimum of $b(q_1 b - 1)/(b - q_1)$ is

$$(q_1 + \sqrt{q_1^2 - 1}) \left(\frac{q_1^2 + q_1 \sqrt{q_1^2 - 1} - 1}{\sqrt{q_1^2 - 1}} \right) = (q_1 + \sqrt{q_1^2 - 1})^2,$$

and the minimum of $a\sqrt{(a^2 - 1)(2 - p_0)/(a^2 - (1 - p_0)) - 1}$ is $q_1 + \sqrt{q_1^2 - 1}$.

Theorem 2.13. Assume that $f_1, f'_1 > 0$, $p(f_1) \geq p_0$, where p_0 is a negative constant, $\Omega \subset \{-f_1(y) < x < f_1(y) | y > 0\}$, and $u \in C^2(\Omega) \cap C^0(\bar{\Omega})$. If $\operatorname{div} Tu \geq 0$ in Ω , $q_1 = 1 - p_0 > 1$, and $u|_{\partial\Omega} \leq b f_1$, where $b \geq q_1 + \sqrt{q_1^2 - 1}$ is a positive constant, then $u \leq \sqrt{b} \sqrt{(b+1)f^2 - x^2}$ in Ω .

Proof. By the assumption that $b \geq q_1 + \sqrt{q_1^2 - 1}$, we have

$$(b - (q_1 + \sqrt{q_1^2 - 1}))(b - (q_1 - \sqrt{q_1^2 - 1})) \geq 0,$$

$b^2 - 2q_1 b + 1 \geq 0$, and $(b+1)(b-q_1) \geq (b-1)(1+q_1)$. If we set $a = b^2$, then $(a^2+1)(a^2-q_1) \geq (a^2-1)(1+q_1)$ and we have $a^2+1 \geq (a^2-1)(1+q_1)/(a^2-q_1)$. Since $u|_{\partial\Omega} \leq a\sqrt{(a^2+1)f_1^2 - x^2}|_{\partial\Omega}$, by Theorem 2.11, $u \leq a\sqrt{(a^2+1)f_1^2 - x^2}$ in Ω and we have $u \leq \sqrt{b} \sqrt{(b+1)f^2 - x^2}$ in Ω .

By Theorems 2.12 and 2.13, we know that the growth rate of u is of the same order as the shape of Ω and $u|_{\partial\Omega}$.

ACKNOWLEDGMENT

The author would like to thank the referee for many helpful comments and suggestions.

REFERENCES

1. R. Finn, *Equilibrium capillary surfaces*, Springer-Verlag, New York, Berlin, Heidelberg, and Tokyo, 1986.
2. D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, second ed., Springer-Verlag, Berlin, Heidelberg, New York, and Tokyo, 1983.
3. J. Hwang, *Phragmén-Lindelöf theorem for the minimal surface equation*, Proc. Amer. Math. Soc. **104** (1988), 825–828.
4. J. C. C. Nitsche, *On new results in the theory of minimal surfaces*, Bull. Amer. Math. Soc. **71** (1965), 195–270.
5. ———, *Vorlesungen über Minimalflächen*, Springer-Verlag, Berlin, Heidelberg, and New York, 1975.
6. R. Osserman, *A survey of minimal surfaces*, Van Nostrand-Reinhold, New York, 1969.
7. M. H. Protter and H. F. Weinberger, *Maximum principles in differential equations*, Prentice-Hall, Englewood Cliffs, NJ, 1967.

INSTITUTE OF MATHEMATICS, ACADEMIA SINICA, TAIPEI 11529, TAIWAN, REPUBLIC OF CHINA