NEAR-RINGS ASSOCIATED WITH MATCHED PAIRS ON RING MODULES

C. J. MAXSON AND A. P. J. VAN DER WALT

(Communicated by Lance W. Small)

ABSTRACT. Let G be a module over a ring R , let $\mathcal{C} = \{C_i\}$, $i \in I$, be a family of submodules of G , and let $\mathcal{H} = \{H_i\}$, $i \in I$, where H_i is a subgroup of $\text{Hom}_R(C_i, G)$ with certain properties. To each such pair $(\mathcal{C}, \mathcal{H})$, a near-ring $M(\mathcal{C}, \mathcal{H})$ is associated, which is a generalization of the near-ring of homogeneous functions determined by (G, R) . The transfer of information from module properties of G_R reflected in $(\mathcal{C}, \mathcal{H})$ to structural properties of $M(\mathcal{C}, \mathcal{H})$ is investigated.

I. INTRODUCTION AND BASIC CONCEPTS

Let R be a ring with identity, and let G be a unitary right R -module. Then, under function addition and function composition, the set $M_R(G) := \{f : G \rightarrow G \mid f(ar) = f(a)r, \forall a \in G, \forall r \in R\}$ is a zero-symmetric near-ring with identity, called the near-ring of homogeneous functions determined by the pair (R, G) . The ring $S := \text{End}_R G$ is a subring of $M_R(G)$. The near-rings $M_R(G)$ have been the subject of several investigations. (See [2] and the references given there.) Recently, sub-near-rings of $M_R(G)$ in which the homogeneous functions can be represented locally as endomorphisms of G have been investigated. We proceed to define these subrings.

Let $\mathcal{C} = \{C_\alpha\}$, $\alpha \in \mathcal{A}$, be a family of submodules of G with the properties: (i) $\bigcup \mathcal{C} = \bigcup_{\alpha \in \mathcal{A}} C_\alpha = G$, and (ii) for each $s \in S$ and each $C_\alpha \in \mathcal{C}$, there exists $\beta \in \mathcal{A}$ such that $s(C_\alpha) \subseteq C_\beta$. In this case we say $\mathcal{C}$ is a cover for G . For each cover $\mathcal{C}$ of G we have a sub-near-ring $PE_R(G, \mathcal{C})$ of $M_R(G)$ given by $PE_R(G, \mathcal{C}) := \{f \in M_R(G) \mid f|_{C_\alpha} \text{ extends to some } s \in \text{End}_R G, \forall \alpha \in \mathcal{A}\}$, called the near-ring of piecewise endomorphisms of G_R determined by $\mathcal{C}$ [3, 4]. In [3] it was shown that for finitely generated modules G over a principal ideal domain D , if one uses the cover $\mathcal{P}$ of cyclic submodules then $M_D(G) = PE_D(G, \mathcal{P})$, i.e., every homogeneous function is locally an endomorphism of G . It is not known if this characterizes finitely generated modules over principal ideal domains.

It is the purpose of this work to initiate an investigation of a generalization of near-rings of piecewise endomorphisms. The main goal of our program of study

Received by the editors September 2, 1992 and, in revised form, February 8, 1993.

1991 *Mathematics Subject Classification.* Primary 16Y30; Secondary 16D10.

©1994 American Mathematical Society
 0002-9939/94 \$1.00 + \$.25 per page

is to develop concepts and techniques which may then be applied to near-rings of piecewise endomorphisms to obtain a better understanding of this situation.

We now present the basic definitions and concepts for our work. Throughout the paper, all rings R will have an identity and all R -modules G will be unitary.

Let $\mathcal{C} = \{C_i\}$, $i \in I$, be a family of submodules of G , where I is an index set and $C_i \neq C_j$ when $i \neq j$. Let $\mathcal{H} = \{H_i\}$, $i \in I$, be a family with H_i subgroups of $\text{Hom}_R(C_i, G)$. The pair $(\mathcal{C}, \mathcal{H})$ is called a *matched pair* on G if: (a) $f \in H_i$ implies $f(C_i) \in \mathcal{C}$, and (b) $f \in H_i$ with $fC_i = C_j$ and $g \in H_j$ implies $gf \in H_i$.

In this definition, as in the sequel unless otherwise stated, unquantified indices like i and j will be understood to range freely over I . We shall find it useful to let $H := \bigcup \mathcal{H}$ and to define the functions $\sigma, \tau : H \rightarrow I$ by $\sigma(h) = i$ if $h \in H_i$ and $\tau(h) = j$ if $hC_{\sigma(h)} = C_j$. This is well defined because $C_i \neq C_j$ when $i \neq j$. Also in the sequel we assume without specific mention that we have some matched pair $(\mathcal{C}, \mathcal{H})$ with accompanying index set I .

Associated with each matched pair $(\mathcal{C}, \mathcal{H})$ on G is a near-ring $M := M(\mathcal{C}, \mathcal{H})$ which we proceed to define. For our set M we take $M = \times_i H_i = \{\{s_i\}, i \in I | s_i \in H_i\}$. Addition on M is defined to be componentwise, and multiplication on M is defined for $s, t \in M$ by $(st)_i := s_{\tau(t_i)}t_i$.

Theorem I.1. $M = M(\mathcal{C}, \mathcal{H})$ is a zero-symmetric abelian near-ring with the property $s(-t) = -(st)$, $\forall s, t \in M$.

Proof. Clearly $(M, +)$ is an abelian group. We verify right distributivity and associativity. Right distributivity follows from

$$\begin{aligned} [(s+t)u]_i &= (s+t)_{\tau(u_i)}u_i = (s_{\tau(u_i)} + t_{\tau(u_i)})u_i \\ &= s_{\tau(u_i)}u_i + t_{\tau(u_i)}u_i = (su)_i + (tu)_i \end{aligned}$$

and associativity from

$$\begin{aligned} [s(tu)]_i &= s_{\tau((tu)_i)}(tu)_i = s_{\tau(t_{\tau(u_i)})}t_{\tau(u_i)}u_i \\ &= (st)_{\tau(u_i)}u_i = [(st)u]_i. \end{aligned}$$

Finally, if $h, k \in H$ and hk is defined, then $h(-k) = -hk$. This completes the proof.

Corollary I.2. If $1_{C_i} \in H_i$ for all $i \in I$, then M is a near-ring with identity e , where $e_i := 1_{C_i}$.

Unless stated to the contrary we always assume $1_{C_i} \in H_i$ for all $i \in I$. This implies that $\varepsilon^i \in M$ where $(\varepsilon^i)_j := 0$ if $j \neq i$ and $(\varepsilon^i)_i = e_i$.

Theorem I.3. The ε^i are mutually orthogonal.

Suppose $\mathcal{P} = \{C_i\}$, $i \in I$, is the collection of all cyclic submodules of an R -module G . If we let $\text{Hom} := \{H_i\}$, $i \in I$, where $H_i := \text{Hom}_R(C_i, G)$, then $(\mathcal{P}, \text{Hom})$ is a matched pair on G . However, $M(\mathcal{P}, \text{Hom})$ does not necessarily represent a near-ring of functions on G since, for $s \in M(\mathcal{P}, \text{Hom})$ and $x \in C_i \cap C_j$, $s_i x$ may be different from $s_j x$. Consequently we define $HF(\mathcal{P}, \text{Hom}) := \{s \in M | s_i | C_i \cap C_j = s_j | C_i \cap C_j, \forall i, j \in I\}$. Straightforward calculations show that $HF(\mathcal{P}, \text{Hom})$ is a sub-near-ring of $M(\mathcal{P}, \text{Hom})$. Then $HF(\mathcal{P}, \text{Hom})$ can be thought of as a near-ring of functions of G by

defining $s(g) = s_i g$ if $g \in C_i$ and $s \in HF(\mathcal{P}, \text{Hom})$. Moreover, since $s_i \in \text{Hom}_R(C_i, G)$, we have $s(gr) = (sg)r$, for each $r \in R$. So, we have a near-ring of homogeneous functions on G . A sub-near-ring, $PE(\mathcal{P}, \text{Hom})$ of $HF(\mathcal{P}, \text{Hom})$ is defined by $PE(\mathcal{P}, \text{Hom}) := \{s \in HF(\mathcal{P}, \text{Hom}) | s_i = \rho|_{C_i} \text{ for some } \rho \in \text{End}_R G\}$. This is a near-ring of piecewise endomorphisms on G which we denoted previously by $PE_R(G, \mathcal{P})$. Thus, in this sense, our near-rings $M(\mathcal{E}, \mathcal{H})$ are indeed generalizations of the near-rings of piecewise endomorphisms.

For $h, g \in H$, if hg is defined, then one has $\sigma(hg) = \sigma(g)$ and $\tau(hg) = \tau(h)$. Now let $s \in M$, and define $\text{suppt}(s) = \{\sigma(s_i) | s_i \neq 0\}$, $\text{null}(s) = \{\sigma(s_i) | s_i = 0\}$, and $\text{target}(s) = \{\tau(s_i) | i \in I\}$. We then have the following lemma, whose proof is left to the reader.

Lemma I.4. For $s, t \in M$,

- (i) $\text{suppt}(st) \subseteq \text{suppt}(t)$,
- (ii) $\text{null}(st) \supseteq \text{null}(t)$,
- (iii) $\text{target}(st) \subseteq \text{target}(s)$, and
- (iv) $|\text{target}(st)| \leq |\text{target}(t)|$.

As an application of the above we take $R = \mathbb{Z}$ and $G = \mathbb{Z}_{\mathbb{Z}}$ and let $\mathcal{E}_{\mathbb{Z}} = \{C_i\}$ be the set of all ideals of $\mathbb{Z}$ where the indexing set $I = \mathbb{N}_0$ and $C_i = \mathbb{Z}i$, the ideal generated by i . Further let $H_i := \text{Hom}_{\mathbb{Z}}(C_i, \mathbb{Z})$.

Lemma I.5. If $s, t \in M(\mathcal{E}_{\mathbb{Z}}, H)$ are such that $|\text{target}(s)| < \infty$ and $|\text{target}(t)| < \infty$, then $|\text{target}(s+t)| < \infty$.

Proof. Define an equivalence relation $\sim$ on I by $i \sim j$ if and only if $\tau(s_i) = \tau(s_j)$ and $\tau(t_i) = \tau(t_j)$. By hypothesis, $\sim$ has finitely many equivalence classes. Let J be one such class. We show that if $i \in J$, then there are just finitely many possibilities for $\tau((s+t)_i)$. If $\tau(s_i) = j$ then $s_i(i) = j$ or $-j$, and likewise if $\tau(t_i) = k$ then $t_i(i) = k$ or $-k$. Hence $\tau((s+t)_i) = |\pm j \pm k|$.

Recall that an additive subgroup A of an arbitrary near-ring is invariant if $NA \subseteq A$ and $AN \subseteq A$. If the only invariant subgroups of N are $\{0\}$ and N , we say N is *invariantly simple*. The previous two lemmas show that $\{s \in M(\mathcal{E}_{\mathbb{Z}}, \mathcal{H}) | |\text{target}(s)| < \infty\}$ is a nontrivial proper invariant subgroup of $M(\mathcal{E}_{\mathbb{Z}}, \mathcal{H})$.

Theorem I.6. $M(\mathcal{E}_{\mathbb{Z}}, \mathcal{H})$ is not invariantly simple.

We remark that this theorem can be generalized to rings in which each nonzero element has a finite number of associates, e.g. $\mathbb{Z}[x]$, $\mathbb{Z}[x_1, \dots, x_n]$, and $F[x_1, \dots, x_n]$ where F is a finite field.

On the other hand we now show that $M(\mathcal{E}_{\mathbb{Z}}, \mathcal{H})$ is a simple near-ring. We note that this also follows from subsequent results.

Theorem I.7. $M(\mathcal{E}_{\mathbb{Z}}, \mathcal{H})$ is a simple near-ring.

Proof. We adopt the following convention. If $s \in M(\mathcal{E}_{\mathbb{Z}}, \mathcal{H})$, $s_i = k$ will mean $s_i(i) = k$. Now suppose $s \neq 0$, say $s_m = k$. We may assume $k > 0$; otherwise use $-s$. Let x and y be defined by $x_i = m$, $y_i = 1$ for $i > 0$. Then $(ysx)_i = 1$ for $i > 0$. Now let u and v be defined by $u_i = i - 1$, $v_i = i(i+1)/2$, $i > 0$. Then $ysx + u = e$, $v(ysx + u) = v$, and $(vu)_i = v_{i-1}$,

$i > 0$. Hence $[v(ysx + u) - vu]_i = v_i - v_{i-1} = i$, $i > 0$, so e is in the ideal generated by s .

We now present a very general way of constructing matched pairs on an R -module G . To this end let $\mathcal{A}$ be any class of right R -modules, and set

$$\mathcal{E}_G(\mathcal{A}) := \{\text{Im } f \mid f \in \text{Hom}_R(A, G), A \in \mathcal{A}\}.$$

If $\mathcal{A} = \{A\}$, we write $\mathcal{E}_G(A)$ for $\mathcal{E}_G(\{A\})$. Suppose $\mathcal{E}_G(\mathcal{A}) = \{C_i\}$, $i \in I$, where this indexing is done in such a way that $C_i \neq C_j$ when $i \neq j$. Two families of sets of mappings are natural candidates to be the second member of the matched pair, viz. $\text{Hom} := \{\text{Hom}_R(C_i, G)\}$, $i \in I$, and $\text{End} := \{\{f \mid C_i \mid f \in \text{End}_R G\}\}$, $i \in I$. However, there are many other possibilities, e.g., if T is any subring of $\text{End}_R G$, then $\{\{f \mid C_i \mid f \in T\}\}$, $i \in I$, is a candidate for $\mathcal{H}$. We note that if $(\mathcal{E}, \text{Hom})$ is a matched pair, then $\mathcal{E} = \mathcal{E}_G(\mathcal{E})$. The next result states a number of obvious facts about matched pairs.

Theorem I.8. (a) Let $(\mathcal{E}, \mathcal{H})$ and $(\mathcal{E}', \mathcal{H}')$ be matched pairs on G . If $\mathcal{E} \subseteq \mathcal{E}'$ and $H \subseteq \mathcal{H}'$, then $M(\mathcal{E}, \mathcal{H})$ is a sub-near-ring of $M(\mathcal{E}', \mathcal{H}')$.

(b) $\mathcal{E}_G(R)$ is the family of all cyclic submodules of G .

(c) $\mathcal{E}_G(\mathcal{S})$, where $\mathcal{S}$ is the class of simple R -modules, is the family of simple submodules of G .

Let $(\mathcal{E}, \mathcal{H})$ be a matched pair on G with relevant indexing set I . We define a relation $\sim$ on I by $i \sim j$ if there exist $h, k \in H$ with $\sigma(h) = \tau(k) = i$ and $\tau(h) = \sigma(k) = j$. It is straightforward to verify that $\sim$ is an equivalence relation. We denote the set of equivalence classes by $I/\sim$ and represent an equivalence class of $\sim$ by $[i]$. We next define a relation $\preceq$ on $I/\sim$ by $[i] \preceq [j]$ if there exists an $h \in H$ with $\sigma(h) \in [j]$ and $\tau(h) \in [i]$. Using the definition of matched pair one verifies that $\preceq$ is a partial order on $I/\sim$.

Theorem I.9. If every $h \in H_i \cap \text{End}_R C_i$ is an automorphism of C_i , then every h with $\sigma(h) \in [i]$ and $\tau(h) \in [i]$ is an isomorphism.

Proof. If $\sigma(h), \tau(h) \in [i]$, then there exist $h_1, h_2 \in H$ with $\sigma(h_1) = i$, $\tau(h_1) = \sigma(h)$, $\sigma(h_2) = \tau(h)$, and $\tau(h_2) = i$. This implies that $h_2 h h_1$ is an endomorphism of C_i and, so by hypothesis, an automorphism. But this in turn implies h is an isomorphism.

Corollary I.10. Let G be Noetherian. Then every $h \in H$ with $\sigma(h), \tau(h) \in [i]$ for some $i \in I$ is an isomorphism. In particular, if $[i]$ is minimal with respect to $\preceq$, then every nonzero $h \in H$ with $\sigma(h) \in [i]$ is an isomorphism.

Proof. Since the mappings in $\mathcal{H}$ are all epimorphisms, the result follows from the above theorem and the well-known theorem that an onto endomorphism of a Noetherian module is an automorphism.

For future use we note that in the case when $\mathcal{H} = \text{Hom}$, if $h \in H$ is an isomorphism, then $h^{-1} \in H$. Hence the above corollary says

Corollary I.11. Let G be Noetherian, let $(\mathcal{E}, \mathcal{H})$ be a matched pair on G , and let $[i]$ be minimal. If $h \in H$ is nonzero with $\sigma(h) \in [i]$, then $h^{-1} \in H$.

Another easy result in this vein is

Theorem I.12. *If $C_i \in \mathcal{C}$ is minimal in G , then $[i]$ is minimal and every $h \in H$ with $\sigma(h), \tau(h) \in [i]$ is an isomorphism.*

Convention. Henceforth, for convenience of exposition we take $0 \in I$ and $C_0 = \{0\}$. Note that $[0] = \{0\}$.

We conclude this section by characterizing when $M(\mathcal{C}, \mathcal{H})$ is a ring. From this, we note that, in general, $M(\mathcal{C}, \mathcal{H})$ is not a ring.

Theorem I.13. *Let $(\mathcal{C}, \mathcal{H})$ be a matched pair on G . The following are equivalent:*

- (1) $M(\mathcal{C}, \mathcal{H})$ is a ring.
- (2) For each nonzero $h \in H$, $\sigma(h) = \tau(h)$.
- (3) $\forall i \in I \setminus \{0\}$, $[i]$ is minimal and $[i] = \{i\}$.

Proof. From the definitions, one observes that (2) and (3) are equivalent, so we show (1) and (2) are equivalent. Let $M(\mathcal{C}, \mathcal{H})$ be a ring, and assume there is $0 \neq h \in H$ with $h: C_j \rightarrow C_i$, $i \neq j$. Define $t \in M$ by $t_j = h$ and $t_i = 0$, $i \neq j$. Since M is a ring, $\varepsilon^i(t + \varepsilon^j) = \varepsilon^i t + \varepsilon^i \varepsilon^j = t$. Let $\tau(t_j + (\varepsilon^j)_j) = l$. Then for each $x \in C_j$, $[\varepsilon^i(t + \varepsilon^j)](x) = (\varepsilon^i)_l(h(x) + x) = t_j(x) = h(x)$. If $l \neq i$, we get $0 = h(x)$, which is a contradiction since $0 \neq h$. If $l = i$, we have $h(x) + x = h(x)$, so $x = 0$, again a contradiction since $C_j \neq \{0\}$. Hence if $M(\mathcal{C}, \mathcal{H})$ is a ring, we must have $\sigma(h) = \tau(h)$, for each $0 \neq h \in H$. Conversely, we note that if $\sigma(h) = \tau(h)$ for each nonzero $h \in H$, then H_i is a subring of $\text{End}_R C_i$ and $M(\mathcal{C}, \mathcal{H}) = \times_I H_i$ with pointwise addition and multiplication, i.e., $M(\mathcal{C}, \mathcal{H})$ is a ring.

II. LEFT IDEALS

Recall that a left ideal L of an arbitrary near-ring N is *strictly minimal* (*strictly maximal*) if L is minimal (maximal) as a left invariant subgroup. In this section we identify some minimal left ideals and strictly minimal left ideals of $M = M(\mathcal{C}, \mathcal{H})$.

For any subset J of I , the set $A(J) := \{s \in M \mid s_j = 0 \text{ if } j \in J\}$ is a left ideal of M . In particular, every $M\varepsilon^i$ is a left ideal since $M\varepsilon^i = A(I \setminus \{i\})$. We investigate when $M\varepsilon^i$ is minimal, and to this end we transfer the inclusion partial order on $\mathcal{C}$ to I , writing $\preceq$ for the resulting partial order on I .

Theorem II.1. *If $[i]$ is minimal with respect to $\preceq$ and there exists $j \in [i]$ such that $j \not\preceq i$, then $M\varepsilon^i$ is a minimal left ideal of M .*

Proof. Let L be a nonzero left ideal of M with $L \subseteq M\varepsilon^i$, and take $0 \neq s \in L$. Then $s_i \neq 0$. We claim that without loss of generality we may assume $\tau(s_i) = j$. For, since $[i]$ is minimal, $\tau(s_i) \in [i]$, so there is some $h \in H$ such that $\tau(hs_i) = j$. If $t \in M$ is such that $(t)_{\tau(s_i)} = h$, then $ts \in L$ and $\tau((ts)_i) = j$. Now let $x \in M$ be arbitrary with $s_i + x_i = e_i$. Therefore, $\tau(x_i) \neq i$; otherwise, for each $c \in C_i$, $s_i(c) = c - x_i(c) \in C_i$, which implies $C_j \subseteq C_i$, i.e., $j \preceq i$. Let $y \in M$ be defined by $y_i = e_i$ and $y_k = 0$ for $k \neq i$. Then $z := y(s+x) - yx \in L$ and $z_i = y_i(s_i + x_i) - y_{\tau(x_i)}x_i = e_i$, so $z = \varepsilon^i$. But this means $\varepsilon^i \in L$ and $L = M\varepsilon^i$.

Corollary II.2. *If $[i]$ is minimal with respect to $\preceq$, then $M\varepsilon^k$ is a minimal left ideal of M for all but (possibly) one $k \in [i]$.*

Proof. If there is more than one element in $[i]$, then for at most one $k \in [i]$ can we have that $j \leq k$ for all $j \in [i]$.

To illustrate the above, let G be any torsion free abelian group regarded as a $\mathbb{Z}$ -module, and consider the matched pair $(\mathcal{E}_G(\mathbb{Z}), \text{Hom})$. Then I has just two equivalence classes, $[0] = \{0\}$ and $I \setminus \{0\}$, the latter being minimal. If G is the infinite cyclic group, then $G \in \mathcal{E}_G(\mathbb{Z})$, say $G = C_k$. Then $j \leq k$ for all $j \in I \setminus \{0\}$, but for each $j \in I \setminus \{0\}$, $j \neq k$ we have $k \not\leq j$, so $M\epsilon^j$ is minimal for each $j \in I \setminus \{0\}$, $j \neq k$. We note that, in this case, it turns out that $M\epsilon^k$ is also minimal (see Theorem 2.5). By contrast, let $G := Q$ and $R := \mathbb{Z}$, and consider the pair $(\mathcal{E}, \mathcal{H})$ where $\mathcal{E} := \{C_0 := \{0\}, C_1 := Q\}$ and $\mathcal{H} := \{0, \mathbb{Z}\}$. Then $M(\mathcal{E}, \mathcal{H}) = M\epsilon^1 \cong \mathbb{Z}$, so $M\epsilon^1$ is not minimal; in fact, M has no minimal left ideal. Note here that $[i] = \{i\}$ for $i \in I$; so if $[i] = \{i\}$ and $[i]$ is minimal, it need not be the case that $M\epsilon^i$ is minimal.

Theorem II.3. *The left ideal $M\epsilon^i$ is strictly minimal if and only if $h \in H_i$, $h \neq 0$, implies $h^{-1} \in H$.*

Proof. This follows from the fact that for $s \in M$, Ms will be properly contained in $M\epsilon^i$ if and only if $s = s\epsilon^i$ and there is no $t \in M$ such that $ts = \epsilon^i$, i.e., such that $(ts)_i = t_{\tau(s)}s_i = e_i$.

Since $M = A(\{i\}) \oplus M\epsilon^i$, we have the following dual result about maximal left ideals.

Theorem II.4. *Let $[i]$ be minimal with respect to $\preceq$.*

- (a) *If there is $j \in [i]$ with $j \not\leq i$, then $A(\{i\})$ is a maximal left ideal of M .*
- (b) *$A(\{k\})$ is maximal for all but (possibly) one $k \in [i]$. Moreover, the left ideal $A(\{j\})$ is strictly maximal if and only if $h \in H_j$, $h \neq 0$, implies $h^{-1} \in H$.*

Combining Theorem II.3, the last part of Theorem II.4, and Corollary I.11 we obtain

Theorem II.5. *Let G be Noetherian and $(\mathcal{E}, H)$ be a matched pair on G . If $[i]$ is minimal with respect to $\preceq$, then $M\epsilon^j$ is strictly minimal and $A(\{j\})$ is strictly maximal for each $j \in [i]$.*

Recall that a near-ring N is 2-semisimple provided the intersection of all strictly maximal ideals of N is $\{0\}$, i.e., if $J_2(N) = \{0\}$. When G is Noetherian and $[i]$ is minimal for each $i \in I$, then $J_2(M) \subseteq \bigcap_{i \in I} A(\{i\}) = \{0\}$. Thus we have

Corollary II.6. *Under the hypothesis of Theorem II.5, if $[i]$ is minimal with respect to $\preceq$ for each nonzero $i \in I$, then M is 2-semisimple.*

III. TWO-SIDED IDEALS

As noted above, the $A(J)$ are obvious examples of left ideals in $M(\mathcal{E}, \mathcal{H})$. In this section we focus on two-sided ideals. We first determine when the $A(J)$ will be two-sided. We then investigate conditions under which $M = M(\mathcal{E}, \mathcal{H})$ is a simple near-ring.

We say a nonempty set $J \subseteq I$ is a *sink* if $\sigma(h) \in J$ implies $\tau(h) \in J$. Note that both $\{0\}$ and I are sinks which we refer to as the trivial sinks.

Theorem III.1. For $\emptyset \neq J \subseteq I$, the left ideal $A(J)$ is a two-sided ideal if and only if J is a sink.

Proof. Let J be a sink. We show $A(J)$ is a right ideal. If $s \in A(J)$ and $t \in M$, then for $j \in J$, $(st)_j = s_k t_j$ where $\tau(t_j) = k$. Since $j \in J$, we have $k \in J$ and thus, since $s \in A(J)$, $s_k = 0$. On the other hand, if $J \neq \emptyset$ is not a sink, then there exists $h \in H$ with $j := \sigma(h)$ and $k := \tau(h)$ with $k \notin J$. Let $s \in A(J)$ with $s_k = e_k$, and let $t \in M$ with $t_j = h$. Then $(st)_j = s_k t_j = h \neq 0$, so $st \notin A(J)$.

In light of this result it is of some importance to determine the sinks in I . The next result gives some information in this regard.

Theorem III.2. (a) If J is any nonempty subset of I , then $\Sigma(J) := \{\tau(h) | h \in H \text{ and } \sigma(h) \in J\}$ is a sink called the sink generated by J .

(b) If $K := \bigcup \{[i] | [i] \preceq [j] \text{ for some } j \in J \subseteq I\}$, then $K = \Sigma(J)$.

(c) If $\mathcal{A}$ is any class of R -modules and $\mathcal{B} \subseteq \mathcal{A}$, then $\Sigma(\mathcal{B}) := \{i \in I | C_i \in \mathcal{E}_G(\mathcal{B})\}$ is a sink with regard to any matched pair $(\mathcal{E}_G(\mathcal{A}), \mathcal{H})$.

(d) If $\mathcal{J}$ is any family of sinks in I , then $\bigcup \mathcal{J}$ and $\bigcap \mathcal{J}$ are also sinks.

Proof. (a) Suppose $h \in H$ with $\sigma(h) \in \Sigma(J)$. Thus there exists $h' \in H$ with $\sigma(h') \in J$ and $\tau(h') = \sigma(h)$. But then $hh' \in H$ and $\sigma(hh') = \sigma(h') \in J$. Hence $\tau(h) = \tau(hh') \in \Sigma(J)$ as desired.

(b) If $i \in \Sigma(J)$, then there is some $h \in H$ with $\sigma(h) \in J$ and $\tau(h) = i$. Consequently, $[i] \preceq [\sigma(h)]$, so $i \in K$. Conversely, if $i \in K$, then $[i] \preceq [j]$, for some $j \in J$, which in turn implies that there exists $h \in H$ with $\sigma(h) = j$ and $\tau(h) = i$. Hence $i \in \Sigma(J)$.

(c) If $\sigma(h) \in \Sigma(\mathcal{B})$, then $\tau(h) \in \Sigma(\mathcal{B})$ since $h(C_{\sigma(h)}) \in \mathcal{E}_G(\mathcal{B})$.

(d) This is clear.

We apply this theorem to obtain some specific instances of sinks.

Corollary III.3. The following are sinks for any matched pair $(\mathcal{E}, \mathcal{H})$ on G :

(a) $[i] \cup \{0\}$ if $[i]$ is minimal.

(b) $\{i | C_i \text{ has property } \mathcal{P}\}$, where $\mathcal{P}$ is any property preserved by homomorphisms, e.g., the property of being simple or $\{0\}$, singular, finitely generated, cyclic, Noetherian, and Artinian.

Proof. Part (a) follows from III.2(b), and part (b) follows from III.2(c).

Another way of getting sinks in the case $(\mathcal{E}, \text{End})$ is given by

Theorem III.4. Let $F \leq G$ be a fully invariant submodule. Then $\{i \in I | C_i \subseteq F\}$ is a sink with respect to the matched pair $(\mathcal{E}, \text{End})$.

This result, whose straightforward proof is omitted, shows, for example, that there are sinks associated with $\text{Rad } G$ and $\text{Soc } G$ and with the singular submodule $Z(G)$ of G .

We now turn to simplicity of $M(\mathcal{E}, \mathcal{H})$. From the above we know if there are two or more nonzero classes in $I/\sim$, then M cannot be simple. However, the existence of just one nonzero class does not guarantee simplicity, as the example with $G = Q$ and $R = \mathbb{Z}$ discussed after Corollary II.2 shows. We start with a decomposition result.

Lemma III.5. Suppose $I = [0] \cup (\bigcup_{k=1}^n [i_k])$ where each $[i_k]$ is minimal. Then $M = \sum_{k=1}^n \oplus M_k$ where $M_k := A(I \setminus [i_k])$ is an ideal.

Proof. Note that $I \setminus [i_k]$ is a sink for each k , $M = \sum_{k=1}^n M_k$, and $M_k \cap (\sum_{k \neq l} M_l) = \{0\}$.

Recall that $\mathcal{S}$ is the class of simple R -modules. We now consider the case $(\mathcal{E}_G(\mathcal{S}), \text{Hom})$.

Theorem III.6. *If G is Noetherian with $\text{Soc } G \neq \{0\}$, then $M(\mathcal{E}_G(\mathcal{S}), \text{Hom})$ is a direct sum of a finite number of ideals, each of which is invariantly simple as a near-ring.*

Proof. Since G is Noetherian, there are only finitely many isomorphism types of simple R -modules represented in $\mathcal{E}_G(\mathcal{S})$. Each isomorphism class gives rise to a class in $I/\sim$ which is minimal with respect to $\preceq$. Since the union of these classes together with $[0]$ is I , we can apply the preceding lemma to obtain a decomposition of M into ideals. To complete the proof, it suffices to show if $I = [0] \cup [i]$, where C_j is simple for $j \in [i]$, then $M(\mathcal{E}_G(\mathcal{S}), \text{Hom})$ is invariantly simple. Since G is Noetherian, so is $F := \text{Soc } G$, and therefore F is the direct sum of some of the C_j . For ease of notation, we let $F = \sum_{i=1}^n \oplus C_i$. Denote by $\eta^i : F \rightarrow C_i$, $i = 1, 2, \dots, n$, the projection onto C_i . Thus $\sum_{i=1}^n \eta^i = 1_F$. Also, η^i represents an element of M by defining $(\eta^i)_l := (\eta^i|_{C_l})$, $l \in I$, and in this context, $\sum_{i=1}^n \eta^i = e$, the identity in M . Now let $s \in M$, $s \neq 0$, with, say, $s_j \neq 0$. From Corollary I.11, $s_j^{-1} \in H$. Therefore, if $t \in M$ is such that $t_{\tau(s_j)} = s_j^{-1}$, then $e^j = ts e^j \in \langle s \rangle$, the invariant subgroup of M generated by s . But from e^j we can easily construct e^i , $i = 1, 2, \dots, n$, by left and right multiplications, again using Corollary I.11. Now $\eta^i = e^i \eta^i$ for $i = 1, 2, \dots, n$, because, since C_i is simple, $\tau((\eta^i)_l)$ is either i or 0 , and therefore $e = \sum_{i=1}^n \eta^i \in \langle s \rangle$. This completes the proof.

Since every simple submodule of an R -module G is cyclic, we have

Corollary III.7. *If G is a semisimple, Noetherian module, then $M(\mathcal{E}_G(R), \text{Hom})$ is a direct sum of a finite number of ideals, each of which is invariantly simple as a near-ring.*

A special case is the following

Corollary III.8. *If V is a finite-dimensional vector space over a division ring, then $M(\mathcal{E}_V(D), \text{Hom})$ is invariantly simple.*

The above result is not true for an infinite-dimensional vector space, V , over a division ring D . In fact, if we define, for $f \in M$, $\mathcal{R}(f) := \text{Span}\{f(C_i) | C_i \in \mathcal{E}\}$ and let $K := \{f \in M | \dim \mathcal{R}(f) < \infty\}$, then K is an invariant subgroup of M , $\{0\} \subsetneq K \subsetneq M$. However, from the next result, which has wide applicability, we will find that in this case $M(\mathcal{E}_V(D), \text{Hom})$ is simple.

Theorem III.9. *Let $(\mathcal{E}, \mathcal{H})$ be a matched pair on G with the following properties:*

- (i) *There is only one nonzero class $[i]$.*
- (ii) *If $h \in H$ is nonzero, then $h^{-1} \in H$.*
- (iii) *There is some $F \leq G$ such that $|\{i \in I | C_i \subseteq F\}| = |I|$.*
- (iv) *There exists a $C \in \mathcal{E}$ with $C \not\subseteq F$.*

Then $M(\mathcal{E}, \mathcal{H})$ is a simple near-ring.

Proof. For convenience we let $C_1 := C$ and $J := \{i \in I \mid C_i \subseteq F\}$. From (iii) there is a bijection between I and J , so without loss of generality we let $\Phi: I \rightarrow J$ be a bijection with $\Phi(0) = 0$. Suppose s is a nonzero element of M . We show that the ideal generated by s , which we denote by T , is M . We first show that $\varepsilon^1 \in T$. Indeed, if $s_i \neq 0$, then we know there exist $u, v \in M$ with $\tau(v_1) = i$ and $v_j = 0$ for $j \neq 1$ and $u_{\tau(s_i)} = v_1^{-1}s_i^{-1}$, $u_j = 0$ otherwise. Then $usv = \varepsilon^1$. Next, let $t \in M$ be such that $\tau(t_i) = 1$ if $i \neq 0$. Hence $\varepsilon^1 t = t$. Now choose $x \in M$ so that $\tau(t_i + x_i) = \Phi(i)$. We claim that $\tau(x_i) \notin J$ for all $i \neq 0$. To establish this for a particular i , let $g \in C_i$ be such that $t_i(g) \notin F$, which is possible because $C_1 \subsetneq F$. Then $(t_i + x_i)(g) = t_i(g) + x_i(g) \in C_{\Phi(i)} \subseteq F$. This implies $x_i(g) \notin F$, which proves our assertion. Finally, choose $y \in M$ with $y_j = (t_{\Phi^{-1}(j)} + x_{\Phi^{-1}(j)})^{-1}$ if $0 \neq j \in J$ and $y_j = 0$ if $j \notin J$ or $j = 0$. Then $yx = 0$ and $y(t + x) = e$, so $e = y(t + x) - yx \in T$, i.e. $T = M$.

We remark that Theorem I.7 follows from Theorem III.9. Further we have

Corollary III.10. *Let V be a vector space over a division ring D . Then $M(\mathcal{E}_V(D), \text{Hom})$ is a simple near-ring.*

Proof. The finite-dimensional case is that of Corollary III.8. For the infinite-dimensional case, let $\mathcal{B} = \{b_k\}$, $k \in \mathcal{K}$, be a basis for V . Then the cardinality of the set of cyclic submodules in $V' := \text{Span}(\mathcal{B} \setminus \{b_{k_0}\})$ is the same as that of the set of all cyclic submodules in V . So, by applying Theorem I.12 and choosing $F := V'$, $C := b_{k_0}D$, we see that the hypotheses of Theorem III.9 are satisfied. Thus $M(\mathcal{E}_V(D), \text{Hom})$ is simple.

As a further application of Theorem III.9 we let K be any infinite field and let $R := K[x]$ and $G := K[x]$. With $\mathcal{E} := \mathcal{E}_G(R)$ and $\mathcal{H} := \text{Hom}$, it is clear that (i), (ii), and (iii) of Theorem III.9 are satisfied if we take $F := xK[x]$. Note that for each $\alpha \in K$, $(x^2 + \alpha x)K[x] \subseteq F$, but $K[x] \not\subseteq F$.

We wish to characterize when $M = M(\mathcal{E}_G(R), \text{Hom})$ is a simple near-ring. As we have noted above, if M is simple, then there is only one nonzero class with respect to $\sim$, i.e., $|I/\sim| = 2$. So, we focus now on $|I/\sim| = 2$. In certain situations this condition is also sufficient.

Theorem III.11. *Let $\mathcal{E} = \mathcal{E}_G(R)$, $\mathcal{H} = \text{Hom}$, and suppose G has finite length. Then the following are equivalent:*

- (1) M is simple.
- (2) M is invariantly simple.
- (3) $|I/\sim| = 2$.

In this case, G is semisimple.

Proof. We know (2) $\Rightarrow$ (1) and (1) $\Rightarrow$ (3), so it remains to show (3) $\Rightarrow$ (2). To this end let $C_j \in \mathcal{E}$. Since $\mathcal{E} = \mathcal{E}_G(R)$, C_j is a cyclic R -module. We show each C_j is simple. Assume the contrary. Then there is a cyclic submodule C_i of C_j , $C_i \subsetneq C_j$. Now, $C_i \in \mathcal{E}$ and $|I/\sim| = 2$ implies there exists $h \in H$, $h: C_j \rightarrow C_i$. Moreover, since G is of finite length, G is Noetherian, so $h^{-1} \in H$. But then, considering $h|C_i$, we get $h(C_i) = C_k \subsetneq C_i$, otherwise $\text{Ker}(h) \neq \{0\}$. In this manner we get an infinite descending chain in G , contrary to G being of finite length. Since each cyclic is simple, $\text{Soc } G \neq \{0\}$, so now as in the proof of Theorem III.6, we get that M is invariantly simple.

To show that G is semisimple, let $a \in G$. If $G = aR$, then we are finished. If not, there exists $b \in G \setminus aR$. Then $aR \cap bR = \{0\}$ since they are both simple.

If $G = aR \oplus bR$, then again we are finished. If not, we find $c \in G \setminus aR \oplus bR$. This process must end since G is of finite length.

Corollary III.12. *If R is an Artinian ring and G is finitely generated, then the following are equivalent:*

- (1) M is simple.
- (2) M is invariantly simple.
- (3) $|I/\sim| = 2$.

Further, if G is a faithful R -module, then R is a semisimple ring.

Proof. Since R is Artinian, R is Noetherian, and G is finitely generated, G has finite length. Thus the first sentence follows from Theorem III.11. Suppose now G is faithful. For each $g \in G \setminus \{0\}$, gR is simple, so $r_R(g) = \{r \in R \mid gr = 0\}$ is a maximal right ideal of R . Therefore, $J(R) \subseteq \bigcap_{g \in R} r_R(g) = A(G) = \{0\}$, hence the result.

We remark that Theorems I.6 and I.7 show that the above corollary is not true if we take R to be Noetherian. We are thus led to the following problem which remains open.

Problem. Let G be Noetherian with matched pair $(\mathcal{E}_G(R), \text{Hom})$ and $|I/\sim| = 2$. Is $M(\mathcal{E}_G(R), \text{Hom})$ a simple near-ring?

We conclude this section and the paper with one further special case. We start here with a few remarks which provide a summary of the above, indicating the present status of the problem. First, we note that all of the nonzero $C_i \in \mathcal{E}$ are isomorphic, and if $h: C_i \rightarrow C_j$ is any isomorphism, then $h \in H$. Moreover, if $h \in H$, $h \neq 0$, then $h^{-1} \in H$.

Suppose any $C \in \mathcal{E}$ is minimal; then it is simple, and hence every $C \in \mathcal{E}$ is simple. So this is the semisimple case, and since G is Noetherian, G is the direct sum of finitely many simple submodules. This case was handled in Theorem III.6. So, henceforth, we assume every nonzero $C \in \mathcal{E}$ contains an infinite descending chain of elements of $\mathcal{E}$. If any one (and therefore all) of them contains $|I|$ elements of $\mathcal{E}$, then we can apply Theorem III.9. We therefore assume that no element of $\mathcal{E}$ contains $|I|$ elements of $\mathcal{E}$. This implies that G is not cyclic, so every cyclic is proper.

Since G is Noetherian, G is the sum (albeit not necessarily direct) of finitely many elements of $\mathcal{E}$. For convenience we take $G = \sum_{i=1}^n C_i$. Let $J_i := \{j \in I \mid C_j \subseteq C_i\}$, $i = 1, 2, \dots, n$, and let C_{j_i} be chosen such that $j_i \notin J_i$, i.e., $C_{j_i} \not\subseteq C_i$.

Let T be any nonzero ideal of M . Then, for each $l \in I$, $\varepsilon^l \in T$. In fact, let $0 \neq s \in T$. For some $i \in I$, $s_i \neq 0$. Furthermore, there exists $h \in H$, $h: C_l \rightarrow C_i$. We let $u \in M$ be defined by $u_l = h$ and $u_j = 0$, $j \neq l$, and let $v \in M$ be defined by $v_{\tau(s_i)} = h^{-1}s_i^{-1}$ and 0 otherwise. Then $vsue^l \in T$ and $vsue^l = \varepsilon^l$. Note that here we have used only the fact that T is an invariant subgroup.

Now, for $i = 1, 2, \dots, n$ define $\delta^i \in M$ by $(\delta^i)_j = e_j$ if $j \in J_i$ and 0 otherwise. We show $\delta^i \in T$. Indeed, let $s \in M$ with $\tau(s_j) = j_i$ if $j \in J_i$ and 0 otherwise. Then $\varepsilon^{j_i}s = s \in T$ since $\varepsilon^{j_i} \in T$. Now choose $x \in M$ such that $s_j + x_j = e_j$ if $j \in J_i$ and 0 otherwise. If $c \in C_j$ is such that $s_j(c) \notin C_i$ (which is possible because $s_j(C_j) = C_{j_i} \not\subseteq C_i$), then $s_j(c) + x_j(c) = c$. Therefore, $x_j(c) \notin C_i$, i.e., $\tau(x_j) \notin J_i$. From this we get $\delta^i(s + x) - \delta^i x = \delta^i \in T$.

We now make an additional assumption, namely, if $C_i, C_j \in \mathcal{C}$ with $C_i = g_i R$ and $C_j = g_j R$, then there exists $h: C_i \rightarrow C_j$ with $h(g_i) = g_j$. From this, we show $e \in T$. Let $K := \{g_i | i \in I\}$ be a set of generators for $\mathcal{C}$, i.e., $C_i = g_i R$. Let $j \in I$, $j \neq 0$, and suppose $g_j = f_{j1} + \cdots + f_{jn}$, where $f_{ji} \in C_i$. Each f_{ji} is a generator for a C_{ji} , although we do not necessarily have $f_{ji} \in K$. However, we do have $ji \in J_i$. Let $h_{ji} \in H$ be that isomorphism of C_j onto C_{ji} which takes g_j to f_{ji} , and let $(\gamma^i)_j = h_{ji}$, for each $j \in I$, $j \neq 0$. Then $\delta^i \gamma^i = \gamma^i \in T$. We claim that $\sum_{i=1}^n \gamma^i = e$. Indeed, $(\sum_{i=1}^n \gamma^i)_j(g_j) = (h_{j1} + \cdots + h_{jn})(g_j) = f_{j1} + \cdots + f_{jn} = g_j$. Hence, $e \in T$, and consequently M is simple. We have established

Theorem III.13. *Let G be a noncyclic Noetherian R -module with matched pair $(\mathcal{E}_G(R), \text{Hom})$ satisfying $|I| \sim |J| = 2$. If for each $C_i, C_j \in \mathcal{C}$, $C_i = g_i R$, $C_j = g_j R$, there exists $h: C_i \rightarrow C_j$ with $h(g_i) = g_j$, then $M(\mathcal{E}_G(R), \text{Hom})$ is a simple near-ring.*

ACKNOWLEDGMENT

This paper was written while the second author was a visiting professor at Texas A & M University in 1991. He wishes to express his gratitude for the financial assistance and hospitality provided by Texas A & M.

REFERENCES

1. F. W. Anderson and K. R. Fuller, *Rings and categories of modules*, Graduate Texts in Math., vol. 13, Springer-Verlag, New York, Heidelberg, and Berlin, 1973.
2. P. Fuchs, C. J. Maxson, and G. Pilz, *On rings for which homogeneous maps are linear*, Proc. Amer. Math. Soc. **112** (1991), 1–7.
3. C. J. Maxson, *Piecewise endomorphisms of PID-modules*, Results in Math. **18** (1990), 125–132.
4. C. J. Maxson and A. P. J. van der Walt, *Homogeneous maps as piecewise endomorphisms*, Comm. Algebra **20** (1992), 2755–2776.
5. J. D. P. Meldrum, *Near-rings and their links with groups*, Pitman Res. Notes Math. Ser., vol. 134, Longman Sci. Tech., Harlow, 1985.
6. G. F. Pilz, *Near-rings*, 2nd ed., North-Holland, Amsterdam, 1983.

DEPARTMENT OF MATHEMATICS, TEXAS A & M UNIVERSITY, COLLEGE STATION, TEXAS 77843
E-mail address: cjmaxson@math.tamu.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF STELLENBOSCH, 7600 STELLENBOSCH, SOUTH AFRICA
E-mail address: apjw@sunvax.sun.ac.za