AMENABLE ACTIONS AND WEAK CONTAINMENT OF CERTAIN REPRESENTATIONS OF DISCRETE GROUPS

M. GABRIELLA KUHN

(Communicated by J. Marshall Ash)

ABSTRACT. We consider a countable discrete group Γ acting ergodically on a standard Borel space S with quasi-invariant measure μ . Let π be a unitary representation of Γ on $L^2(S, d\mu, \mathcal{H})$ “nicely” related with S . We prove that if Γ acts amenably on S then π is weakly contained in the regular representation.

Let Γ be a countable discrete group acting (on the right) on a standard Borel space S . Suppose that the action is ergodic with μ as a quasi-invariant probability measure on S . Let $P(s, g)$ be the Radon-Nikodym cocycle of the action, that is, $P(s, g) = \frac{d\mu(sg)}{d\mu(s)}$. Construct a representation of Γ as follows:

- (1) choose any Hilbert space $\mathcal{H}$,
- (2) choose any (Borel) cocycle $A(s, g) : S \times G \rightarrow \mathcal{U}(\mathcal{H})$ where $\mathcal{U}(\mathcal{H})$ denotes the group of unitary operators on $\mathcal{H}$,
- (3) construct $\mathcal{H}_\pi = L^2(S, d\mu, \mathcal{H})$ with the usual inner product given by $\langle F, G \rangle = \int_S \langle F(s), G(s) \rangle_{\mathcal{H}} d\mu(s)$,

and finally define

$$(\pi(g)F)(s) = P(s, g)^{\frac{1}{2}} A(s, g) F(sg).$$

Observe now that although the group *is not* amenable it turns out that π is weakly contained in the regular representation for many natural constructions. In particular, this is true when

- (a) Γ is a free group and S is its Martin boundary (see [K-S, Appendix]),
- (b) Γ is a hyperbolic group and S is its boundary (see [A] and use the techniques of [K-S]), and
- (c) Γ is a lattice in a semisimple Lie group G and $S = G/B$ is the maximal Furstenberg boundary of G (see [Q-S, §4]).

Recall that a group is amenable if and only if every unitary representation of Γ is weakly contained in ${}^\infty\pi_{\text{reg}}$ (where π_{reg} denotes the regular representation) [H] and [Z3, Proposition 7.3.6].

Received by the editors February 10, 1993.

1991 *Mathematics Subject Classification.* Primary 43A07, 22D10.

Key words and phrases. Amenable actions, weak containment of representations.

©1994 American Mathematical Society
0002-9939/94 \$1.00 + \$.25 per page

Zimmer introduced a class of Borel spaces which make a nonamenable group G “look like” an amenable one when acting on them. These spaces are called *amenable G -spaces* (G is said to act amenably on them).

To see that cases (a) and (c) deal with amenable actions of Γ on S recall the following

Proposition [Z3]. *Let G be a locally compact group, H a closed subgroup, and Γ a lattice in G . Then Γ acts amenably on G/H if and only if H is amenable.*

Case (c) follows since minimal parabolic subgroups of semisimple Lie groups are amenable. For case (a) the reader should think of the free group acting on a homogeneous tree as a discrete cocompact subgroup of the full automorphism group G of the tree. The Martin boundary of Γ is the set of all “ends” of the tree, and it was proved by Nebbia [N] that the stabilizer (in G) of an end is amenable. Case (b) was proved by Adams in [A].

In this paper we shall prove the following

Theorem. *Suppose that a countable discrete group Γ acts ergodically on (S, μ) with quasi-invariant measure μ . Suppose that the action is amenable. Then every representation π constructed as above is weakly contained in π_{reg} .*

We also believe that the converse is true; namely, we believe that if for every Hilbert space $\mathcal{H}$ the above-constructed representation $(L^2(S, d\mu, \mathcal{H}), \pi)$ is weakly contained in π_{reg} then the action is amenable.

Proof of the result. Denote by π_{reg} the right regular representation of Γ . Construct $\pi_{\text{reg}} \otimes \pi$ acting on $\ell^2(\Gamma) \otimes \mathcal{H}_\pi$. Recall that $U : \ell^2(\Gamma) \otimes \mathcal{H}_\pi \rightarrow \ell^2(\Gamma, \mathcal{H}_\pi)$ given by

$$(Uf)(x) = \pi(x)f(x)$$

intertwines $\pi_{\text{reg}} \otimes \pi$ to $\pi_{\text{reg}} \otimes I$, where I denotes the trivial representation. Thus $\pi_{\text{reg}} \otimes \pi$ is equivalent to a direct sum of copies of π_{reg} , and it is sufficient to prove that π is weakly contained in $\pi_{\text{reg}} \otimes \pi$.

Fix any finite subset E of Γ . Let $F \in L^2(S, d\mu, \mathcal{H})$. Inside $\ell^2(\Gamma) \otimes L^2(S, d\mu, \mathcal{H})$ we shall construct a sequence of functions $(f_n)_{n=0}^\infty$ such that $\langle \pi_{\text{reg}} \otimes \pi(x)f_n, f_n \rangle$ approaches $\langle \pi(x)F, F \rangle$ for x in E .

Identify $\ell^2(\Gamma) \otimes L^2(S, d\mu, \mathcal{H})$ with $L^2(S, d\mu, \ell^2(\Gamma) \otimes \mathcal{H})$. Set $f_n(s) = \phi_n(s) \otimes F(s)$ where the ϕ_n are elements of

$$\ell^2(\Gamma) \otimes L^2(S, d\mu) = L^2(S, d\mu, \ell^2(\Gamma)).$$

Compute

$$\begin{aligned} & \langle \pi_{\text{reg}} \otimes \pi(g)f_n, f_n \rangle \\ &= \int_S P(s, g)^{\frac{1}{2}} \langle \pi_{\text{reg}}(g)\phi_n(sg), \phi_n(s) \rangle_{\ell^2(\Gamma)} \langle A(s, g)F(sg), F(s) \rangle_{\mathcal{H}} d\mu(s) \end{aligned}$$

and

$$\langle \pi(g)F, F \rangle = \int_S P(s, g)^{\frac{1}{2}} \langle A(s, g)F(sg), F(s) \rangle_{\mathcal{H}} d\mu(s).$$

In order to prove our result we need a sequence of functions $\phi_n(s, x) \in L^2(S, d\mu, \ell^2(\Gamma))$ such that $\sum_{x \in \Gamma} \phi_n(sg, xg)\phi_n(s, x)$ is weakly converging, in $L^2(S, d\mu)$, to the function identically one on S .

For amenable groups the existence of such a sequence is related to existence of an invariant mean in $(L^\infty(G))^*$.

An analogue of the invariant mean property was proved for discrete groups by Zimmer in [Z1, Z2] in the case of an amenable action. The reader may take the following as a definition of an amenable action:

Proposition 1 (Zimmer). *Suppose that Γ is a countable discrete group acting ergodically on (S, μ) . Then S is an amenable Γ -space if and only if there is a norm one linear map $\sigma : L^\infty(S \times \Gamma) \rightarrow L^\infty(S)$ such that:*

- (1) $\sigma(\mathbf{1}) = \mathbf{1}$, $f \geq 0$ implies $\sigma(f) \geq 0$;
- (2) if $A \subseteq S$ is measurable, $\sigma(f \cdot \chi_{p^{-1}A}) = \sigma(f)\chi_A$ where $p : S \times \Gamma \rightarrow S$ is the projection; and
- (3) $\sigma(f \cdot g) = \sigma(f) \cdot g$, where $f \cdot g(s, x) = f(sg, xg)$ and $\phi \cdot g(s) = \phi(sg)$.

In [Z1] it is shown how to construct σ once we know that the action of Γ on S is amenable. From these we may deduce other properties of σ which will be needed. For the reader's convenience we briefly recall the construction. Let $\tilde{U}_g : L^2(S \times \Gamma) \rightarrow L^2(S \times \Gamma)$ be defined by

$$(\tilde{U}_g f)(s, x) = f(sg, xg)P(s, g)^{\frac{1}{2}} ; .$$

For $f \in L^\infty(S)$ define a multiplication operator M_f on $L^2(S \times \Gamma)$ by letting

$$(M_f h)(s, x) = f(s)h(s, x) .$$

Let R denote the von Neumann algebra generated by $\{\tilde{U}_g, M_f\}$ in $\mathcal{B}(L^2(S \times \Gamma))$. It was proved in [Z2] that if Γ acts amenably on S then there is a norm one projection and hence a conditional expectation $P : \mathcal{B}(L^2(S \times \Gamma)) \rightarrow R$.

Consider now the following decomposition for $L^2(S \times \Gamma)$. For $f \in L^2(S \times \Gamma)$ let $f_s(x) = f(sx, x)P(s, x)^{\frac{1}{2}}$. Then $f \rightarrow \int_S^\oplus f_s d\mu$ is a unitary isomorphism of $L^2(S \times \Gamma) \simeq \int_S^\oplus \ell^2(\Gamma) d\mu$. Further every element of R is decomposable with respect to this direct integral decomposition. (Namely, $\tilde{U}_g$ corresponds to $\int_S^\oplus \pi_{\text{reg}}(g)$ and M_f to $\int_S^\oplus f^s$ where $(f^s h_s)(x) = f(sx)h_s(x)$.) For every $f \in L^\infty(S \times \Gamma)$ we have the multiplication operator $M_f \in \mathcal{B}(L^2(S \times \Gamma))$. Now $P(M_f)$ is decomposable; hence, we can write $P(M_f) = \int_S^\oplus T_s^f d\mu$. Our map σ is now given by

$$(\sigma f)(s) = \langle T_s^f(\delta_e), \delta_e \rangle_{\ell^2(\Gamma)} ,$$

where δ_e denotes the characteristic function of the identity e .

In particular, if F is an element of $L^\infty(S)$ and we let $F(s, g) = F(s)$ for every $g \in \Gamma$ then

$$(i) \quad (\sigma F)(s) = F(s) ;$$

furthermore, for every $h \in L^\infty(S \times \Gamma)$

$$(ii) \quad (\sigma h F)(s) = (\sigma h)(s)F(s) ,$$

since $P(M_h M_F) = P(M_h)M_F$ because F is an element of R .

In order to use our function σ we need the Radon-Nikodym derivative $P(s, g)^{\frac{1}{2}}$ to be a bounded function of s for any fixed g . This, in general, may or may not be true. However, we shall prove that it is always possible to find a measure ν equivalent to μ , for which this is true.

Lemma 1. *Let Γ be a discrete group. There exists a length function $x \rightarrow \|x\|$ such that the cardinality of the sets $\{\|x\| \leq N\}$ is less than or equal to 3^N . Given such a length function choose $K > 3$ and set*

$$\nu(E) = \sum_{x \in \Gamma} K^{-\|x\|} \mu_x(E) \quad \text{where } \mu_x(E) = \mu(Ex)$$

for every measurable E . Then ν is equivalent to μ and the Radon-Nikodym derivative $\frac{d\nu(sg)}{d\nu(s)}$ is less than or equal to $K^{\|g\|}$.

Proof. Choose any ordering $\{x_n\}_{n=0}^\infty$ for the elements of Γ satisfying $x_0 = e$. Set

$$\|x\| = \inf \left\{ \sum_{j=1}^J n_j : x = x_{n_1}^\pm x_{n_2}^\pm \cdots x_{n_J}^\pm \right\}.$$

It is immediate that $\|\cdot\|$ is a length function. Set $\Phi(N) = \#\{x : \|x\| = N\}$. We observe that the number of strictly positive solutions $(n_1, n_2, \dots, n_k)$ of the equation

$$n_1 + n_2 + \cdots + n_k = N$$

is $\binom{N-1}{k-1}$. Hence

$$\Phi(N) \leq \sum_{k=1}^N \binom{N-1}{k-1} 2^k = 2 \cdot 3^{N-1}.$$

Finally

$$\#\{x : \|x\| \leq N\} \leq 1 + 2 + 2 \cdot 3 + \cdots + 2 \cdot 3^{N-1} = 3^N.$$

It is obvious from the definition of ν and from the quasi-invariance of μ that ν and μ are equivalent measures. Let E be a measurable set. Compute

$$\begin{aligned} \nu(Eg) &= \sum_{x \in \Gamma} K^{-\|x\|} \mu(Egx) = \sum_{x \in \Gamma} K^{-\|x\| - \|g\| + \|g\|} \mu(Egx) \\ &\leq K^{\|g\|} \sum_{x \in \Gamma} K^{-\|gx\|} \mu(Egx) = K^{\|g\|} \nu(E). \end{aligned}$$

This completes the proof. $\square$

From this point on we shall assume that the Radon-Nikodym derivative $P(s, g)$ is a bounded function of s for any given g .

Proof of the theorem. Let

$$P = \left\{ \phi \in L^1(S \times \Gamma) \mid \phi(s, x) \geq 0 \text{ and } \int_S \sum_{x \in \Gamma} \phi(s, x) d\mu(s) = 1 \right\}.$$

First of all observe that P is weak- $*$ dense in the unit ball of $(L^\infty(S \times \Gamma))^*$. Construct a linear functional σ^* on $L^\infty(S \times \Gamma)$ by letting

$$\sigma^*(f) = \int_S (\sigma f)(s) d\mu(s).$$

Recall that $(h \cdot g)(s, x) = h(sg, xg)$ and $(\psi \cdot)g(s) = \psi(sg)$.

Choose h in $L^\infty(S \times \Gamma)$, ϵ positive, and g in a finite set E of Γ . First of all we shall show that it is possible to find $\phi \in L^1(S \times \Gamma)$ such that

$$(1.1) \quad \left| \int_S \left(\sum_{x \in \Gamma} (\phi(sg, xg) - \phi(s, x)) h(s, x) \right) d\mu(s) \right| < \epsilon \quad \text{for every } g \text{ in } E.$$

In fact, by property (c) of Proposition 1, we have

$$\begin{aligned} \int_S (\sigma h)(s) d\mu(s) &= \int_S (\sigma h \cdot g^{-1})(s) P(s, g^{-1}) d\mu(s) \quad (\text{because of (ii)}) \\ &= \int_S (\sigma h \cdot g^{-1} P(s, g^{-1}))(s) d\mu(s). \end{aligned}$$

Hence it is possible to choose $\phi \in L^1(S \times \Gamma)$ so that

$$(1.2) \quad \left| \int (\sigma h)(s) - \left(\sum_{x \in \Gamma} \phi(s, x) h(s, x) \right) d\mu(s) \right| < \epsilon$$

and

$$(1.3) \quad \left| \int_S (\sigma h \cdot g^{-1} P(s, g^{-1}))(s) - \left(\sum_{x \in \Gamma} \phi(s, x) h(sg^{-1}, xg^{-1}) P(s, g^{-1}) \right) d\mu(s) \right| < \epsilon.$$

But

$$\begin{aligned} &\int_S \left(\sum_{x \in \Gamma} \phi(s, x) h(sg^{-1}, xg^{-1}) P(s, g^{-1}) \right) d\mu(s) \\ &= \int_S P(s, g^{-1}) \sum_{x \in \Gamma} \phi(s, xg) h(sg^{-1}, x) d\mu(s) \end{aligned}$$

and letting $sg^{-1} = s'$

$$= \int_S \sum_{x \in \Gamma} \phi(sg, xg) h(s, x) d\mu(s).$$

Subtracting (1.3) from (1.2) we get (1.1)

Moreover, since $\sigma(1) = 1$, we may also require that for any given $h' \in L^\infty(S)$

$$(1.4) \quad \left| \int_S h'(s) \left(1 - \sum_{x \in \Gamma} \phi(s, x) \right) d\mu(s) \right| < \epsilon.$$

We shall use now Namioka's argument to pass from the weak to the strong closure. Let $F = L^1(S) \times \prod_{g \in \Gamma} L^1(S \times \Gamma) = F_0 \times \prod_{g \in \Gamma} F_g$ where $L^1(S)$ and each copy of $L^1(S \times \Gamma)$ has the norm topology. The weak topology on F is the product topology for the weak topology on $L^1(S)$ and on $L^1(S \times \Gamma)$. Define a linear map $T : P \rightarrow F$ by letting

$$(T(\phi))_0 = \sum_{x \in \Gamma} \phi(s, x) \quad \text{and} \quad (T(\phi))_g = \phi(sg, xg) - \phi(s, x).$$

Then (1.1) and (1.4) say that the point $(1, 0, \dots, 0, \dots)$ is in the weak closure of $T(P)$, and since $T(P)$ is convex, it follows that the same point is in the strong closure of $T(P)$. Hence we can find a net of elements $\psi_j \in P$ such that

$$\sum_{x \in \Gamma} \psi_j(s, x) \rightarrow 1 \quad \text{and} \quad \psi_j(sg, xg) - \psi_j(s, x) \rightarrow 0$$

strongly, respectively, in $L^1(S)$ and in $L^1(S \times \Gamma)$.

Finally let $\phi_j(s, x) = \sqrt{\psi_j(s, x)}$. By standard arguments

$$\|\phi_j \cdot g - \phi_j\|_{L^2(S \times \Gamma)} \leq \|\psi_j \cdot - \psi_j\|_{L^1(S \times \Gamma)}^{\frac{1}{2}}.$$

Choose any F in $L^2(S, d\mu, \mathscr{H})$. Set $f_n(s, x) = \phi_n(s, x) \otimes F(s)$. Let

$$G(s, g) = P(s, g)^{\frac{1}{2}} \langle A(s, g)F(sg), F(s) \rangle_{\mathscr{H}}.$$

Since the set of functions $F : S \rightarrow \mathscr{H}$ such that $\langle (\pi(g)F)(s), F(s) \rangle$ is a bounded function of s is dense in $L^2(S, d\mu, \mathscr{H})$, we may assume that $G(s, g)$ is a bounded function of s for any given $g \in \Gamma$. We may also assume that $G(s, g) \geq 0$. Observe that $\|\sqrt{G(s, g)}(\phi_n \cdot g - \phi_n)\|_{L^2(S \times \Gamma)} \rightarrow 0$. We have

$$\begin{aligned} & \|\sqrt{G(s, g)}(\phi_n \cdot g - \phi_n)\|_{L^2(S \times \Gamma)} \\ &= \int_S G(s, g) \sum_{x \in \Gamma} \psi_n(sg, xg) d\mu(s) + \int_S G(s, g) \sum_{x \in \Gamma} \psi_n(s, x) d\mu(s) \\ & \quad - 2 \int_S G(s, g) \sum_{x \in \Gamma} (\phi_n(sg, xg) \phi_n(s, x)) d\mu(s). \end{aligned}$$

But $\sum_{x \in \Gamma} \psi_n(s, x)$ converges to 1 in $L^1(S)$, and so does $\sum_{x \in \Gamma} \psi_n(sg, xg)$ since $\|\psi_n \cdot g - \psi_n\|_{L^1(S \times \Gamma)} \rightarrow 0$. Hence,

$$\int_S G(s, g) \sum_{x \in \Gamma} (\phi_n(sg, xg) \phi_n(s, x)) d\mu(s) \rightarrow \int_S G(s, g) d\mu(s). \quad \square$$

ACKNOWLEDGMENT

We are happy to thank Tim Steger for suggesting the proof of Lemma 1. A special thanks to Robert Zimmer for helpful conversations and remarks and for providing us with [Z1] and [Z2].

REFERENCES

- [A] S. Adams, *Boundary amenability for hyperbolic groups and an application to smooth dynamic of simple groups*, preprint.
- [H] A. Hulanicki, *Means and Folner conditions on locally compact groups*, *Studia Math.* **27** (1966), 87–104.
- [K-S] M. G. Kuhn and T. Steger, *More irreducible boundary representations of free groups*, preprint.
- [N] C. Nebbia, *Amenability and Kunze-Stein property for groups acting on a tree*, *Pacific J. Math.* **135** (1988), 371–380.
- [Q-S] J. C. Quigg and J. Spielberg, *Regularity and hyporegularity in C^* -dynamical systems*, *Houston J. Math.* **18** (1992), 139–152.

- [Z1] R. J. Zimmer, *On the von Neumann algebra of an ergodic group action*, Proc. Amer. Math. Soc. **66** (1977), 289–293.
- [Z2] ———, *Hyperfinite factors and amenable ergodic actions*, Invent. Math. **41** (1977), 23–31.
- [Z3] ———, *Ergodic theory and semisimple groups*, Birkhäuser, Boston, MA, 1984.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DEGLI STUDI DI MILANO, VIA C. SADINI, 50 20133
MILANO, ITALY

E-mail address: `kuhn@vmimat.mat.unimi.it`