SOME ESTIMATES OF THE KOBAYASHI METRIC IN THE NORMAL DIRECTION

SIQI FU

(Communicated by Eric Bedford)

ABSTRACT. In this paper, we study the behavior of the Kobayashi metric in the normal direction near a Levi-pseudoconvex boundary point of a smoothly bounded domain without assuming global pseudoconvexity. As a corollary, we obtain a characterization of pseudoconvexity by the rate of the growth of the Kobayashi metric in the normal direction.

I. INTRODUCTION AND THEOREMS

Let $\Omega \subset \subset \mathbb{C}^n$ be a bounded domain with smooth boundary near $z_0 \in \partial\Omega$. Let U be a neighborhood of z_0 . Let $r(z)$ be a local defining function of Ω on U , i.e.,

$$\Omega \cap U = \{z \in U \mid r(z) < 0\}$$

and $r(z) \in \mathcal{C}^\infty(U)$, $\nabla r(z)|_{\partial\Omega \cap U} \equiv (\partial r(z)/\partial z_1, \partial r(z)/\partial z_2, \dots, \partial r(z)/\partial z_n) \neq 0$.

Let Δ be the unit disc and let $\Delta_\gamma \equiv \{\gamma\zeta; \zeta \in \Delta\}$. The Kobayashi metric of Ω is defined by

$$F_\Omega(z, X) \equiv \inf \{1/\lambda \mid \exists f: \Delta \rightarrow \Omega \text{ is holomorphic and } f(0) = z, f'(0) = \lambda X, \lambda > 0\}.$$

For $z = (z_1, z_2, \dots, z_n) \in \mathbb{C}^n$, set $z' = (z_1, z_2, \dots, z_{n-1})$. Let $d(z) = \text{dist}(z, \partial\Omega)$ and let $\pi(z)$ be the projection to the boundary for z near z_0 such that $d(z) = |z - \pi(z)|$. Let $N_{\pi(z)}$ be the inward normal direction at $\pi(z)$. Denote

$$H_{z_0} \equiv \left\{ X \in \mathbb{C}^n \mid \langle \partial r(z_0), X \rangle \equiv \sum_{i=1}^n \frac{\partial r}{\partial z_i}(z_0) X_i = 0 \right\}.$$

We call $z_0 \in \partial\Omega$ a Levi-pseudoconvex point if

$$\sum_{i,j=1}^n \frac{\partial^2 r(z_0)}{\partial z_i \partial \bar{z}_j} X_i \bar{X}_j \geq 0 \quad \text{for all } X \in H_{z_0}.$$

Received by the editors March 24, 1993.

1991 *Mathematics Subject Classification.* Primary 32H15.

Key words and phrases. Kobayashi metric, pseudoconvex domain.

In [K], Krantz has proven that if Ω is a bounded domain with smooth boundary near z_0 , then there exists constant $C > 0$ such that

$$F_{\Omega}(z, \nabla r(\pi(z))) \geq C \frac{|\nabla r(\pi(z))|}{d^{3/4}(z)} \quad \text{for } z \in \Omega \text{ near } z_0.$$

In this note, we shall prove the following theorems.

Theorem A. Let $\Omega \subset \subset \mathbb{C}^n$ be a bounded domain with smooth boundary near $z_0 \in \partial\Omega$. Suppose that there exist $\alpha > 3/4$, $C > 0$, $X \in \mathbb{C}^n \setminus H_{z_0}$, and $\{z_k\}_{k=1}^{\infty}$ with $z_k \rightarrow z_0$ nontangentially (i.e., z_k stays in some cone Λ with vertex at z_0 and axis N_{z_0}) such that

$$(1.1) \quad F_{\Omega}(z_k, X) \geq C \frac{|\langle \partial r(z_0), X \rangle|}{d^{\alpha}(z_k)} \quad \text{for all } k \in \mathbb{N}.$$

Then z_0 is a Levi-pseudoconvex point.

Theorem B. Let $\Omega \subset \subset \mathbb{C}^n$ be a bounded domain with smooth boundary near $z_0 \in \partial\Omega$, and let Λ be a cone with vertex at z_0 and axis N_{z_0} . Assume that z_0 is the origin. Suppose that the local defining function of Ω has the following form near z_0 :

$$r(z) = \operatorname{Re} z_n + \mathcal{O}(|z'|^m + |z_n| \cdot |z|).$$

Then there exist a neighborhood V of z_0 and a constant $C > 0$ such that

$$(1.2) \quad F_{\Omega}(z, X) \geq C \frac{|X_n|}{d^{1-\frac{1}{m}}(z)}$$

for $z \in \Lambda \cap \Omega \cap V$ and all $X \in \mathbb{C}^n$. Furthermore, there exists $C_1 > 0$ such that

$$(1.3) \quad F_{\Omega}(z, X) \geq C_1 \frac{|X_n|}{d^{1-\frac{1}{2m}}(z)}$$

for $z \in \Lambda \cap \Omega \cap V$ and $X \in \mathbb{C}^n$ with $|X| \leq K|X_n|$ (C_1 may depend on the constant K).

As corollaries of Theorem A and Theorem B, we obtain

Corollary 1. Let $\Omega \subset \subset \mathbb{C}^n$ be a bounded domain with smooth boundary near $z_0 \in \partial\Omega$, and let Λ be a cone with vertex at z_0 and axis N_{z_0} . Then z_0 is a Levi-pseudoconvex point if and only if there exist a neighborhood V of z_0 , $\alpha > 3/4$, and $C > 0$ such that

$$(1.4) \quad F_{\Omega}(z, \nabla r(z_0)) \geq C \frac{|\nabla r(z_0)|}{d^{\alpha}(z)}$$

for all $z \in \Lambda \cap \Omega \cap V$.

Corollary 2. Let $\Omega \subset \subset \mathbb{C}^2$ be a bounded domain with smooth pseudoconvex boundary near $z_0 \in \partial\Omega$ and let Λ be a cone with vertex at z_0 and axis N_{z_0} . Then for each $\alpha \in (0, 1)$, there exist a neighborhood V of z_0 and a constant $C > 0$ such that

$$(1.5) \quad F_{\Omega}(z, X) \geq C \frac{|\langle \partial r(z_0), X \rangle|}{d^{\alpha}(z)}$$

for all $z \in \Lambda \cap \Omega \cap V$ and $X \in \mathbb{C}^n$.

II. PROOFS OF THE THEOREMS

Some ideas in the proofs of Theorem A and Theorem B come from [K]. We will also need the following lemma, which can be proven directly from Lemma 2 in [R].

Lemma. *Let Ω' be a subdomain of a bounded domain Ω with $\partial\Omega' \cap \partial\Omega \supset U \cap \partial\Omega$ for some neighborhood U of $z_0 \in \partial\Omega$. Then there exist a neighborhood $V \subset\subset U$ of z_0 and a constant $C > 0$ such that*

$$(2.1) \quad F_{\Omega'}(z, X) \leq CF_{\Omega}(z, X)$$

for $z \in \Omega' \cap V$ and $X \in \mathbb{C}^n$.

We will use C to denote constants which may be different in different appearances.

Proof of Theorem A. After a translation and a unitary transformation, we may assume that z_0 is the origin and $\partial\Omega$ is locally defined by

$$(2.2) \quad r(z) = \operatorname{Re} z_n + \sum_{i,j=1}^n a_{ij} z_i \bar{z}_j + \mathcal{O}(|z|^3)$$

for z near z_0 .

Suppose that $\partial\Omega$ is not Levi-pseudoconvex at z_0 . Then the matrix

$$\left(\frac{\partial^2 r(z_0)}{\partial z_i \partial \bar{z}_j} \right)_{1 \leq i, j \leq n-1}$$

has at least one negative eigenvalue. Therefore, after a unitary transformation in $z' = (z_1, z_2, \dots, z_{n-1})$ and a simple change of coordinate system, we may assume that

$$r(z) = \operatorname{Re} z_n - |z_1|^2 + \sum_{i,j=2}^n a_{ij} z_i \bar{z}_j + \mathcal{O}(|z|^3)$$

for $z \in U$, where U is some neighborhood of z_0 . Shrinking U , we have

$$(2.3) \quad r(z) \leq \operatorname{Re} z_n - \frac{|z_1|^2}{2} + C \sum_{i=2}^n |z_i|^2, \quad \text{for } z \in U.$$

Let $\Lambda = \{-\operatorname{Re} z_n > k|z|\}$ ($0 < k < 1$) be the cone. By the Implicit Function Theorem,

$$(2.4) \quad \lim_{\substack{z \rightarrow 0 \\ z \in \Lambda \cap \Omega}} \frac{-\operatorname{Re} z_n}{d(z)} = 1.$$

By the homogeneity of the Kobayashi metric, we may assume that $X = (X_1, X_2, \dots, X_{n-1}, 1)$. For $z = (z', z_n) = (z_1, z_2, \dots, z_n) \in \Lambda \cap \Omega \cap U$, let $\delta = -\operatorname{Re} z_n$. Define $\Phi_{\delta}(\zeta) = (\Phi_{1\delta}(\zeta), \Phi_{2\delta}(\zeta), \dots, \Phi_{n\delta}(\zeta))$ by

$$\begin{aligned} \Phi_{1\delta} &= z_1 + \frac{\delta^{3/4}}{2} X_1 \zeta + 2\zeta^2; \\ \Phi_{k\delta} &= z_k + \frac{\delta^{3/4}}{2} X_k \zeta, \quad \text{for } 2 \leq k \leq n-1; \\ \Phi_{n\delta} &= z_n + \frac{\delta^{3/4}}{2} \zeta. \end{aligned}$$

Then $\Phi_{\delta}(0) = z$, $\Phi'_{\delta}(0) = \frac{\delta^{3/4}}{2} X$.

Claim. *There exists $\gamma \in (0, 1)$ such that for all $\delta > 0$ sufficiently small, we have $\Phi_\delta(\Delta_\gamma) \subset \Omega \cap U$.*

Proof of the Claim. By choosing $\gamma \in (0, 1)$ small enough, we have $\Phi_\delta(\Delta_\gamma) \subset U$. It follows from (2.3) that

$$r(\Phi_\delta(\zeta)) \leq -\delta + \frac{\delta^{3/4}}{2} \operatorname{Re} \zeta - \frac{|\Phi_{1\delta}(\zeta)|^2}{2} + C \sum_{i=2}^n |\Phi_{i\delta}(\zeta)|^2.$$

Since $\delta > k|z|$, we see that when δ is sufficiently small,

$$(2.5) \quad r(\Phi_\delta(\zeta)) < -\frac{3\delta}{4} + \frac{\delta^{3/4}}{2} \operatorname{Re} \zeta - \frac{|\Phi_{1\delta}(\zeta)|^2}{2}.$$

For $|\zeta| < \delta^{1/4}$, by (2.5),

$$\begin{aligned} r(\Phi_\delta(\zeta)) &< -\frac{3\delta}{4} + \frac{\delta^{3/4}}{2} \cdot \delta^{1/4} - \frac{|\Phi_{1\delta}(\zeta)|^2}{2} \\ &= -\frac{\delta}{4} - \frac{|\Phi_{1\delta}(\zeta)|^2}{2} < 0. \end{aligned}$$

For $|\zeta| \geq \delta^{1/4}$, we have

$$\begin{aligned} |\Phi_{1\delta}(\zeta)|^2 &\geq 2|\zeta|^4 - |z_1 + \frac{\delta^{3/4}}{2} X_1 \zeta|^2 \\ &\geq 2|\zeta|^4 - C\delta^{3/2}. \end{aligned}$$

Thus (2.5) implies that when δ sufficiently small,

$$\begin{aligned} r(\Phi_\delta(\zeta)) &< -\frac{3\delta}{4} + \frac{\delta^{3/4}}{2} \operatorname{Re} \zeta - |\zeta|^4 + \frac{C\delta^{3/2}}{2} \\ &\leq -\frac{3\delta}{4} + \frac{C\delta^{3/2}}{2} + \frac{|\zeta|^4}{2} - |\zeta|^4 < 0. \end{aligned}$$

This concludes the proof of the Claim.

Next, by the Claim and the definition of the Kobayashi metric,

$$F_{\Omega \cap U}(z, X) \leq \frac{C}{\delta^{3/4}}.$$

Thus, combining (2.4) and the length-decreasing property of Kobayashi metric, we obtain

$$F_\Omega(z, X) \leq \frac{C}{d^{3/4}(z)},$$

which contradicts (1.1). Therefore, $\partial\Omega$ is Levi-pseudoconvex at z_0 . $\square$

Proof of Theorem B. By the assumption, there exists a neighborhood U of z_0 such that

$$(2.6) \quad \Omega \cap U \subset \{z \in U \mid \operatorname{Re} z_n - C(|z'|^m + |z_n| \cdot |z|) < 0\}.$$

Let $\Lambda = \{-\operatorname{Re} z_n > k|z|\}$ ($k \in (0, 1)$). For $z \in \Lambda \cap \Omega \cap U$ and $X \in \mathbb{C}^n$ (by homogeneity of the Kobayashi metric, we may assume that $|X| \leq 1$), let

$\Phi(\zeta) = (\tilde{\Phi}(\zeta), \Phi_n(\zeta)) = (\Phi_1(\zeta), \Phi_2(\zeta), \dots, \Phi_n(\zeta)) : \Delta \rightarrow \Omega \cap U$ be an analytic disc satisfying

$$(2.7) \quad \Phi(0) = z, \quad \Phi'(0) = \lambda X,$$

where $\lambda > 0$ is a constant to be estimated. By the Cauchy Integral Formula, we have

$$(2.8) \quad |\Phi_i(\zeta) - z_i| \leq C|\zeta|, \quad 1 \leq i \leq n,$$

and

$$(2.8') \quad |\Phi_i(\zeta) - z_i - \lambda X_i \zeta| \leq C|\zeta|^2, \quad 1 \leq i \leq n,$$

for $|\zeta| < 1/2$. Also, by (2.6), we have

$$(2.9) \quad \operatorname{Re} \Phi_n(\zeta) < C \left(|\tilde{\Phi}(\zeta)|^m + |\Phi_n(\zeta)| \cdot |\Phi(\zeta)| \right).$$

Denote $\delta = -\operatorname{Re} z_n$. By (2.8), (2.8'), and the fact that $k|z| < \delta$, we have

$$(2.10) \quad \begin{aligned} |\Phi(\zeta)| &\leq C(|z| + |\zeta|) \\ &\leq C \left((1/k)\delta + c\delta^{1/m} \right) \\ &\leq c^{1/2}\delta^{1/m}, \quad \text{for } |\zeta| < c\delta^{1/m} \end{aligned}$$

and

$$(2.10') \quad \begin{aligned} |\Phi(\zeta)| &\leq C \left(|z| + (\lambda|X|)|\zeta| + |\zeta|^2 \right) \\ &\leq C \left((1/k)\delta + (\lambda|X|)c\delta^{1/2m} + c^2\delta^{1/m} \right) \\ &\leq c^{1/2} \left(\delta^{1/m} + (\lambda|X|)\delta^{1/2m} \right), \quad \text{for } |\zeta| < c\delta^{1/2m} \end{aligned}$$

when c, δ are sufficiently small.

Now, it follows from (2.10), (2.10'), and (2.9) that

$$(2.11) \quad \operatorname{Re} \Phi_n(\zeta) < \frac{\delta}{2} + \frac{1}{2}|\Phi_n(\zeta)|, \quad \text{for } |\zeta| < c\delta^{1/m}$$

and

$$(2.11') \quad \operatorname{Re} \Phi_n(\zeta) < \frac{\delta + (\lambda|X|)^m \delta^{1/2}}{2} + \frac{1}{2}|\Phi_n(\zeta)|, \quad \text{for } |\zeta| < c\delta^{1/2m}$$

when c, δ are sufficiently small.

Denote

$$D_{i\delta} = \left\{ w \in \mathbb{C} \mid \operatorname{Re} w < \frac{\delta + \varepsilon_i(\lambda|X|)^m \delta^{1/2}}{2} + \frac{1}{2}|w| \right\}, \quad \text{for } i = 1, 2$$

where $\varepsilon_1 = 0, \varepsilon_2 = 1$. Let $g_1(\zeta) \equiv \Phi_n(c\delta^{1/m}\zeta)$ and $g_2(\zeta) \equiv \Phi_n(c\delta^{1/2m}\zeta)$. By (2.7), (2.11), and (2.11'), we have $g_i(\Delta) \subset D_{i\delta}$, $g_i(0) = z_n$ ($i = 1, 2$), $g'_1(0) = \lambda X_n c\delta^{1/m}$, and $g'_2(0) = \lambda X_n c\delta^{1/2m}$. However, it is clear that

$$D_{i\delta} \subset \tilde{D}_{i\delta} \equiv \mathbb{C} \setminus \left\{ w \in \mathbb{C} \mid \operatorname{Im} w = 0, \operatorname{Re} w \geq \delta + \varepsilon_i(\lambda|X|)^m \delta^{1/2} \right\}.$$

Thus $g_i(\Delta) \subset \tilde{D}_{i\delta}$.

Since

$$F_{D,\delta}^{\sim}(z_n, 1) \geq \frac{C}{\delta + \varepsilon_i(\lambda|X|)^m \delta^{1/2}}$$

where $C > 0$ is a constant independent on δ , it follows that

$$|g'_i(0)| \leq C(\delta + \varepsilon_i(\lambda|X|)^m \delta^{1/2}).$$

Therefore,

$$(2.12) \quad \lambda|X_n| \delta^{1/m} \leq C\delta$$

and

$$(2.13) \quad \lambda|X_n| \delta^{1/2m} \leq C(\delta + (\lambda|X|)^m \delta^{1/2}).$$

Now by (2.12), $\lambda|X_n| \leq C\delta^{1-1/m}$. Thus (1.2) is valid. Furthermore, when $|X| < K|X_n|$, it follows from (2.13) that $\lambda|X_n| \leq C\delta^{1-1/2m}$ where $C > 0$ may depend on K . Therefore, (1.3) follows. $\square$

Proof of Corollary 1. The sufficiency comes directly from Theorem A. We now prove the necessity. Suppose that z_0 is a Levi-pseudoconvex point. After a change of coordinates, we may assume that z_0 is the origin and $\partial\Omega$ is locally defined by

$$r(z) = \operatorname{Re} z_n + \sum_{i,j=1}^n a_{ij} z_i \bar{z}_j + \mathcal{O}(|z|^3)$$

near z_0 .

Since the matrix $(a_{ij})_{1 \leq i,j \leq n-1}$ is positive semidefinite,

$$\begin{aligned} r(z) &\geq \operatorname{Re} z_n + 2 \operatorname{Re} \left(\sum_{j=1}^{n-1} a_{nj} z_n \bar{z}_j \right) + a_{nn} |z_n|^2 + \mathcal{O}(|z|^3) \\ &= \operatorname{Re} z_n + \mathcal{O}(|z'|^3 + |z_n| \cdot |z|). \end{aligned}$$

Therefore, (1.4) is valid for $\alpha = 5/6$ by Theorem B. $\square$

Proof of Corollary 2. When z_0 is of finite type, as in Property 2.1 in [M], we can construct a smoothly bounded pseudoconvex domain $\Omega' \subset \Omega$ such that $\partial\Omega' \cap \partial\Omega \supset \partial\Omega \cap U$ for some neighborhood U of z_0 . By the Lemma, there exist a neighborhood V of z_0 and a constant $C > 0$ such that

$$F_{\Omega}(z, X) \geq CF_{\Omega'}(z, X)$$

for $z \in V \cap \Omega'$, $X \in \mathbb{C}^n$. Applying Theorem 1 in [C] to Ω' , we then obtain that (1.5) is valid for $\alpha = 1$.

When z_0 is of infinite type, then for each $m \in \mathbb{N}$, after a change of coordinates, the local defining function of Ω near z_0 has the form

$$r(z) = \operatorname{Re} z_2 + \mathcal{O}(|z_1|^m + |z_2| \cdot |z|).$$

Thus, (1.5) follows from Theorem B. $\square$

Remark. The estimates in Theorem B are sharp. Suppose that Ω is a bounded domain such that $\partial\Omega$ is locally defined by $r(z) = \operatorname{Re} z_2 - |z_1|^{2m}$ near the origin. Then for $P_{\delta} = (0, -\delta)$, $X = (\delta^{-1+1/2m}, 1)$, and $Y = (0, 1)$, we have

$$F_{\Omega}(P_{\delta}, X) \approx \frac{1}{\delta^{1-1/2m}}, \quad F_{\Omega}(P_{\delta}, Y) \approx \frac{1}{\delta^{1-1/4m}}.$$

ACKNOWLEDGMENT

The author thanks Professor Steven Krantz for his advice and encouragement.

REFERENCES

- [C] D. Catlin, *Estimates of invariant metrics on pseudoconvex domains of dimension two*, Math. Z. **200** (1989), 429–466.
- [K] S. Krantz, *The boundary behavior of the Kobayashi metric*, Rocky Mountain J. Math. **22** (1992).
- [M] J. McNeal, *Lower bounds on Bergman metric near a point of finite type*, Ann. of Math. (2) **136** (1992), 339–360.
- [R] H. Royden, *Remarks on the Kobayashi metric*, Several Complex Variables II, Lecture Notes in Math., vol 185, Springer, New York, 1971, pp. 125–137.

DEPARTMENT OF MATHEMATICS, WASHINGTON UNIVERSITY, ST. LOUIS, MISSOURI 63130
E-mail address: `sfu@math.wustl.edu`